REVIEW

The long (G_{sa-L}) and short (G_{sa-S}) variants of the stimulatory guanine nucleotide-binding protein. Do they behave in an identical way?

J Novotny and P Svoboda

Institute of Physiology, Czech Academy of Sciences, Videnska 1083, 142 20 Prague 4, Czech Republic
(Requests for offprints should be addressed to P Svoboda)

ABSTRACT

The relative proportions and tissue distribution of the long (G_{sa-L}) and short (G_{sa-S}) variants of the α subunit of the stimulatory G-protein (G_{sa}) change under a wide range of metabolic conditions, such as cellular differentiation, ontogenetic development, ageing and various adaptive processes. Although the two variants of G_{sa} are generally regarded to be functionally identical, this review summarizes recent experimental support for the non-identical behaviour of these proteins. Similarly, there is no consistent evidence for the functional meaning of these changes as far as regulation of adenylate

cyclase activity is concerned. Since it is hard to believe that the complicated scheme of alternative splicing and the energy-consuming synthesis of proteins would be performed for no reason, it is suggested that G_{sa} variants might be involved in controlling other effector molecules and processes besides adenylate cyclase and cAMP metabolism. Such an idea is indirectly supported by specific alterations in the G_{sa-L}/G_{sa-S} ratio under various physiological and pathophysiological conditions. *Journal of Molecular Endocrinology* (1998) **20**, 163–173

INTRODUCTION

Cell responses to a wide variety of extracellular signals are mediated by heterotrimeric guanine nucleotide-binding proteins (G-proteins), which are composed of α , β and γ subunits. The α subunit is responsible for specific interactions with both the receptor and effector molecules and it primarily determines the function of a G-protein (Gilman 1987, Helmreich & Hofmann 1996). In this review we focus our attention on the properties and behaviour of the individual variants (subforms, isoforms, species) of the α subunits of the stimulatory G-protein (G_{sa}), which have been established as the stimulatory regulatory components of adenylate cyclase and may also be involved in activation of dihydropyridine-sensitive voltage-gated Ca^{2+} channels in skeletal muscle and inactivation of cardiac Na^+ channels (Mattera *et al.* 1989, Schubert *et al.* 1989, Birnbaumer *et al.* 1990). As for other G-proteins, the molecular interactions responsible for G_s -modulated transmembrane sig-

nalling are driven by a cycle of guanine nucleotide exchange and hydrolysis (Fig. 1).

Two forms of G_{sa} ubiquitously expressed in various tissues have been identified by their ability to be ADP-ribosylated by cholera toxin (Northup *et al.* 1980) or by immunoblotting with specific antibodies (Mumby *et al.* 1986). These proteins have been reported to migrate in polyacrylamide gels with apparent M_r values of 52 000 and 45 000 by some authors (Jones & Reed 1987, Feldman *et al.* 1990, Sethi *et al.* 1993) or 45 000 and 42 000 by others (Scherer *et al.* 1987, Milligan 1990, McFarlane-Anderson *et al.* 1992, Negishi *et al.* 1992), depending on the materials and experimental conditions used. Both the long and short forms of G_{sa} (G_{sa-L} and G_{sa-S}) have been shown to be produced by alternative splicing of a single pre-mRNA transcript (Robishaw *et al.* 1986, Kozasa *et al.* 1988). The human G_{sa} gene is a split gene composed of 13 exons and 12 introns that span a region of about 20 kb of genomic DNA (Kozasa *et al.* 1988). The long and short forms of G_{sa} differ by 14 or 15 amino

FIGURE 1. A simplified scheme of the G_s regulatory cycle. Activation of G_s is substantiated by exchange of GTP for GDP on α_s . The transition from inactive ($\alpha_{s(GDP)}$) to active ($\alpha_{s*(GTP)}$) conformation is accelerated by interaction of G_s with a cognate ligand-activated receptor, which stimulates GDP release and facilitates GTP binding. Upon activation, the $\alpha_s^*\beta\gamma$ complex dissociates and both free activated G_{sa} and $G_{\beta\gamma}$ can interact with appropriate effectors. Hydrolysis of GTP by G_{sa} deactivates it, increases its affinity for $G_{\beta\gamma}$, and leads to reassociation to give an inactive GDP-bound G_s .

acid residues, which are coded by exon 3 (Fig. 2). Originally it was thought that there were four splicing products (I–IV) of a single G_{sa} gene, resulting from alternative use of exon 3 and of two 3' splice sites of intron 3 (Bray *et al.* 1986), but several additional splicing variants of G_{sa} have since been described (Ishikawa *et al.* 1990, Swaroop *et al.* 1991, Ali *et al.* 1992, Crawford *et al.* 1993, Habecker *et al.* 1993). It is not known if all splicing variants are expressed as mature proteins. Some of these couple receptors to stimulation of adenylate cyclase and Ca^{2+} channels, whereas others encode truncated proteins the functions of which are not currently defined.

Besides typically ubiquitous long and short forms of G_{sa} , there exists a closely related α subunit of the olfactory GTP-binding protein (G_{aolf} , $M_r \sim 42\,000$), which is distributed in vertebrate chemosensory neurons only (Pace & Lancet 1986, Jones & Reed 1987, 1989, Jones *et al.* 1990). Two forms of G_{sa} -like subunit ($M_r \sim 50\,000$ and $45\,000$) have been described in *Drosophila melanogaster*, which function in a manner similar to their mammalian homologues (Quan *et al.* 1991), and another functional G_{sa} -like protein ($M_r \sim 42\,000$) has recently been identified in the free-living protist *Euglena gracilis* (Torresmarquez *et al.* 1996). In

FIGURE 2. Origin of possible subtypes of G_{sa} cDNAs by alternative splicing of 5'-exons (modified from Swaroop *et al.* 1991). The first four G_{sa} cDNA species (I–IV) differ from each other by the presence of an exon 3 and/or a CAG trinucleotide at the boundary of exons 3 and 4. The next eight variants (V–XII) contain novel exons upstream of exon 2 (exon A and/or B) and they may be generated either by alternative splicing or by using an alternative promoter. In addition, aberrant splicing events involving internal deletions at non-consensus sites have been also described (Ali *et al.* 1992), and Crawford *et al.* (1993) reported neural expression of the $G_{sa}N1$ transcript generated by alternative splicing and polyadenylation of a novel terminal exon downstream of exon 3.

addition, a tightly membrane-associated 'extra large' G-protein, XLas ($M_r \sim 94\,000$), has been revealed in the *trans*-Golgi network, which consists of a new 51 kDa XL-portion linked to a G_{sa} truncated at the N-terminus (Kehlenbach *et al.* 1994).

During the last decade, a large amount of information has been gathered about the unequal distribution of G_{sa} variants in various tissues. To give some examples, G_{sa-L} markedly predominates in kidney, placenta, adrenal medulla, cortex and cerebellum (Evans *et al.* 1986, Mumby *et al.* 1986, Granneman & Kapatos 1990, Feinstein *et al.* 1992, Michel *et al.* 1994), whereas G_{sa-S} is the prevailing form in heart, liver, neostriatum and platelets (Mumby *et al.* 1986, Molina y Vedia *et al.* 1989, Cooper *et al.* 1990, Kawai & Arinze 1996). The existence of different forms of G_{sa} with varying tissue distribution invites speculation that they may have evolved to perform different regulatory functions. To date, however, there is only a small

amount of evidence to indicate that the G_{sa} variants have more than marginal differences in properties and function.

G_{sa-L} AND G_{sa-S} ARE FUNCTIONALLY IDENTICAL

Studies with purified G_{sa} produced in bacteria have shown only minor functional differences between G_{sa-L} and G_{sa-S} (Graziano *et al.* 1987, 1989, Mattera *et al.* 1989, Freissmuth 1991). Recombinant G_{sa} proteins interact functionally with $G_{\beta\gamma}$, β -adrenergic receptors, adenylate cyclase and Ca^{2+} channels and they can be equally ADP-ribosylated by cholera toxin. Both forms of recombinant G_{sa} hydrolyse GTP and have essentially the same k_{cat} for GTP hydrolysis (about 4 min^{-1}). The only difference between bacterially expressed G_{sa-L} and G_{sa-S} is a modest difference in the rate of GDP dissociation, otherwise they are biochemically indistinguishable in *in vitro* reconstitution assays. Further experiments revealed that the affinity of recombinant G_{sa} for adenylate cyclase is roughly 10 times lower than that of native G_{sa} purified from the liver, while the intrinsic capacity of the recombinant protein to activate adenylate cyclase is normal (Graziano *et al.* 1989). This can probably be seen as a consequence of the failure of G_{sa} to undergo a post-translational modification that is essential to achieve high-affinity interaction of the G-protein with adenylate cyclase.

Studies using G_{sa} expression in mammalian systems have provided a similar picture. No substantial difference has been found in the ability of G_{sa-L} and G_{sa-S} expressed in S49 cyc^- cells to stimulate adenylate cyclase activity and to couple to β -adrenergic receptors (Jones *et al.* 1990, O'Donnell *et al.* 1991). In another study, the region of variation between the long and short variants of G_{sa} was modified by genetic engineering to produce an epitope-tagged variant of G_{sa-L} (Levis & Bourne 1992). The cellular distribution (membrane-bound vs soluble form) of tagged and untagged G_{sa} constructs expressed in S49 cyc^- cells was identical, suggesting that the epitope does not disturb targeting of the protein to the plasma membrane, and no significant alterations were observed in the method of regulation of adenylate cyclase by these two variants.

G_{sa-L} AND G_{sa-S} ARE NOT FUNCTIONALLY IDENTICAL

The first evidence in favour of the idea that the two variants of G_{sa} are not identical from a functional

point of view was put forward by Sternweis *et al.* (1981), who, by testing partially purified preparations of G_{sa-L} and G_{sa-S} from rabbit liver, found that the larger species has a greater ability to support hormone-stimulated adenylate cyclase activity. Similar results have been reported by Yagami (1995), who studied the interaction of G_{sa} subforms with β -adrenergic and glucagon receptors in liver plasma membranes. It has been shown that activated glucagon receptors, by accelerating the rate of exchange of bound GDP for GTP on G_{sa} , enhance substantially the sensitivity of both G_{sa} variants to tryptic digestion, while activated β -adrenergic receptors have a similar effect on G_{sa-L} only. This observation supports the notion that β -adrenergic receptors are preferentially coupled to the long form of G_{sa} .

In contradiction, a higher capability of the short isoform of G_{sa} to transmit signals has been suggested by Walseth *et al.* (1989). This departure from the commonly held view that the two subunits have similar efficacies in stimulating adenylate cyclase was based on the increase in responsiveness to isoprenaline, cholera toxin and forskolin in master pancreatic islet β cells as a function of passage number. The increased number of passages in the course of cultivation of these cells was accompanied by a decrease in the G_{sa-L}/G_{sa-S} ratio from 1.0 to 0.22, and the extracts derived from later passages reconstituted adenylate cyclase activity in S49 cyc^- membranes 3-4 times more effectively.

The idea of functional 'non-identity' of G_{sa-L} and G_{sa-S} has also been supported by the different behaviour of the two isoforms in the course of subcellular redistribution of the G_{sa} proteins induced by prolonged agonist (isoprenaline) stimulation of S49 lymphoma cells (Kvapil *et al.* 1994). The transfer of G_{sa} functional activity (measured by a cyc^- reconstitution assay) from plasma membranes to low-density membranes (light vesicles) was accompanied by an increase in the G_{sa-L}/G_{sa-S} ratio in the latter membrane fraction. In addition, the light-vesicular fractions enriched in G_{sa-L} proved to be more effective in protecting adenylate cyclase against its thermal inactivation (Kvapil *et al.* 1995a). This observation may indicate different quality of the interaction between adenylate cyclase and the long and short G_{sa} subforms. Similar subcellular redistribution of G_{sa} has been recently described as a consequence of long-term forskolin treatment of astroglial cells (El Jamali *et al.* 1996). In membrane preparations from placental vascular smooth muscle, vasoactive intestinal peptide in the presence of guanosine 5'-[γ -thio]triphosphate (GTP γ S) also triggered a dose-dependent release of both G_{sa} subforms into the cytosolic fraction, while

the amount of G_{sa-S} released appeared to be much higher than that of G_{sa-L} (Bourgeois *et al.* 1996).

An interesting difference in the behaviour of the G_{sa} variants was also observed in experiments performed with reconstituted phospholipid vesicles, where different recoveries (different molar ratios of α to $\beta\gamma$) for G_{sa-L} and G_{sa-S} were determined for different preparations of G-protein $\beta\gamma$ subunits (Rubenstein *et al.* 1991). Two preparations of $\beta\gamma$ subunits differed reproducibly in their ability to efficiently reconstitute G_{sa-S} , while no substantial change was detected with respect to G_{sa-L} . These results indicate for the first time that different combinations of β and γ subtypes can interact unequally with the long and short G_{sa} subforms.

IN VITRO REGULATION OF G_{sa-L} AND G_{sa-S} : STUDIES ON TISSUE CULTURES

Maintained exposure of NG108-15 cells (neuroblastoma–glioma hybrid) to agonists at the prostanoïd IP receptor results in substantial reduction (down-regulation) of total cellular levels of G_{sa} (McKenzie & Milligan 1990, Adie *et al.* 1992). Agonist-induced down-regulation of G_{sa} protein levels is dependent on the level of receptor expression and is generally restricted to the G-protein(s) with which the receptor interacts (Milligan 1993, Adie & Milligan 1994). Alterations in the individual variants of G_{sa} , however, have not been studied in this respect and it remains to be clarified whether G_{sa-L} and G_{sa-S} respond to long-term agonist treatment in the same way.

Foster *et al.* (1990) have studied the effect of K^+ -induced depolarization on G-protein expression in spontaneously contracting neonatal rat myocytes. They observed that within 3 days of cultivation in medium containing high concentrations of KCl, the cells began to accumulate G_{sa-S} , which was virtually not expressed under control conditions. The level of G_{sa-L} in myocyte membranes was not substantially influenced and the changes in G_{sa-S} induced by KCl depolarization appeared to be reversible.

When selective patterns of expression of G-protein α subunits during *in vitro* development of primary hypothalamic neuron-enriched cultures was studied, no G_{sa-S} was detected (Viollet *et al.* 1994). Nevertheless, remarkably dynamic short-term changes in G_{sa} mRNA as well as G_{sa-L} protein levels were demonstrated. While the cellular concentration of G_{sa} mRNA increased by roughly 75% after 5 days and this level was maintained for the next 2 weeks of cultivation, G_{sa-L} immunoreactivity doubled during the first 5 days of

development and then fell sharply. Thus the time-related decrease in G_{sa} protein level was not paralleled by the corresponding mRNA. Similarly, rapid disappearance of G_{sa} proteins has been reported by Cussac *et al.* (1990) in mouse hypothalamus shortly after birth and in fetal hypothalamic cells cultivated *in vitro* (Kitamura *et al.* 1989).

Alterations in the relative amounts of G_{sa} variants have also been observed during maturation of reticulocytes. Although the concentration of both isoforms of G_{sa} decreased with maturation, a greater reduction in G_{sa-L} was detected (Larner & Ross 1981).

When characterizing the potential roles of G_a proteins in differentiation of white adipocytes cultivated in primary culture, Denis-Henriot *et al.* (1996) detected mainly G_{sa-L} in confluent preadipocytes, whereas mature adipocytes expressed both the long and short variants. Differential expression of the G_{sa} isoforms has also been found during differentiation of 3T3-L1 fibroblasts, which can readily differentiate into cells possessing the morphological and biochemical properties of adipocytes (Watkins *et al.* 1982, 1987, 1989). In 3T3-F442 fibroblasts, the levels of G_{sa-L} and G_{sa-S} increased during differentiation by about 125% and 750% respectively (Kilgour & Anderson 1993).

IN VIVO REGULATION OF G_{sa-L} AND G_{sa-S}

Cardiovascular system

Under 'more physiological' conditions, non-identical behaviour (regulation) of the two G_{sa} variants has also been demonstrated. Significant alterations have been determined during ontogenetic development and ageing. The amounts of G_{sa-L} in rabbit heart muscle do not change during the first 6 weeks of life, while those of G_{sa-S} substantially increase during this period of postnatal development (Kawai & Arinze 1996). Interestingly, the G_{sa-L}/G_{sa-S} ratio (about 0.4) in plasma membranes derived from rabbit liver remained unchanged during the first 6 weeks of life, but the total amount of G_{sa} was markedly elevated in 6-week-old animals (Kawai & Arinze 1991).

The opposite type of alteration has been described in rat hearts during ageing. A significant decrease in G_{sa-S} paralleled by an increase in G_{sa-L} was determined by cholera toxin-induced ADP ribosylation in samples prepared from 24-month-old rats, compared with 6-month-old animals (Urasawa *et al.* 1991). Similar but less pronounced age-related changes in the distribution of G_{sa} variants in rat heart have been shown by

immunodetection techniques (Shu & Scarpace 1994). Interestingly, basal, isoprenaline- and forskolin-stimulated adenylate cyclase activity in myocardial membranes decreased during ageing, whereas adenylate cyclase activity stimulated by the non-hydrolysable GTP analogues GTP γ S and guanosine 5'-[β,γ -imido]triphosphate (p[NH]ppG) was significantly potentiated. An age-related decline in G_{sa} gene transcripts, which was associated with down-regulation of all the G_{sa} protein species, has been reported by Miyamoto *et al.* (1994) in rat ventricular myocardium. When studying expression of G-proteins in cardiocytes, Foster *et al.* (1990) observed that neonatal rat cardiac myocytes and non-muscle heart cells contained preferentially G_{sa-L} and almost no G_{sa-S} . In contrast, atrial membranes from adult rat hearts had approximately equal amounts of the two G_{sa} variants, and adult ventricles had predominantly G_{sa-S} .

In the aorta of adult and senescent rats, no significant difference has been detected in the levels of G_{sa} mRNAs (Johnson *et al.* 1995), while an immunoblot analysis using specific G_{sa} -oriented antibodies revealed a significant age-related decrease in the content of the G_{sa-S} protein (but not G_{sa-L}) in 24-month-old rats, compared with 6-month-old animals.

Kidney

Marked alterations in the relative proportion of G_{sa} variants have also been found in rat kidney during ontogenesis. While the levels of G_{sa-L} in membrane preparations prepared from 3-week-old and 28-week-old rats were roughly identical, the content of G_{sa-S} doubled in preparations from older animals (Michel *et al.* 1994). Almost the opposite phenomenon has been observed in age-matched spontaneously hypersensitive rats; G_{sa-S} did not change, whereas G_{sa-L} significantly decreased in membranes derived from 28-week-old rats (Michel *et al.* 1994). The reduced G_{sa-L} content may well be the molecular basis for the previously observed unchanged isoprenaline-stimulated cAMP formation despite the increased β -adrenergic receptor number (Michel *et al.* 1993).

Liver

The effect of partial hepatectomy on the levels of G_{sa} variants in rat liver plasma membranes has been studied by Yagami *et al.* (1994). The amounts of both G_{sa} species increased after hepatectomy, with a maximum at 48-72 h, and subsequently decreased. The increase in G_{sa-L} was more significant than that in G_{sa-S} . The maximal amounts of G_{sa-L} and G_{sa-S}

were 2.4-fold and 1.8-fold higher respectively when compared with samples from animals before sham operation. The abundance of the G_{sa} mRNAs reached a maximum at 24-48 h, and the increase in G_{sa-L} mRNA was higher than that in G_{sa-S} mRNA, which may well reflect the differences in the elevated levels of G_{sa} proteins. Perhaps it is not such a surprise that the time courses of quantitative changes in G_{sa} proteins were accompanied by corresponding changes in catecholamine-responsive adenylate cyclase activity.

When studying the adenylate cyclase signalling system in liver plasma membranes from lean and genetically diabetic (*db/db*) mice, Palmer & Houslay (1991) noted that treatment of animals with pertussis toxin significantly influenced levels of G_{sa} . In membranes prepared from pertussis toxin-treated lean rats, the expression of G_{sa-L} and G_{sa-S} was increased 2.5-fold and 2.9-fold respectively, whereas the levels of these variants in membranes from diabetic animals were reduced by 20% and 15% respectively. A similar enhancement of the ability of glucagon, isoprenaline and p[NH]ppG to activate adenylate cyclase was observed in samples from both lean and diabetic rats. It therefore seems likely that *in vivo* treatment of rats with pertussis toxin elicits alterations extending far beyond the simple modification of G_{ia} by ADP-ribosylation. Nevertheless, the reason for its opposing effects on the expression of G_{sa} variants in lean and diabetic animals is unclear.

Brain

The ratios of G_{sa} variants in mouse brain change dramatically during development. The G_{sa-L} content was found to be more than twice that of the adult in the embryo, whereas G_{sa-S} represented no more than 10% of the level determined in adult animals (Rius *et al.* 1991). The amount of G_{sa-L} decreased as G_{sa-S} increased during the postnatal period and at day 14 the concentrations of the long and short G_{sa} isoforms in whole brain membranes were nearly equal. These changes were accompanied by a significantly enhanced sensitivity of adenylate cyclase to stimulation by p[NH]ppG. The reciprocal relationship between the two isoforms of G_{sa} suggests that the alternative splicing mechanism of G_{sa} might be involved in ontogenetic development of brain tissue.

In rat telencephalon, a gradually increasing concentration of G_{sa-L} has been detected during the first few days after birth by Kitamura *et al.* (1989). After reaching its peak level at day 12, G_{sa} decreased to the fetal level, which was then maintained at the adult stage.

Differential expression of G_{sa} variants has also been described in the cerebellum and neostriatum, in which G_{sa-L} (cerebellum) and G_{sa-S} (neostriatum) are predominantly expressed (Cooper *et al.* 1990). Since neostriatum is far less rich in Ca^{2+} /calmodulin-regulated adenylate cyclase than most brain regions (including cerebellum), it can be hypothesized that preferential expression of G_{sa-S} might be selectively associated with Ca^{2+} /calmodulin-independent types of adenylate cyclase.

Ozawa *et al.* (1993) have found a reduction in the concentration of G_{sa-L} but not G_{sa-S} in synaptic membranes derived from the temporal cortex of alcoholics. A decrease in G_{sa} in anterior pituitary and cerebellar membranes has also been reported after chronic ethanol administration to long-sleep mice (Wand & Levine 1991). Immunoblotting techniques have revealed different distributions of G_{sa} variants in post-mortem samples of frontal cortex of suicide victims (Cowburn *et al.* 1994). The G_{sa-S} content showed a tendency to be increased in the both violent death and depressed suicide subgroups. The levels of G_{sa-L} exhibited a significant positive correlation (increase) with the age of the experimental subject. This observation is in contrast with that reported by Young *et al.* (1991a) for human parietal cortex, where a strikingly higher ratio of G_{sa-L}/G_{sa-S} was found in infants than in adults. This resulted from lower amounts of G_{sa-L} together with much higher concentrations of G_{sa-S} in adulthood. Patients with bipolar affective disorder exhibited an increased G_{sa} -related immunoreactivity in cerebral cortex, which was mainly due to an increased concentration of G_{sa-L} compared with that in age- and sex-matched controls (Young *et al.* 1991b).

In brain membranes of genetically obese (*ob/ob*) mice, the total amount of immunochemically detectable G_{sa} was increased and the G_{sa-L}/G_{sa-S} ratio shifted from 1.64 to 2.8, compared with lean controls (McFarlane-Anderson *et al.* 1992). Interestingly, liver membranes from the same obese mice contained roughly half the amount of the G_{sa} determined in lean controls, and the G_{sa-L}/G_{sa-S} ratio remained almost unchanged (McFarlane-Anderson *et al.* 1992).

Myometrium

Different patterns of expression of G_{sa} mRNA splice variants as well as G_{sa} proteins have been observed in human myometrium during gestation (Europe-Finner *et al.* 1993, 1994, 1996). The G_{sa} levels were considerably higher in myometrium taken from pregnant than from non-pregnant women, while the increase in G_{sa-L} was more

pronounced than that in G_{sa-S} . Interestingly enough, the amounts of both G_{sa} species returned to control levels at the spontaneous onset of term or preterm labour.

White and brown adipose tissue

Specific age-related changes in the long and short isoforms of G_{sa} have been found in epididymal white adipose tissue. Adipocyte membranes isolated from 18-month-old rats exhibited a 2-fold increase in G_{sa-S} as compared with 9-week-old animals (Green & Johnson 1989). No change was detected in the content of G_{sa-L} . Similar results were obtained by both immunoblotting and cholera toxin-induced ADP-ribosylation.

Quantification of G_{sa} variants by immunodetection in white adipose tissue of genetically diabetic (*db/db*) mice revealed that the long isoform was significantly less abundant than in non-diabetic controls, whereas no difference was detected in the amount of the short isoform (Begin-Heick 1992, 1996). A similar result was obtained for genetically obese (*ob/ob*) mice as compared with lean littermates (Begin-Heick 1990). An identical reduction in G_{sa-L} and G_{sa-S} levels was found by Strassheim *et al.* (1991) in the adipocytes of obese (*fa/fa*) Zucker rats. This reduction in G_{sa} proteins was accompanied by attenuated activation of adenylate cyclase by stimulatory ligands.

Another type of evidence in favour of differential regulation of the steady-state levels of the short and long variants of G_{sa} was obtained in studies of brown adipose tissue. The perinatal stimulation (recruitment) of brown fat was associated with changes in the splicing pattern of G_{sa} mRNA, and these changes were reflected in G_{sa} protein expression (Granneman *et al.* 1990). G_{sa-L} mRNA increased significantly without any effect on the level of G_{sa-L} protein, whereas G_{sa-S} mRNA did not change and G_{sa-S} protein concentration declined to 40% of the control level. The increase in the G_{sa-L}/G_{sa-S} ratio was therefore achieved by the preferential decrease in G_{sa-S} , with little or no change in G_{sa-L} protein (Chaudhry & Granneman 1991).

Granneman & Bannon (1989) and Granneman *et al.* (1990) also suggested that the increased G_{sa} mRNA levels that occur in brown adipose tissue during periods of stimulation serve to maintain membrane levels of G_{sa} protein. The reason why this is needed may well involve the agonist-induced down-regulation of the G_{sa} subunits, which proceeds primarily as increased proteolytic degradation of this protein (McKenzie & Milligan 1990, Milligan 1993, Mitchell *et al.* 1993).

A similar change in the G_{sa-L}/G_{sa-S} ratio has been found in another 'recruited' state of brown adipose tissue metabolism, the one induced by long-term adaptation to cold (cold acclimation). Cold acclimation was found to be associated with a preferential decrease in G_{sa-S} in brown adipose tissue membranes, whereas no change was detected in the amount of G_{sa-L} (Kvapil *et al.* 1995b). This result was based on resolution of plasma membrane proteins by standard SDS-PAGE and immunoblot analysis with antiserum raised against an internal sequence (amino acids 326-335) of G_{sa} . When another type of antiserum (raised against the C-terminal decapeptide of G_{sa}) and high-resolution urea-SDS-PAGE was used, the results were very similar: cold acclimation was associated with an increase in the G_{sa-L}/G_{sa-S} ratio from 2.39 to 3.03 as compared with controls (P Svoboda, L Bourova & J Novotny, unpublished results).

Thus the two types of recruitment of brown adipose tissue, i.e. cold acclimation and perinatal cold stress, seem to be associated with similar changes in the G_{sa-L}/G_{sa-S} ratio. The increase in the G_{sa-L}/G_{sa-S} ratio induced by perinatal cold stress, however, is accompanied by an increase in noradrenaline- and fluoride-stimulated adenylate cyclase (Chaudhry & Granneman 1991) while the isoprenaline-, fluoride- and GTP γ S-stimulated adenylate cyclase activity in plasma membranes isolated from cold-acclimated animals is decreased; the same applies to the functional activity of G_{sa} assessed by a cyc^- reconstitution assay (Svoboda *et al.* 1993). The reason for this discrepancy is not known.

CONCLUDING REMARKS

Until recently there was little evidence that the splice variations in G_{sa} have more than marginal significance with respect to their properties and function (Graziano *et al.* 1987, 1989, Mattera *et al.* 1989, O'Donnell *et al.* 1991, Jones *et al.* 1990, Freissmuth *et al.* 1991). In addition, the region of variation between the long and short variants of G_{sa} has been modified by genetic engineering to produce an epitope-tagged variant of G_{sa} which behaves in a similar manner to the wild-type protein (Levis & Bourne 1992).

It is hard to believe, however, that the complicated scheme of alternative splicing and energy-consuming synthesis of proteins from corresponding transcripts would proceed without any functional use. As summarized in this review, abundant data in the literature show that the steady-state mRNA and protein levels of the long and short variants of

G_{sa} change dramatically during ontogenetic development, ageing, cellular differentiation, gestation, cold acclimation and pathophysiological states such as obesity, hypertension, diabetes and alcoholism. This descriptive evidence, although indirect and circumstantial, supports the idea that the expression of the two G_{sa} variants is regulated according to the functional requirements of a given cell or tissue.

Similarly, there is no consistent evidence that the changes in the relative proportion of G_{sa-L} and G_{sa-S} regulate effector activity. The increase in the G_{sa-L}/G_{sa-S} ratio was found to be associated with an increase (Granneman *et al.* 1990, Chaudry & Granneman 1991, Urasawa *et al.* 1991) as well as a decrease in adenylate cyclase activity (Svoboda *et al.* 1993, Kvapil *et al.* 1995b), and the decrease in the G_{sa-L}/G_{sa-S} ratio was accompanied by an increase (Rius *et al.* 1991, Kilgour & Anderson 1993), no change (Michel *et al.* 1994) or a decrease (Green & Johnson 1989, Begin-Heick 1992) in the enzyme activity. More specifically, a dramatic increase in G_{sa-S} , accompanied by little or no change in G_{sa-L} , was found to proceed together with an increase (Walseth *et al.* 1989, Rius *et al.* 1991, Kilgour & Anderson 1993), no change (Michel *et al.* 1994) or a decrease (Green & Johnson 1989) in adenylate cyclase activity. On the one hand this is not surprising, because the final effect of G_{sa} proteins on adenylate cyclase is modulated by other constituents of a complex G-protein-regulated signalling pathway (such as G_{ia} and $G_{\beta\gamma}$, to name just two) but, on the other hand, the lack of correlation between the G_{sa-L}/G_{sa-S} ratio and adenylate cyclase activity might indicate that the functional meaning of alternative splicing of pre-mRNA is not just regulation of coupling *per se* but regulation of some other process(es) not directly involved in the receptor-G-protein-adenylate cyclase-cAMP cascade.

According to such an alternative view, the differential expression of G_{sa-L} and G_{sa-S} would be part of some physiological or pathophysiological 'programme' (differentiation, recruitment, obesity, etc.). Interesting parallels supporting this view may be found in brown and white adipose tissue. In brown adipose tissue from cold-acclimated animals (Svoboda *et al.* 1993, 1996a, Kvapil *et al.* 1995b), the ratio between the long and short splice variants of G_{sa} protein is increased. During postnatal development of brown adipose tissue, a change in the proportion between these isoforms has also been observed, with a higher G_{sa-L}/G_{sa-S} ratio being found in 1-week-old pups (high degree of recruitment) and a lower ratio in 1-month-old pups (lower degree of recruitment) (Granneman *et al.* 1990,

Chaudry & Granneman 1991). If the postnatal change is understood as being a response to the environmental conditions (Nedergaard *et al.* 1986, Obregon *et al.* 1989) rather than being an ontogenetic change after birth, then both the perinatal changes and the changes during cold acclimation follow a new general pattern of decreased relative levels of G_{sa-S} vs G_{sa-L} in functionally recruited states of brown adipose tissue.

Interesting parallels can be found in white adipose tissue. During differentiation of white adipocytes in culture, the G_{sa-L}/G_{sa-S} ratio is decreased (Dennis-Henriot *et al.* 1996). Similarly, Green & Johnson (1989) found a higher G_{sa-L}/G_{sa-S} ratio in white adipocytes from young rats than in those from aged rats, and a higher G_{sa-L}/G_{sa-S} ratio was observed in controls than in *ob/ob* and *db/db* mice (Begin-Heick 1990, 1992, 1996). Thus, in adipose tissue, a higher degree of 'recruitment' is probably in general associated with a higher G_{sa-L}/G_{sa-S} ratio.

A more provocative view is that the changes in G-protein levels during recruitment and/or differentiation should not be seen just as passive events but as causative ones. Such an opinion has been recently promoted especially by Malbon and co-workers, who suggested that the balance between G_{sa} and G_{ia} controls adipogenesis independently of adenylate cyclase and cAMP (Wang & Malbon 1996, Malbon 1997). Related or not, in human atrium, cAMP was found to be an unlikely candidate for regulation of splicing events or post-translational modification of G_{sa} (Monteith *et al.* 1995). Therefore it might be of interest for future experiments to study the regulation of G_{sa-L} and G_{sa-S} expression not as a passive event but as a causative factor of physiological functions.

Since different tissues and cells possess different complements of receptor subtypes, adenylate cyclase isoforms and $G_{p\gamma}$ combinations with potentially different properties with respect to their interactions with G_{sa} splice variants, preferential coupling of G_{sa-L} or G_{sa-S} to certain subtypes of individual components (receptors and adenylate cyclase) of the adenylate cyclase signalling system may help to explain the possible variability in G_{sa} splice variant function. Such a hypothesis, however, remains to be addressed in the future, because no data have been published to date.

ACKNOWLEDGEMENTS

This work was supported by grant 71120 from the Czech Academy of Sciences and grant 305/96/0678 from the Grant Agency of the Czech Republic.

REFERENCES

- Adie EJ & Milligan G 1994 Agonist regulation of cellular G_{sa} subunit levels in neuroblastoma x glioma hybrid NG108-15 cells transfected to express different levels of the human β_2 -adrenoceptor. *Biochemical Journal* **300** 709–715.
- Adie EJ, Mullaney I, McKenzie FR & Milligan G 1992 Concurrent down-regulation of IP prostanoid receptors and the alpha-subunit of the stimulatory guanine-nucleotide-binding protein (Gs) during prolonged exposure of neuroblastoma x glioma cells to prostanoid agonists. *Biochemical Journal* **285** 529–536.
- Ali IU, Reinhold W, Salvador C & Aguano S 1992 Aberrant splicing of G_{sa} transcript in transformed human astroglial and glioblastoma cell lines. *Nucleic Acid Research* **20** 4263–4267.
- Begin-Heick N 1990 Quantification of the α and β subunits of the transducing elements (G_s and G_i) of adenylate cyclase in adipocyte membranes from lean and obese (*ob/ob*) mice. *Biochemical Journal* **268** 83–89.
- Begin-Heick N 1992 Alpha-subunits of Gs and Gi in adipocyte plasma membranes of genetically diabetic (*db/db*) mice. *American Journal of Physiology* **263** C121–C129.
- Begin-Heick N 1996 Beta-adrenergic receptors and G-proteins in the *ob/ob* mouse. *International Journal of Obesity* **20** S32–S35.
- Birnbaumer L, Abramowitz J & Brown AM 1990 Receptor-effector coupling by G proteins. *Biochimica et Biophysica Acta* **1031** 163–224.
- Bourgeois C, Duc-Gorain P, Robert B, Mondon F & Ferre F 1996 G protein expression in human fetoplacental vascularization. Functional evidence for G_{sa} and G_{ia} subunits. *Journal of Molecular and Cellular Cardiology* **28** 1009–1021.
- Bray P, Carter A, Simons C, Guo V, Puckett C, Kamholz J, Spiegel A & Nirenberg M 1986 Human cDNA clones for four species of G_{sa} signal transduction protein. *Proceedings of the National Academy of Sciences of the USA* **83** 8893–8897.
- Chaudhry A & Granneman JG 1991 Developmental changes in adenylate cyclase and GTP binding proteins in brown fat. *American Journal of Physiology* **261** R403–R411.
- Cooper D, Boyajian C, Goldsmith P, Unson C & Spiegel A 1990 Differential expression of low molecular weight form of G_s - α in neostriatum and cerebellum: correlation with expression of calmodulin-independent adenylate cyclase. *Brain Research* **523** 143–146.
- Cowburn RC, Marcusson JO, Eriksson A, Wiehager B & O'Neill C 1994 Adenylate cyclase activity and G-protein subunit levels in postmortem frontal cortex of suicide victims. *Brain Research* **633** 297–304.
- Crawford JA, Mutchler KJ, Sullivan BE, Lanigan TM, Clark MS & Russo AF 1993 Neural expression of a novel alternatively spliced and polyadenylated G_{sa} transcript. *Journal of Biological Chemistry* **268** 9879–9885.
- Cussac D, Faivre-Bauman A, Kordon C, Loudes C & Saltarelli D 1990 G protein ontogenesis in the mouse hypothalamus assessed by ADP-ribosylation. *Neuroendocrinology* **52** (Suppl 1) P3-128.
- Denis-Henriot D, de Mazancourt P, Goldsmith PK & Giudicelli Y 1996 G proteins in adipocytes and preadipocytes: characterization, subcellular distribution, and potential roles for Gi2 and/or Gi3 in the control of cell proliferation. *Cellular Signalling* **8** 225–234.
- El Jamali A, Rachdaoui N, Jacquemin C & Correze C 1996 Long-term effect of forskolin on the activation of adenylate cyclase in astrocytes. *Journal of Neurochemistry* **67** 2532–2539.

- Europe-Finner GN, Phaneuf S, Watson SP & Bernal AL 1993 Identification and expression of G-proteins in human myometrium: up-regulation of G_s in pregnancy. *Endocrinology* **132** 2484–2490.
- Europe-Finner GN, Phaneuf S, Tolkovsky AM, Watson SP & Bernal AL 1994 Down-regulation of G_s in human myometrium in term and preterm labor: a mechanism for parturition. *Journal of Clinical Endocrinology and Metabolism* **79** 1835–1839.
- Europe-Finner GN, Phaneuf S, Mardon HJ & Bernal AL 1996 Human myometrial G_s-small (with serine) and G_s-large (with serine) messenger ribonucleic acid splice variants promote the increased expression of 46- and 54-kilodalton G_s protein isoforms in pregnancy and their down-regulation during labor. *Journal of Clinical Endocrinology and Metabolism* **81** 1069–1075.
- Evans T, Brown ML, Fraser ED & Northup JK 1986 Purification of the major GTP-binding proteins from human placental membranes. *Journal of Biological Chemistry* **261** 7052–7059.
- Feinstein DL, Mumby SM & Milner MJ 1992 Characterization of G_{sa} mRNA transcripts in primary cultures of rat brain astrocytes. *Glia* **5** 139–145.
- Feldman AM, Tena RG, Kessler PD, Weisman HF, Schulman SP, Blumenthal RS, Jackson DG & Van Dop C 1990 Diminished β -adrenergic receptor responsiveness and cardiac dilation in hearts of myopathic Syrian hamsters (Bio 53-58) are associated with functional abnormality of the G stimulatory protein. *Circulation* **81** 1341–1352.
- Foster KA, McDermott PJ & Robishaw JD 1990 Expression of G proteins in rat cardiac myocytes: effect of KCl depolarization. *American Journal of Physiology* **259** H432–H441.
- Freissmuth M, Selzer E, Marulo S, Schutz W & Strosberg AD 1991 Expression of two human β -adrenergic receptors in *E. coli*: functional interaction with two forms of the stimulatory G protein. *Journal of Biological Chemistry* **88** 8548–8552.
- Gilman AG 1987 G proteins: transducers of receptor-generated signals. *Annual Reviews in Biochemistry* **56** 615–649.
- Granneman JG & Bannon MJ 1989 Neural control of the α subunit of G_s messenger ribonucleic acid in rat brown adipose tissue. *Endocrinology* **125** 2328–2334.
- Granneman JG & Kapatos G 1990 Developmental expression of G_o in neuronal cultures from rat mesencephalon and hypothalamus. *Journal of Neurochemistry* **54** 1995–2001.
- Granneman JG, Haverstick DM & Chaudry A 1990 Relationship between G_{sa} messenger ribonucleic acid splice variants and the molecular forms of G_{sa} protein in rat brown adipose tissue. *Endocrinology* **127** 1596–1561.
- Graziano MP, Casey PJ & Gilman AG 1987 Expression of cDNAs for G proteins in *Escherichia coli*. Two forms of G_{sa} stimulate adenylate cyclase. *Journal of Biological Chemistry* **262** 11375–11381.
- Graziano MP, Freissmuth M & Gilman AG 1989 Expression of G_{sa} in *Escherichia coli*. Purification and properties of the two forms of the protein. *Journal of Biological Chemistry* **264** 409–418.
- Green A & Johnson JL 1989 Evidence for altered expression of the GTP-binding regulatory proteins, G_s and G_i, in adipocytes of aged rats. *Biochemical Journal* **258** 607–610.
- Habecker BA, Martin JM & Nathanson NM 1993 Isolation and characterization of a novel cDNA which identifies both neural-specific and ubiquitously expressed G_{sa} mRNAs. *Journal of Neurochemistry* **61** 712–717.
- Helmreich EJM & Hofmann K-P 1996 Structure and function of proteins in G-protein-coupled signal transfer. *Biochimica et Biophysica Acta* **1286** 285–322.
- Ishikawa Y, Bianchi C, Nadal-Ginard B & Homcy C 1990 Alternative promoter and 5' exon generate a novel G_{sa} mRNA. *Journal of Biological Chemistry* **265** 8458–8462.
- Johnson MD, Zhou Y, Friedman E & Roberts J 1995 Expression of G protein α subunits in the aging cardiovascular system. *Journal of Gerontology (Biological Sciences)* **50A** B14–B19.
- Jones DT & Reed RR 1987 Molecular cloning of five GTP-binding protein cDNA species from rat olfactory neuroepithelium. *Journal of Biological Chemistry* **262** 14241–14249.
- Jones DT & Reed RR 1989 G_{olf}: an olfactory neuron-specific G protein involved in odorant signal transduction. *Science* **244** 790–795.
- Jones DT, Masters SB, Bourne HR & Reed RR 1990 Biochemical characterization of three stimulatory GTP-binding proteins. The large and small forms of G_s and the olfactory specific G protein, G_{olf}. *Journal of Biological Chemistry* **265** 2671–2676.
- Kawai Y & Arinze IJ 1991 Ontogeny of guanine-nucleotide-binding regulatory proteins in rabbit liver. *Biochemical Journal* **274** 439–444.
- Kawai Y & Arinze IJ 1996 Differential localization and development-dependent expression of G protein subunits, G α and G β , in rabbit heart. *Journal of Molecular and Cellular Cardiology* **28** 1555–1564.
- Kehlenbach RH, Matthey J & Wieland BH 1994 XL α s is a new type of G protein. *Nature* **372** 804–808.
- Kilgour E & Anderson NG 1993 Changes in the expression of guanine nucleotide-binding proteins during differentiation of 3T3-F442A cells in a hormonally defined medium. *FEBS Letters* **328** 271–274.
- Kitamura Y, Mochii M, Kodama R, Agata K, Watanabe K, Eguchi G & Nomura Y 1989 Ontogenesis of α_2 -adrenoceptor coupling with GTP-binding proteins in the rat telencephalon. *Journal of Neurochemistry* **53** 249–257.
- Kozasa T, Itoh H, Tsukamoto T & Kaziro Y 1988 Isolation and characterization of the human G_{sa} gene. *Proceedings of the National Academy of Sciences of the USA* **85** 2081–2085.
- Kvapil P, Novotny J, Svoboda P & Ransnas LA 1994 The short and long forms of the α subunit of the stimulatory guanine-nucleotide-binding protein are unequally redistributed during (-)-isoproterenol mediated desensitization of intact S49 lymphoma cells. *European Journal of Biochemistry* **226** 193–199.
- Kvapil P, Novotny J & Ransnas LA 1995a The splicing variants of the α subunit of the stimulatory G-protein (G_{sa}) differ in their ability to interact with adenyl cyclase in S49 lymphoma cells. *FASEB Journal* **9** A1411.
- Kvapil P, Novotny J & Ransnas LA 1995b Prolonged exposure of hamsters to cold changes the levels of G proteins in brown adipose tissue plasma membranes. *Life Sciences* **57** 311–318.
- Larner AC & Ross EM 1981 Alteration in the protein components of catecholamine-sensitive adenylate cyclase during maturation of rat reticulocytes. *Journal of Biological Chemistry* **256** 9551–9557.
- Levis MJ & Bourne HR 1992 Activation of the α subunit of G_s in intact cells alters its abundance, rate of degradation and membrane avidity. *Journal of Cellular Biology* **119** 1297–1307.
- McFarlane-Anderson N, Bailly J & Begin-Heick NB 1992 Levels of G-proteins in liver and brain of lean and obese (*ob/ob*) mice. *Biochemical Journal* **282** 15–23.
- McKenzie FR & Milligan G 1990 Prostaglandin E₁-mediated, cyclic AMP-independent, down-regulation of G_{sa} in neuroblastoma x glioma hybrid cells. *Journal of Biological Chemistry* **265** 17084–17093.
- Malbon CC 1997 Heterotrimeric G-proteins and development. *Biochemical Pharmacology* **53** 1–4.

- Mattera R, Graziano MP, Yatani A, Zhou Z, Graf R, Codina J, Birnbaumer L, Gilman A & Brown AM 1989 Splice variants of the α subunit of the G protein G_s activate both adenylyl cyclase and calcium channels. *Science* **243** 804–807.
- Michel MC, Siepmann F, Buscher R, Philipp T & Brodde O-E 1993 Ontogenesis of sympathetic responsiveness in spontaneously hypersensitive rats. I. Renal α_1 -, α_2 - and β -adrenergic receptors and their signalling. *Hypertension* **22** 169–177.
- Michel MC, Farke W, Erdbrugger W, Philipp T & Brodde O-E 1994 Ontogenesis of sympathetic responsiveness in spontaneously hypersensitive rats. II. Renal G proteins in male and female rats. *Hypertension* **23** 653–658.
- Milligan G 1990 Immunological probes and the identification of guanine nucleotide-binding proteins. In *G proteins as Mediators of Cellular Signalling Processes*, pp 31–46. Eds MD Houslay & G Milligan. New York: John Wiley & Sons. Ltd.
- Milligan G 1993 Agonist regulation of cellular G protein levels and distribution: mechanisms and functional implications. *Trends in Pharmacological Sciences* **14** 413–418.
- Mitchell FM, Buckley NJ & Milligan G 1993 Enhanced degradation of the phosphoinositidase C-linked guanine-nucleotide-binding protein Gq/G11 α following activation of the human M1 muscarinic acetylcholine receptor expressed in CHO cells. *Biochemical Journal* **293** 495–499.
- Miyamoto A, Kawana S, Kimura H & Ohshika H 1994 Impaired expression of G_{sa} protein mRNA in rat ventricular myocardium with aging. *European Journal of Pharmacology* **266** 147–154.
- Molina y Vedia L, Nolan RD & Lapetina EG 1989 The effect of iloprost on the ADP-ribosylation of G_{sa} (the α subunit of G_s). *Biochemical Journal* **261** 841–845.
- Monteith MS, Wang T & Brown MJ 1995 Differences in transcription and translation of long and short G_{sa} , the stimulatory G-protein, in human atrium. *Clinical Sciences* **89** 487–495.
- Mumby SM, Kahn RA, Manning DR & Gilman AG 1986 Antisera of designed specificity for subunits of guanine-nucleotide-binding regulatory proteins. *Proceedings of the National Academy of Sciences of the USA* **83** 265–269.
- Nedergaard J, Connolly E & Cannon B 1986 Brown adipose tissue in the mammalian neonate. In *Brown Adipose Tissue*, pp 152–213. Eds P Trayhurn & DG Nicholls. London: Edward Arnold, Ltd.
- Negishi M, Hashimoto H & Ichikawa A 1992 Translocation of α subunits of stimulatory guanine nucleotide-binding proteins through stimulation of the prostacyclin receptor in mouse mastocytoma cells. *Journal of Biological Chemistry* **267** 2364–2369.
- Northup JK, Sternweis PC, Smigel MD, Shleifer LS, Ross EM & Gilman AG 1980 Purification of the regulatory component of adenylyl cyclase. *Proceedings of the National Academy of Sciences of the USA* **77** 6516–6520.
- Obregon MJ, Jacobsson A, Kirchgessner T, Schotz MC, Cannon B & Nedergaard J 1989 Postnatal recruitment of brown adipose tissue is induced by the cold stress experienced by the pups. An analysis of mRNA levels for thermogenin and lipoprotein lipase. *Biochemical Journal* **259** 341–346.
- O'Donnell JK, Sweet RW & Stadel JM 1991 Expression and characterization of the long and short splice variants of G_{sa} in S49 cyc⁻ cells. *Molecular Pharmacology* **39** 702–710.
- Ozawa H, Katamura Y, Hatta S, Saito T, Katada T, Gsell W, Froelich L, Takahata N & Riederer P 1993 Alterations of guanine nucleotide-binding proteins in post-mortem human brain in alcoholics. *Brain Research* **620** 174–180.
- Pace U & Lancet D 1986 Olfactory GTP-binding protein: signal-transducing polypeptide of vertebrate chemosensory neurons. *Proceedings of the National Academy of Sciences of the USA* **83** 4947–4951.
- Palmer TM & Houslay MD 1991 Determination of G-protein levels, ADP-ribosylation by cholera and pertussis toxins and the regulation of adenylyl cyclase activity in liver plasma membranes from lean and genetically diabetic (*db/db*) mice. *Biochimica et Biophysica Acta* **1097** 193–204.
- Quan F, Thomas L & Forte M 1991 Drosophila stimulatory G protein α subunit activates mammalian adenylyl cyclase but interacts poorly with mammalian receptors: implications for receptor-G protein interaction. *Proceedings of the National Academy of Sciences of the USA* **88** 1898–1902.
- Rius RA, Streety RA, Loh YP & Klee WA 1991 Developmental expression of G proteins that differentially modulate adenylyl cyclase activity in mouse brain. *FEBS Letters* **288** 51–54.
- Robishaw JD, Russell DV, Harris BA, Smigel MD & Gilman AG 1986 Deduced primary structure of the α subunit of the GTP-binding stimulatory protein of adenylyl cyclase. *Proceedings of the National Academy of Sciences of the USA* **83** 1251–1255.
- Rubenstein RC, Linder ME & Ross EM 1991 Selectivity of the β -adrenergic receptor among G_s , G_i 's, and G_o : assay using recombinant α subunits in reconstituted phospholipid vesicles. *Biochemistry* **30** 10769–10777.
- Scherer NM, Toro M-J, Entman ML & Birnbaumer L 1987 G-protein distribution in canine cardiac sarcoplasmic reticulum and sarcolemma: comparison to rabbit skeletal muscle membranes and to brain and erythrocyte G-proteins. *Archives of Biochemistry and Biophysics* **259** 431–440.
- Schubert B, Van Dongen AMJ, Kirsch GE & Brown AM 1989 β -adrenergic inhibition of cardiac sodium channels by dual G-protein pathways. *Science* **245** 516–519.
- Sethi R, Dhalla KS, Shah KR & Dhalla NS 1993 Characterization of adenylyl cyclase in heart sarcolemma in the absence or presence of alamethicin. *Molecular and Cellular Biochemistry* **119** 185–193.
- Shu Y & Scarpace PJ 1994 Forskolin binding sites and G-protein immunoreactivity in rat hearts during aging. *Journal of Cardiovascular Pharmacology* **23** 188–193.
- Sternweis PC, Northup JK, Smigel MD & Gilman AG 1981 The regulatory component of adenylyl cyclase. Purification and properties. *Journal of Biological Chemistry* **256** 11517–11526.
- Strassheim D, Palmer T, Milligan G & Houslay MD 1991 Alterations in G-protein expression and the hormonal regulation of adenylyl cyclase in the adipocytes of obese (*fa/fa*) Zucker rats. *Biochemical Journal* **276** 197–202.
- Svoboda P, Unnelius L, Cannon B & Nedergaard J 1993 Attenuation of G_{sa} coupling efficiency in brown-adipose tissue plasma membranes from cold-acclimated hamsters. *Biochemical Journal* **295** 655–661.
- Svoboda P, Unnelius L, Dicker A, Cannon B & Nedergaard J 1996a Cold-induced reduction in G_{sa} proteins in brown adipose tissue. Effects of cellular hypersensitization to noradrenaline caused by pertussis-toxin treatment. *Biochemical Journal* **313** 761–768.
- Svoboda P, Kim G-D, Crassie MA, Eidne KA & Milligan G 1996b Thyrotropin-releasing hormone-induced subcellular redistribution and down-regulation of G11 α : analysis of agonist regulation of coexpressed G11 α species variants. *Molecular Pharmacology* **49** 646–655.
- Swaroop A, Agarwal N, Gruen JR, Bick D & Weissman SM 1991 Differential expression of novel G_{sa} signal transduction protein cDNA species. *Nucleic Acid Research* **19** 4725–4729.

- Torresmarquez ME, Maciassilva M & Vegasegura A 1996 Identification of a functional Gs protein in *Euglena gracilis*. *Comparative Biochemistry and Physiology C* **115** 233–237.
- Urasawa K, Murakami T & Yasuda H 1991 Age-related alterations in adenylyl cyclase system of rat hearts. *Japanese Circulation Journal* **55** 676–684.
- Viollet C, Loudes C, Kordon C & Faivre-Bauman A 1994 Selective patterns of expression of G protein α subunits during *in vitro* development of hypothalamic neurons. *Journal of Neurochemistry* **63** 2231–2238.
- Walseth TF, Zhang H-J, Olson LK, Schroeder WA & Robertson RP 1989 Increase in Gs and cyclic AMP generation in HIT cells. Evidence that the 45 kDa α subunit of Gs has greater functional activity than the 52 kDa α subunit. *Journal of Biological Chemistry* **264** 21106–21111.
- Wand GS & Levine MA 1991 Hormonal tolerance to ethanol is associated with decreased expression of the GTP-binding protein, Gs α and adenylylase activity in ethanol-treated LS mice. *Alcohol Clinical and Experimental Research* **15** 705–710.
- Wang H & Malbon CC 1996 The Gs α /G α 2 axis controls adipogenesis independently of adenylyl cyclase. *International Journal of Obesity* **20** (Suppl 3), S26–S31.
- Watkins PA, Moss J, Pekala PH & Lane MD 1982 Effect of differentiation on the adenylylase system of 3T3-C2 and 3T3-L1 cells. *Journal of Biological Chemistry* **257** 14719–14722.
- Watkins PA, Northup JK & Malbon CC 1987 Regulation of G-proteins in differentiation. Altered ratio of α to β -subunits in 3T3-L1 cells. *Journal of Biological Chemistry* **262** 10651–10657.
- Watkins PA, Rapiejko PJ, Ros M, Wang H-Y & Malbon CC 1989 G-protein mRNA levels during adipocyte differentiation. *Biochemical and Biophysical Research Communications* **165** 929–934.
- Yagami T 1995 Differential coupling of glucagon and β -adrenergic receptors with the small and large forms of the stimulatory G protein. *Molecular Pharmacology* **48** 849–854.
- Yagami T, Kirita S, Matsushita A, Kawasaki K & Mizushima Y 1994 Alterations in the stimulatory G protein of the rat liver after partial hepatectomy. *Biochimica et Biophysica Acta* **1220** 81–87.
- Young LT, Warsh JJ, Li PP, Po Siu K, Becker L, Gilbert J, Hornykiewicz O & Kish SJ 1991a Maturation and aging effects on guanine nucleotide binding protein immunoreactivity in human brain. *Developmental Brain Research* **61** 243–248.
- Young LT, Li PP, Kish SJ, Siu KP & Warsh JJ 1991b Postmortem cerebral cortex G α -subunit levels are elevated in bipolar affective disorder. *Brain Research* **553** 323–326.

RECEIVED 28 July 1997