Suppression of P450 aromatase gene expression in sex-reversed males produced by rearing genetically female larvae at a high water temperature during a period of sex differentiation in the Japanese flounder (*Paralichthys olivaceus*)

T Kitano^{1,2}, K Takamune³, T Kobayashi⁴, Y Nagahama⁴ and S-I Abe²

¹Kumamoto Prefectural Fisheries Research Center, Ohyano, Amakusa, Kumamoto, 869–3603, Japan

²Department of Materials and Life Science, Graduate School of Science and Technology,

Kumamoto University, Kumamoto, 860–8555, Japan

³Department of Biological Science, Faculty of Science, Kumamoto University, Kumamoto, 860–8555, Japan

⁴Laboratory of Reproductive Biology, National Institute for Basic Biology, Okazaki, 444–8585, Japan

(Requests for offprints should be addressed to T Kitano, Kumamoto Prefectural Fisheries Research Center, Ohyano, Amakusa, Kumamoto, 869–3603, Japan)

ABSTRACT

The phenotypic sex of many teleost fishes including flounders can be experimentally altered by treating embryos or larvae with varied temperatures or sex-steroid hormones. To analyse the sex determination mechanism, especially the role of cytochrome P450 aromatase (P450arom), an enzyme that catalyses the conversion of androgens to estrogens, in temperature-dependent gonadal sex differentiation in the Japanese flounder, we generated two populations of larvae, both having XX (genetic females) but each growing up to display all phenotypic females or males, by rearing the larvae at normal (18 °C) or high (27 °C) water temperatures from days 30 to 100 after hatching respectively. The larvae (XX) were produced artificially by mating normal females (XX) with gynogenetic diploid males (XX) which had been sex-reversed to phenotypic males by 17 α -methyltestosterone.

To study the role of P450arom in sex determination in the flounder, we first isolated a P450arom cDNA containing the complete open reading frame from the ovary. RT-PCR showed that P450arom

mRNA was highly expressed in the ovary and spleen but weakly in the testis and brain. Semi-quantitative analyses of P450arom mRNA in gonads during sex differentiation showed that there was no difference in the levels of P450arom mRNA between the female and male groups when the gonad was sexually indifferent (day 50 after hatching). However, after the initiation of sex differentiation (day 60), the mRNA levels increased rapidly in the female group, whereas they decreased slightly in the male group. Similarly, estradiol-17 β levels rose remarkably in the female group, yet remained constant in the male group.

These results suggest that induction of sex reversal of genetically female larvae to phenotypic males by rearing them at a high water temperature caused a suppression of P450arom gene expression. Furthermore, we suggest that the maintenance of P450arom mRNA at very low levels is a prerequisite for testicular differentiation, while the increased levels are indispensable for ovarian differentiation.

Journal of Molecular Endocrinology (1999) **23**, 167–176

INTRODUCTION

In vertebrates, including fishes, sex is often determined by genotype under normal circumstances. In some fishes, however, sex determination is greatly influenced by environmental factors

(Conover & Kynard 1981, Rubin 1985, Sullivan & Schultz 1986). For example, the genetic sex determination mechanism in the Japanese flounder (*Paralichthys olivaceus*) is basically XX(female)–XY(male) type (Tabata 1991, Yamamoto *et al.* 1992). The female to male ratio in the Japanese

flounder decreases drastically with exposure of the larvae to higher or lower water temperature than normal (17.5–22.5 °C) during sex differentiation (Yamamoto 1995, 1999). In addition, the phenotypic sex of many teleost fishes (Hunter & Donaldson 1983, Yamazaki 1983) including the flounder (Tanaka 1988, Tabata 1991) can be experimentally altered by treatment of the embryos or larvae with sex-steroid hormones, suggesting an important role of sex-steroid hormones in phenotypic sex differentiation in fishes.

Cytochrome P450 aromatase (P450arom) is a steroidogenic enzyme responsible for conversion of androgens to estrogens. In vertebrates, this enzyme is expressed in various tissues and plays an important physiological function by regulating estrogen biosynthesis (Pasmanik & Callard 1985, Matsumine *et al.* 1986, Sasano *et al.* 1997). Moreover, the estradiol-17 β produced by the action of P450arom is essential for ovarian development (Wallace 1985). Recently, several reports have described the expression of P450arom during sexual differentiation in several species. P450arom mRNA is detected in the gonads of genetic females but not of males during sexual differentiation in the chicken (Yoshida *et al.* 1996, Smith *et al.* 1997). P450arom activity is significantly higher during ovarian differentiation than during testicular differentiation in reptiles and amphibians which exhibit temperature-dependent sex determination (Desvages & Pieau 1992, Desvages *et al.* 1993, Chardard *et al.* 1995). Positive immunoreaction against P450arom appears in differentiating ovaries, but not in differentiating testes in tilapia (*Oreochromis niloticus*) (Chang *et al.* 1997a). However, the role of P450arom in gonadal sex differentiation of fishes remains unclear, particularly in species with temperature-sensitive, phenotypic sexual differentiation.

To address the role of P450arom in gonadal sex differentiation in the Japanese flounder, we cloned a P450arom cDNA and examined mRNA expression in various adult organs. Furthermore, we analysed the expression level of P450arom mRNA in the flounder gonad and concentrations of sex-steroid hormones during sex differentiation in females (XX) reared under normal water temperature and in sex-reversed males (XX) reared under higher water temperature.

MATERIALS AND METHODS

Animals

Japanese flounder (*Paralichthys olivaceus*) were reared at natural sea water temperature (12–28 °C) in our laboratory.

We produced two populations of larvae, both having XX (genetic females), but each growing up to display all phenotypic females or males, making use of meiotic gynogenetic diploid. First, the meiotic gynogenetic diploid was induced by cold shock treatment (1 °C for 30 min, 4 min after insemination) of the eggs which had been inseminated by UV-irradiated (1000 erg/mm², 1 min) sperm. All the resultant larvae were genetically XX. The phenotypic sex was then reversed to male by feeding the larvae from days 30 to 100 after hatching with an artificial diet (Love Larva, Maruha Co., Yamaguchi, Japan) containing 17 α -methyltestosterone (10 μ g/g diet; Nakarai Tesque, Inc., Kyoto, Japan). These sex-reversed phenotypic males of meiotic gynogenetic diploid were then mated with normal females to produce all genetic female broods. These larvae (all XX) were reared at normal (18 °C) or high (27 °C) water temperatures from days 30 to 100 after hatching, the thermo-sensitive period for sex differentiation (Yamamoto 1995, 1999), to produce phenotypic females and males respectively. The percentage of the phenotypic females (males) among the resultant adult fishes reared at 18 °C (27 °C) was 100%, based on the morphological and histological observations of the adult gonads at 10 months of age (Yamamoto 1995).

Ten to twenty larvae or juveniles were sampled every 10 days after hatching. After measurements of the total body lengths, the samples were frozen at –80 °C for quantification of P450arom mRNA and sex-steroid hormones.

Construction of cDNA library

Total RNA was extracted by the guanidinium isothiocyanate–CsCl method (Chirgwin *et al.* 1979) from an ovary containing vitellogenic oocytes in a 2-year-old flounder. Poly(A)⁺ RNA was purified by affinity chromatography on an oligo(dT)-cellulose column using an mRNA purification kit (Pharmacia Biotech, Tokyo, Japan). Double-stranded cDNA was synthesized according to the method of Gubler & Hoffman (1983) with the following modifications. Oligo(dT) primer with XhoI site at the 5'-end was used, and dCTP in the buffer for first-strand synthesis was substituted by 5-methyl dCTP for the protection of cDNA against the following XhoI treatment. After ligation with EcoRI adaptor (Stratagene, La Jolla, CA, USA), the cDNA was treated with XhoI and inserted into the EcoRI–XhoI site of λ ZAP phage vector (Uni-ZAP XR vector; Stratagene). The resulting constructs were packaged in Gigapack III packaging extract (Stratagene).

Isolation of P450arom cDNA

A flounder ovarian cDNA library was screened with an EcoRI-XhoI fragment (1.8 kbp) of tilapia P450arom cDNA (Chang *et al.* 1997b) as a probe which had been labeled with digoxigenin (DIG) by the method of Feinberg & Vogelstein (1983) using a DIG DNA labeling kit (Boehringer Mannheim, Tokyo, Japan). Plaque replicas on nylon membranes (Hybond-N⁺; Amersham, Tokyo, Japan) were hybridized with the probe in DIG Easy Hyb solution (Boehringer Mannheim) at 42 °C for 16 h. The membranes were washed twice with 2 × SSC (0.3 M NaCl, 30 mM sodium citrate)/0.1% SDS at room temperature for 5 min and twice with 0.1 × SSC/0.1% SDS at 68 °C for 15 min. Detection of DIG was performed by immunoreaction with 1:10 000 diluted alkaline phosphatase-conjugated anti-DIG antibody, followed by treatment with 337.5 µg/ml nitroblue tetrazolium salt and 175 µg/ml 5-bromo-4-chloro-3-indolyl phosphate, using a DIG nucleic acid detection kit (Boehringer Mannheim).

DNA sequencing

The pBluescript phagemid containing the desired cDNA was excised from P450arom-positive phages by co-infection of *Escherichia coli* (XL1-Blue MRF') with the positive phage and the helper phage R408 (Stratagene). Sequencing of both strands of the insert cDNA was performed by the dideoxy chain-termination method of Sanger *et al.* (1977) using the Cy5 AutoRead sequencing kit (Pharmacia Biotech) and an ALFexpress DNA sequencer (Pharmacia Biotech).

Amplification of the 5'-end of P450arom cDNA

The 5'-end of P450arom cDNA was amplified by rapid amplification of cDNA ends (RACE) as described by Frohman *et al.* (1988), using a 5'-RACE system (Gibco BRL, Grand Island, NY, USA). First-strand cDNA was synthesized from 0.5 µg ovarian poly(A)⁺ RNA using 200 units of an RnaseH deficient Moloney murine leukaemia virus reverse transcriptase (Superscript II RT; Gibco BRL) and a specific primer for flounder P450arom corresponding to nucleotides 498–517 (5'-ATGATGCCTC TCTCATACAT-3'; Fig. 1) in the reaction buffer (0.4 mM of each dNTP, 50 mM KCl, 2.5 mM MgCl₂, 20 mM Tris-HCl (pH 8.4), 10 mM dithiothreitol). After addition of oligo-dC to the 3'-end of the first-strand cDNA by treatment with terminal

deoxynucleotide transferase, polymerase chain reaction (PCR) was performed with an abridged anchor primer from the kit (5'-GGCCACGCGTCGACTAGTAC-3') and another specific primer for P450arom corresponding to nucleotides 469–488 (5'-GCTCAGTCCCTGCT TGCTCC-3'; Fig. 1) in the PCR mixture (0.2 mM each of dNTP, 50 mM KCl, 1.5 mM MgCl₂, 10 mM Tris-HCl (pH 8.3), 1 unit Taq DNA polymerase (AmpliTaq Gold; Perkin Elmer, Foster City, CA, USA)). The product was then reamplified with the abridged universal amplification primer from the kit (5'-GGCCACGCGTCGACTAGTAC-3') and a third specific primer for P450arom corresponding to nucleotides 407–426 (5'-TGGC TGATGCTCTGCTGAGG-3'; Fig. 1). PCR conditions were as follows: preheating at 95 °C for 10 min, 30 cycles of PCR at 94 °C for 30 s, 59 °C for 30 s, 72 °C for 1 min, and a final extension at 72 °C for 5 min. The resulting PCR product was electrophoresed on a 2% agarose gel, purified and inserted into the TA-vector pCR II (Invitrogen, San Diego, CA, USA). Determination and confirmation of base sequence of the PCR product were carried out by sequencing of four clones derived from the same PCR product.

Semi-quantification of P450arom mRNA

One microgram total RNA extracted from the juvenile gonads at 50, 60, 80 and 100 days after hatching, or various adult tissues was reverse-transcribed in 20 µl of the reaction mixture (1 mM of each dNTP, 50 mM KCl, 5 mM MgCl₂, 10 mM Tris-HCl (pH 8.3), 2.5 µM oligo (dT)₁₆, 1 unit RNase inhibitor, 2.5 units MuLV reverse transcriptase (Perkin Elmer)) at 42 °C for 30 min. Following dilution of the reaction mixture with 80 µl distilled water, 1 µl of the diluted mixture served as a template for a 20 µl PCR reaction containing 50 pmol of specific primers for P450arom and flounder elongation factor-1α (EF-1α). Primer sequences were based on the sequences of the flounder P450arom (Fig. 1) and the flounder EF-1α (DNA Data Bank of Japan, accession no. AB017183). Forward primers for P450arom and EF-1α were 5'-ATCGGATCCCTGCCTG TGAC-3' and 5'-AGTTTCGAGAAAGAAGCTG CC-3' corresponding to nucleotides 73–92 and 4–23 respectively. Reverse primers for P450arom and EF-1α were 5'-TGGCTGATGCTCTGC TGAGG-3' and 5'-ATCCAGAGCATCCAGC AGTG-3' corresponding to nucleotides 407–426 and 562–581 respectively. After preheating of the PCR mixture at 95 °C for 10 min, 25 cycles

CATCACTTGGCCTGGTCGTCAGTTTGTGCAGTCTTGTGCAGGTTGTTTTAAATTCAGTCTTCCCATGGATCGGATCCCTGCTGTG 90
ACCTGGCATGACTCCCGTAGGTTGGGGGCGCACTGGGGACCTGGTCTCCACGTCCCAAGCCACCGCAGTGAGAACACCGGCA 180
LAAMTTPVGLGLGAAALGDLVSTSPNATAVRRTPGFI
TCTCGTGGCTCAAGGACCTTGATCTGCTGGTGTCTGCTGCTGCTGGAGCCACACAGACAGGAACTCTGGCAGTCCAC 270
SVDAASRIPLILLVCLVLLVAWSHTDRRTVPFGFP
CTTCTGTTTGGGTTTGGGCACTTCTCTCATATGTGAGATTCTCTGGACGGGTATAGCCACAGCTGCACTACTACACAGAGGT 360
FCLGLGLPLLSYVRFIWTGIGTACNNYYNKRKY
ATGGAGACATCTGCAGACTTGGATCGATGGAGAGGAGACACTCTCTCAGCAGAGCATCAGCCATTACCACGTGCTGAAGAATGGAC 450
GDLTVRVVWTDGEEELLLSRAASAITYHVLRKNGH
ATTACACGTCACTTTGGGAGCAAGCAGGAGTACGATGCATCGGCATGTATGAGAGGCGATCATCTTCAACAACAATGTGTCTCT
YTSTRFGSKQGLSCIGMYERGIIFNNVSLW
GGAAAAAGATACGACCACTTACACAGAGCTCTGACAGGTCCAGGTTTGCAGAAGACAGTGGAGGTCTGCTCTCCACAGACTC
KKIRLTHFTRALRTGPGGLQKTVVEVCVSSSTQTH
ACCTGGACGACCTGGACGGTTTGGGTACGTTGACGTCCTCAGTTTGTGCTGCGCTGCACCGTGTGACATCTCCAACAGACTCTTCCTTG
LDDDLDGHLGHVVDVLSLLRCTVVDISNRLFLD
ATGCGCCATCAATGAGAAAGAGCTGGTGAAGATTCTGAAGTATTTGACACGTGGCAACTGTGTATCAACACAGACATTTACT
VPIINKEELLVKILKYFDTWQTVLIKPDIYF
TCAAGTTTACTGGATCCATCAGCAGCAAGGCGGACGTCAGCATGCCATGGGACCTTGTGGAACAGAGAGCGGAGAG
KFDWIHQHRAAAVQELHDAITGDLVEQRRRD
ACGTGGAGCAGGCAGATAAATGGACAACATCACTTACCACAGGCTCATATTTGCACAGAACCTGGCGAGCTGTCTGGGAGAAGC
NFEKTAAPRRYFQPFSGSPRSCVKGKHIAMAM
TGGTGACGTGCGTGGAGATGGTATCGCGGCGCCGACACTCTGTCGTCAGCTCTTCTCATGCTGCTGCTCAAGCAGATC
VQCVLEMVIAAPDTLSVSLFFFMILLRLKQNP
CAGATGTGGAGCTGCAGTGTGCGAGAGATTGACACTTAGTAGTGAGAGGAGCTGCAGAACGTCACCTGCAGAACGTCAGGTGC
DVELQLLREIDTVVGERQLQNGDLQKLQVL
TGGAGAGCTTCATCAAGCAGTGTCTGCGCTTCCACCTGTGGTGGACTTCAGCATGCGCGCGGCGCTGTCCGATGACATCATAGATGGCT
ESFINECLEFRHPVVDFSMRRALSDDIIDGY
ACAGGTACCAAGGCGCAACATCATCTCAACGCGCGGCGGATGACCGCAGAGATTCTTCTGCAAGCTAATGAATCCCGCTGG
RVPKGTNIIILNCRMHRTTEFFCKPNEFRRLD
ACAACTTTGAGAAAACCGCTCTCGACGTTACTTCCAGCAATTCAGGTTTCAGGCCCCGCTCTCGTTGGCAACACATCGCCATGGCA
1440
TGATGAAATCCATCTGGTACATTTGCTCTCCAGTACTCTGTGTGCCCCATGAGGGTCTGACCTGGAGTGCCTCCACAGACCAACA
1530
MKAASILVTLLSQYSVCPHEGLTLLDCLPQTN
ACCTGTCCAGAGCCTGTGGAGATCAGCAGGAGGCCACATCTCAACATGAGATTCTTACCAGACAGAGAGAGCTGGCAACAC
1620
LSQQQPVVEHQEAPHLLNMRFLPRQRGSWQTTL
TCTGAGACTCGACCTTAGCTTCTCCCTGCACATTTACACATATACGCTACATACATATGATCCACACTGCTCATATATGTTAT
1710
*
CTCATGACTGTACAAAGCTAACTTTTATAATTTAATGTGTGTTAACTTGTATTGCTACTGGACCTAATATGCATAATGTAAACGTTTA
1800
GATTAATCTGGATTAAATGATAATTTGTGCAATATTATATATTGTGTCTAATTTTTCTACATTCGGTACAAAATCAGATGAAAT
1890
AACTGGTGCACTTCCAACTAGGAACTGACTGTTTTTCATTTTAAATCTCTCTCATTTTAGACAGCTTATATTTAATTGCTTTTT
1980
TGTAGTTTTCAGTCTGTTTTTTGTTGAAAGTTCTTTACATAAATAAGGTGATTTTCATCATCGAAAAAATTTTTTTTTT
2067

FIGURE 1. Nucleotide and deduced amino acid sequences of Japanese flounder ovarian P450arom cDNA. The portion of the cDNA fragment (426 bp) determined by 5'-RACE is underlined. The polyadenylation signal is double underlined.

of PCR were performed at 94 °C for 30 s, 59 °C for 30 s, 72 °C for 1 min, followed by a final extension at 72 °C for 5 min. The PCR products were electrophoresed on a 2% agarose gel, transferred to nylon membrane (Hybond-N⁺; Amersham), and fixed to the membrane by baking the membrane at 80 °C for 2 h. Following hybridization with either the DIG-labeled 5'-RACE product or EF-1 α cDNA in DIG Easy Hyb solution at 42 °C for 16 h, membranes were washed and visualized according to the methods described above.

For quantification of P450arom mRNA during the period of sex differentiation, the intensity of the hybridization signal was determined by image analyzer software (PDI Inc., Huntington Station, NY, USA). The relative amounts of P450arom mRNA in the gonads at various stages were normalized to EF-1 α . The mean difference between female and male groups was analysed by Student's *t*-test.

Determination of the amounts of sex-steroid hormones in a whole body

Larvae or juveniles were homogenized in 10 ml phosphate-buffered saline (137 mM NaCl, 2.68 mM KCl, 8.1 mM Na₂HPO₄, 1.47 mM KH₂PO₄; pH 7.4)/g body weight. Estradiol-17 β and testosterone were extracted three times from the homogenates into diethyl ether, evaporated by vacuum centrifugation and concentrations were determined according to the enzyme immunoassay described by Van Weemen *et al.* (1979), using enzyme immunoassay kits (Cayman Chemical Co., Ann Arbor, MI, USA). The mean difference between female and male groups was analysed by Student's *t*-test.

RESULTS

Isolation of P450arom cDNA from the Japanese flounder ovary

To isolate P450arom cDNA from Japanese flounder, one million independent cDNA clones in the ovarian

cDNA library were screened with tilapia ovarian P450arom cDNA as a probe, and two positive clones were obtained. The longer clone was 1672 bp which contained the 3'-untranslated region of 444 bp. Compared with the open reading frame (ORF) length (more than 1500 bp) of other P450arom cDNAs, the ORF (1228 bp) of this clone did not seem sufficient for the complete ORF. Therefore, the 5'-end of the cDNA fragment (426 bp) was isolated by 5'-RACE. As there was no difference in the overlapping region (31 bp) of the 5'-RACE clone and library clone, the 5'-RACE clone was considered to be the 5'-end of flounder P450arom. The combined size of the two clones was 2067 bp (Fig. 1) which contained an ORF (69–1623 bp) of 1554 bp encoding a potential protein of 518 amino acid residues. However, there was another potential initiation codon 30 bp downstream of the first ATG. Both start sites resulted in a sequence similar to the consensus sequence proposed by Kozak (1986). The true initiation codon remains uncertain for flounder P450arom. The 3'-untranslated region was 444 bp long and terminated with a poly(A)⁺ tail. The polyadenylation signal, AATAAA, was 18 bp from the first A of the poly(A) track.

The deduced amino acid sequence of the Japanese flounder P450arom was compared with other species (Fig. 2). The overall homology between flounder P450arom and human (Corbin *et al.* 1988, Harada 1988), rat (Hickey *et al.* 1990, Lephart *et al.* 1990), mouse (Terashima *et al.* 1991), bovine (Hinshelwood *et al.* 1993) and chicken (McPhaul *et al.* 1988) P450arom is 50.3, 51.2, 51.9, 50.9 and 50.4% respectively. The homology between flounder and rainbow trout (Tanaka *et al.* 1992), catfish (Trant 1994), medaka (Tanaka *et al.* 1995) or tilapia (Chang *et al.* 1997b) is significantly higher (74.9, 61.5, 76.7, 76.2% respectively). The membrane-spanning (37 to 56 residues), α -helix (302 to 336 residues), Ozols peptide (361 to 383 residues), aromatic (419 to 430 residues) and heme-binding (444 to 467 residues) regions in the flounder P450arom are easily identified by comparison with the corresponding regions in other species.

Tissue-specific expression of P450arom mRNA in adult Japanese flounder

Expression of P450arom mRNA in various adult tissues was examined by RT-PCR. P450arom mRNA was expressed highly in the ovary and spleen, but weakly in the testis and brain (Fig. 3). No difference in the expression pattern in brain or spleen was observed between sexes (data not shown). Signal was barely detectable in the liver, kidney, heart and muscle of both sexes. When a

random primer and a different primer set were used for the reverse transcription reaction and PCR, respectively, the expression pattern of P450arom mRNA was similar (data not shown).

Differential expression of P450arom mRNA in gonads during sex differentiation

The expression pattern of P450arom mRNA during gonadal sex differentiation in both sexes was analysed semi-quantitatively by RT-PCR (Fig. 4). On day 50 after hatching there was no difference in the levels of P450arom mRNA between female and male groups. After day 60, significant differences ($P < 0.01$) in the expression level of P450arom mRNA between the female and male groups were seen. The level of P450arom mRNA in the female group increased rapidly, whereas it decreased slowly in the male group.

Sex-steroid hormone levels during development

To investigate steroid hormone levels during development, whole body concentrations of estradiol-17 β and testosterone were measured (Fig. 5). There were no differences in testosterone levels between the female and male groups during development. Concomitant with the change of P450arom mRNA levels, estradiol-17 β levels in the female group were higher than in the male group after day 60 following hatching. Subsequently, on day 100 there was a significant difference between the female and male groups ($P < 0.05$).

DISCUSSION

In the present study, we isolated P450arom cDNA containing the complete ORF from the Japanese flounder ovary. Similar to rainbow trout, medaka and tilapia, the predicted amino acid sequence of the flounder P450arom has two potential initiating codons. Catfish P450arom has three potential initiating codons. However, the true initiation codon remains uncertain in teleost fish P450arom. The deduced amino acid sequence size of flounder P450arom is 518 amino acid residues if the first potential initiation codon is used or 508 residues if the second codon is used. The deduced amino acid sequence sizes of other teleost fish P450arom cDNA range from 517 to 524 residues if the first potential initiation codon is used or 507–517 residues if the second codon is used. In comparison with the deduced amino acid sequence size of mammalian and chicken P450arom (502–508 residues), teleost fish P450arom tends to be longer.

		I	
flounder	M-DRIPA-CDLMTFVGLGAALGD-LVSTSPNATAVRTGISVASRTLLVLCVLLVAV-SHTDRRTVPGPFCLGLGPLLSYVRFIWTG	86	
tilapia	M-DLISA-CEQAMNFVGL---DA-VVARSLCDLKCHPIDGISMATRLILLVCLLLVAV-SHTDKKIVPGPSFCLGLGPLLSYLRFIWTG	83	
medaka	M-DLIPA-CDRTMSSSL---AE-LVS IAPNTVGLPSGIPMATRSLILLVGLLLMVS-SHSEKKTIPGPSFCLGLGPLMSYLRFIWTG	82	
catfish	MAAHVFPCECERTRKVFHFSETVMEILLREARNGTDPRIENPRGITLLLLCLVLLTVVNRHEKKCSIPGPSFCLGLGPLMSYCRFIWTG	90	
trout	M-DLLSPVCGRVMAVVCDDTVIADLLVSESRNATATRSEGISLATGSLLLLLCLLAAW-RHTDNNVPGPFCLGLGPLLSYLRFIWTG	88	
chicken	-----MIPETLNLNLYF-TSLVPDLMPVATVPIIILICFLFLIWN-HEETSSIPGPGYCMGIGPLISHGRFLWMG	68	
bovine	-----MLLEVLNPRHYNVTSMVSEVVP IAS IAILLTGFLLLVWN-YEDTSSIPGPSYFLGIGPLISHGRFLWMG	69	
mouse	-----MFLEMLNPMQYNVTIMVPEVTVPVSAMPELLIMGLLLLIWN-CESSSSIPGPGYCLGIGPLISHGRFLWMG	69	
rat	-----MFLEMLNPMHYNVTIMVPEVTVPVSAMPELLIMGLLLLIWN-CESSSSIPGPGYCLGIGPLISHGRFLWMG	69	
human	-----MVLEMLNPIHYNITSIPEAMPATMPVLLLTGLFLLLVWN-YEGTSSIPGPGYCMGIGPLISHGRFLWMG	69	
		*** * * * * *	
flounder	IGTACNYNKKRYGDIVRVWIDGEETLILSRASAIYHVLKNGHYTSRFGSKQLGSCIGMYERGIIFNNVSLWKKIRTHFTALTGPGLQK	176	
tilapia	IGTASNYNKKYGDIVRVWINGEETLILSRSSAVHHVLKNGNYTSRFGSIQGLSYLGMNERGIIFNNVTLWKKIRTYFAKALTGPGLNQ	173	
medaka	IGTASNYNKKYGDIVRVWINGEETLILSRASAVHHVLKNGHYTSRFGSKQLGSCIGMNEKGIIFNNVSLWKKIRTYFAKALTGPGLNQ	172	
catfish	IGTASNYNKKYGDIVRVWISGEETLILSRPSAVHHVLKHSQYTSRFGSKQLGQICGMHEQGIIFNSNVLWKKIRTYFAKALTGPGLQ	180	
trout	IGTASNYNKKYGDIVRVWINGEETLILSSSAVHHVLKNGHYTSRFGSKQLGSCIGMDERGIIFNSNMLWKKIRTYFAKALTGPGLQ	178	
chicken	VGNACNYNKKYGEFVRVWISGEETLIISSSSVFHVMKHWNYVSFRGSKLGLQCIGMYENGIIFNNPFAHWKEIRPFTKALSGPGLVR	158	
bovine	IGSACNYNKKYGEFMRVWVCGEETLIISSSSMFHVMKHSYISFRGSKLGLQFICGMHEKGIIFNNPALWKAIRPFTKALSGPGLVR	159	
mouse	IGSACNYNKKYGEFMRVWISGEETLIISSSSMFHVMKHSYISFRGSKRGLQCIGMHENGIIFNNPNSLWRTIRPFFMKALTGPGLVR	159	
rat	IGSACNYNKKYGEFMRVWISGEETLIISSSSMFHVMKHSYISFRGSKRGLQCIGMHENGIIFNNPNSLWRTIRPFFMKALTGPGLVR	159	
human	IGSACNYNRRVYGEFMRVWISGEETLIISSSSMFHIMKHNHYSRFGSKLGLQCIGMHEKGIIFNNPELWKTIRPFFMKALSGPGLVR	159	
		* * * * *	
flounder	TVEVCVSSQTTHLDDLGL----GHVDVLSLLRCTVVDISNRLFLDVPINEKELLVKILKYFDTWQTVLIKPDYIFKFDWIHQHKA	261	
tilapia	TVDVCVSSIQAHLDHLDL----GHVDVLSLLRCTVVDISNRLFLNVLNKEKELMLKIQKYFHTWQDVLIKPDYIFKFWIHHRRKAT	258	
medaka	TVEVCVTSQTTHLDNLSSL----SYVDVLGLFRGTVVDISNRLFLGVPVDEKELQKIHKYFDTWQTVLIKPDYIFKFWIHHRRKAT	257	
catfish	TLEICTMSTNTHLDGLSRLTDAQ-GHVDVLSLLRCTVVDISNRLFLDVLNQLLFFKIHRYFETWQTVLIKPDYIFKRLKWLHDKHRNA	269	
trout	TVDVCVSSQTTHLDAQPGDGLMGQVVDLSLLRCTVVDISNRLFLGVPVDEKELQKIQKYFDTWQTVLIKPDYIFKDLWHEKHRTA	268	
chicken	MIAICVESITVHLDLLEEVTEV-GNVNVLNLMRRLIMLDTSNLFLGVPVDESAIVLKIQNYFDAWQALLKPDIFFKISWLKRYKESV	247	
bovine	MVTICADSITKHLDRLEEVNDL-GYVDVLTLMRRLIMLDTSNLFLGIPDESAIVVNIQGYFDAWQALLKPDIFFKISWLKRYKESV	248	
mouse	MVEVCVESIKQHLDRLEEVTDTS-GYVDVLTLMRRLIMLDTSNLFLGIPDESAIVKIQGYFNAWQALLIKPNIFFKISWLRYKERSV	248	
rat	MVEVCVESIKQHLDRLEEVTDNS-GYVDVLTLMRRLIMLDTSNLFLGIPDESAIVKIQGYFNAWQALLIKPNIFFKISWLRYKERSV	248	
human	MVTVCAESLTKHLDRLEEVTDNS-GYVDVLTLMRRLIMLDTSNLFLGIPDESAIVKIQGYFDAWQALLIKPDIFFKISWLRYKERSV	248	
		* * * * *	
		II	
flounder	QELHDAIGDLVEQKRRDVEQADKL-DNINFTTGLIFAQNHGELSAENVQCVLEMVIAAPDTLSVSLFFMMLLLKQNPDELQLLREIDT	350	
tilapia	QELQDAIKRLVDQKRRMEQADKL-DNINFTTGLIFAQNHGELSAENVQCVLEMVIAAPDTLSLFFMMLLLKQNPDELQLLREIDT	347	
medaka	QELQDAIESLVERKKRMEQAEKL-DNINFTTGLIFAQNHGELSAENVQCVLEMVIAAPDTLSLFFMMLLLKQNPDELQLLREIDT	346	
catfish	QELHDAIEDLIEQKRTLQQAELK-DNINFTTGLIFAQNHGELSAENVQCVLEMVIAAPDTLSISVFFMMLLLKQNPDELQLLREIDT	358	
trout	QELDAEDLSLVQKRRGLQADKL-DHINFTADLIFAQNHGELSAENVQCVLEMVIAAPDTLSLFFMMLLLKQNPDELQLLREIDT	357	
chicken	KDLKGAMEILIEQKQKSLTVEKLEDMDFASQLIFAQNRGDLTAENVQCVLEMMIAAPDTLSVTLFIMLLIADDPVVEKMMREIET	337	
bovine	KDLKDAMEILIAEKRRHISTAEKLEDSIDFATELIFAERKRGELTRENVNQCILEMLIAAPDTMSVSFFMFLIAKHPQVEEAIIREIQT	338	
mouse	KDLKDEIAVLVEKRRKVSIAEKLEDCMDFATDLIFAERRGDLTKENVNQCILEMLIAAPDTMSVTLYFMLLLVAEYPEVEAAILKEIHT	338	
rat	KDLKDEIEILVEKRRKQVSSAEKLEDCMDFATDLIFAERRGDLTKENVNQCILEMLIAAPDTMSVTLYFMLLLIAEYPEVEATILKEIHT	338	
human	KDLKDAIEVLIAEKRRRISTEAEKLEDCMDFATELILAEKRGDLTRENVNQCILEMLIAAPDTMSVSFFMFLIAKHPNVEEAIKEIQT	338	
		* * * * *	
		III	
flounder	VVGERQLQNGDLQKLVLESFNECLRFHPVVDFTMRRLSDDIIDGYRVPKGTNIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	439	
tilapia	VVGERQLQNGDLQKLVLESFNECLRFHPVVDFTMRRLSDDIIDGYRVPKGTNIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	437	
medaka	IVGDSQLQNGDLQKLVLESFNECLRFHPVVDFTMRRLSDDIIDGHRVQKGTNIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	435	
catfish	VLGDTLQSHLSLQKLVLECFINEALRFHPVVDFTMRRLSDDIIDGHRVQKGTNIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	447	
trout	AIGDRELHNSDLQNLRLVESFNECLRFHPVVDFTMRRLSDDIIDGHRVQKGTNIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	446	
chicken	VMGHRVQSDDMFNKLVENFIYESMRYQPVVDLIMRKALQDDVIDGYPVKKGNTIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	426	
bovine	VVGERDIRIDDMQKLVENFIYESMRYQPVVDLIMRKALQDDVIDGYPVKKGNTIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	427	
mouse	VVGDRDIRIDDMQKLVENFIYESMRYQPVVDLIMRKALQDDVIDGYPVKKGNTIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	427	
rat	VVGDRDIRIDDMQKLVENFIYESMRYQPVVDLIMRKALQDDVIDGYPVKKGNTIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	427	
human	VIGERDIKIDDMQKLVENFIYESMRYQPVVDLIMRKALQDDVIDGYPVKKGNTIILNIGRMHRTTEFFCKPNEFLDNFEKTAPR-RYF	427	
		* * * * *	
		IV	
flounder	QPFSGSPRSCVKGHIAMAMMKSILVTLSSQYSVCPHEGLTLDCLPQTNNSLQQPVEHQ--EAPHLNMRFLPRQGSQWTL-----	518	
tilapia	QPFSGSPRACIGHMAAMMKSILVTLSSQYSVCTHEGLTLDCLPQTNNSLQQPVEHQQA-ETEHLNMRFLPRQGSQWTL-----	522	
medaka	QPFSGSPRACVGNIAAMMKSILVTLSSQYSVCPHEGLTLDCLPQTNNSLQQPVEHQ--EADHLSMTFLPRQGIWES-PSF-F	517	
catfish	QPFSGSPRSCVKGHIAMAMMKSILVTLSSQYSVCPHEGLTLDCLPQTNNSLQQPVEHQ--EADHLSMTFLPRQGIWES-PSF-F	524	
trout	QPFSGSPRSCVKGHIAMAMMKSILVTLSSQYSVCPHEGLTLDCLPQTNNSLQQPVEHQ--EADHLSMTFLPRQGIWES-PSF-F	522	
chicken	QPFSGSPRSCVKGHIAMAMMKSILVTLSSQYSVCPHEGLTLDCLPQTNNSLQQPVEHQ--EADHLSMTFLPRQGIWES-PSF-F	507	
bovine	QPFSGSPRSCVKGHIAMAMMKSILVTLSSQYSVCPHEGLTLDCLPQTNNSLQQPVEHQ--EADHLSMTFLPRQGIWES-PSF-F	503	
mouse	QPFSGSPRSCVKGHIAMAMMKSILVTLSSQYSVCPHEGLTLDCLPQTNNSLQQPVEHQ--EADHLSMTFLPRQGIWES-PSF-F	503	
rat	QPFSGSPRSCVKGHIAMAMMKSILVTLSSQYSVCPHEGLTLDCLPQTNNSLQQPVEHQ--EADHLSMTFLPRQGIWES-PSF-F	508	
human	QPFSGSPRSCVKGHIAMAMMKSILVTLSSQYSVCPHEGLTLDCLPQTNNSLQQPVEHQ--EADHLSMTFLPRQGIWES-PSF-F	503	

		V	

FIGURE 2. Alignment of the amino acid sequence of Japanese flounder P450arom with tilapia, medaka, catfish, rainbow trout, chicken, bovine, mouse, rat and human. Asterisks indicate identical amino acid residues. Roman numerals indicate the membrane-spanning region (I), the helix region (II), the Ozols peptide region (III), the aromatic region (IV) and the heme-binding region (V).

FIGURE 3. Tissue-specific expression of P450arom transcript in adult Japanese flounder. EF-1 α serves as an internal control for loading differences. RT-PCR amplified fragments specific for P450arom and EF-1 α are predicted to be 344 bp and 577 bp respectively. PCR products were fractionated on a 2% agarose gel and confirmed by Southern blotting with labeled P450arom or EF-1 α cDNA.

RT-PCR shows that the flounder P450arom mRNA is expressed highly in the ovary and spleen but weakly in the testis and brain. All transcripts are detected as a single band. In catfish, the expression of P450arom mRNA is observed as a single band in the testis and brain, but as double bands in the ovary (Trant *et al.* 1997). Recently, two CYP19 genes encoding different P450arom isozymes in brain and ovary have been identified in goldfish. RT-PCR analysis has shown that brain- and ovary-type P450arom cDNA variants are found in the forebrain, whereas only the ovary-type variant is detected in the ovary (Callard & Tchoudakova 1997, Tchoudakova & Callard 1998). In the present study, there was no difference in the sequences (344 bp) of RT-PCR products among the ovary, spleen, testis and brain (data not shown), suggesting that none of the RT-PCR products is a brain-type variant. It therefore remains to be clarified whether the brain-type variant exists in the flounder.

Expression of the P450arom gene in vertebrate spleen has only been reported in human fetal spleen (Price *et al.* 1992). Although it is known that P450arom plays important physiological functions in various tissues by regulating estrogen biosynthesis, the role of P450arom in spleen remains unclear. However, the presence of an mRNA encoding P450 cholesterol side-chain cleavage was detected in the spleen, suggesting a steroidogenic function similar to gonads and the adrenal cortex (K-I Morohashi, personal communication). The significance of P450arom mRNA expression in the flounder spleen needs to be studied further.

The expression of P450arom mRNA in the gonads during sexual differentiation was analysed by RT-PCR. Previous histological observations showed that ovarian and testicular differentiation in the flounder were initiated in juveniles about 30 mm

FIGURE 4. Differential expression of P450arom transcripts in female and male gonads during sex differentiation. (A) P450arom transcript levels in total RNA isolated from ten pooled gonads from either the female or male group were examined by RT-PCR at 50, 60, 80 or 100 days after hatching. RT-PCR and Southern blotting were done as described in Fig. 3. (B) Relative amounts of P450arom mRNA in female (solid bars) and male (open bars) gonads were normalized to EF-1 α . Vertical bars indicate the mean $\pm$ standard error of the five replicates. ** $P < 0.01$ compared with the male group (Student's *t*-test).

in total length (TL) and 37 mm in TL respectively (Tanaka 1987, Yamamoto 1995, 1999). In the present study, the average TLs of the female and male groups at day 60 after hatching were 31.6 mm and 39.0 mm respectively. Thus, this period is thought to be the time of initial sex differentiation for both sexes. On day 50 after hatching when the gonad is sexually indifferent, there was no difference in the levels of P450arom mRNA between the female and male groups. After the initiation of sex differentiation (day 60), the level of P450arom mRNA increased rapidly in the female group, whereas it decreased slowly in the male group. In this study, females were produced artificially by mating normal females with gynogenetic diploid

FIGURE 5. Whole body concentrations of estradiol-17 β (A) and testosterone (B) in the female (solid bars) and male (open bars) groups during development.

Sex-steroid hormones extracted from five to twenty pooled larvae or juveniles on day 20, 40, 60, 80 or 100 after hatching were measured by enzyme immunoassay. Vertical bars indicate the mean $\pm$ standard error of the five replicates. * $P < 0.05$ compared with the male group (Student's *t*-test).

males, reared at normal water temperature (18 °C), resulting in all genetic females (XX). Males were produced by rearing the same genetic females as described above at higher water temperature (27 °C) during sex differentiation, resulting in conversion to all phenotypic males. Thus, we present, to the best of our knowledge, the first demonstration that the expression of P450arom mRNA in genetically identical (XX) gonads is inhibited by rearing the larvae at a high water temperature, resulting in conversion of genetically female larvae into phenotypic males.

In reptiles with temperature-dependent sex determination (TSD), P450arom mRNA levels in both sexes have been examined during sexual differentiation (Jeyasuria & Place 1997). In embryos of the diamondback terrapin (*Malaclemys terrapin*), P450arom mRNA levels in female adrenal/kidney/gonadal complexes are significantly higher than in

male complexes at the sexually differentiated stage. Thus, temperature appears to play an important role in the regulation of P450arom gene expression in fishes and reptiles with TSD. Studies on the regulation of P450arom gene expression by temperature would provide insight into the sex determination mechanisms in lower vertebrates.

Estradiol-17 β levels in the whole body of the female group were elevated relative to the male group after sex differentiation was initiated. In contrast, no differences in the whole body testosterone levels in the female and male groups were observed throughout the course of sex differentiation. P450arom catalyses the conversion from testosterone to estradiol-17 β . Consequently, the change of estradiol-17 β levels in female and male groups during sex differentiation is consistent with the changes in P450arom mRNA levels. The rate of increase in estradiol-17 β levels in the female group, however, was moderate relative to the marked change in P450arom mRNA levels. There was basal estradiol-17 β before and after sex differentiation. In addition to the ovary, a significant amount of P450arom mRNA was expressed in the spleen. Since there were no differences in P450arom mRNA levels in brain and spleen between the female and male groups during the sex differentiation (data not shown), the increase of estradiol-17 β levels above basal levels in the female group may be derived from the increased P450arom mRNA levels in ovaries. In tilapia, P450arom-like immunoreactivity appears in differentiating ovaries, but not in differentiating testes (Chang *et al.* 1997a). In the embryos of the European pond turtle (*Emys orbicularis*), aromatase activity remains low in differentiating testes, but increases in differentiating ovaries (Desvages & Pieau 1992). In the marine turtle (*Dermochelys coriacea*), P450arom activity and estrogen content increase in gonads of the embryos incubated at a feminizing temperature, whereas they remain low in gonads of the embryos incubated at a masculinizing temperature (Desvages *et al.* 1993). In the larvae of urodele amphibia (*Pleurodeles waltl*), aromatase activity is significantly higher in the gonads of genetic females (ZW) than of genetic males (ZZ) during sex differentiation. The gonads of sex-reversed males (ZW) produced by exposure of the larvae to a high temperature have low aromatase activity similar to those of genetic males (Chardard *et al.* 1995). These results suggest that endogenous estrogen plays a key role in gonadal differentiation of fishes, amphibians and reptiles with TSD.

In chinook salmon (*Oncorhynchus tshawytscha*), brief treatment with an aromatase inhibitor during sex differentiation causes genetic females to develop

into normal phenotypic males (Piferrer *et al.* 1994). This result indicates that P450arom activity and estradiol-17 β biosynthesis are indispensable for sex differentiation into females in some teleost fishes. Based on increases seen during ovarian differentiation, we suggest that expression of P450arom mRNA and estradiol-17 β biosynthesis is indispensable for ovarian differentiation. In contrast, the expression of P450arom mRNA and estradiol-17 β biosynthesis is inhibited during testicular differentiation, suggesting suppression as a prerequisite for testicular differentiation. Future studies will focus on the effect of aromatase inhibitors in the sex differentiation of the flounder.

ACKNOWLEDGEMENTS

We thank the staff of the Kumamoto Prefectural Fisheries Research Center for the rearing of fish. This work was supported by Grants-in-Aid for Scientific Research (no. 09480206 to S A) and priority areas (no. 07283104 to S A and no. 10160216 to K T) from the Ministry of Education, Science, Sports and Culture of Japan and Grants-in-Aid for Research for the Future (JSPS-RFTF 96L00401 to Y N) from the Japan Society for the Promotion of Science.

REFERENCES

- Callard GV & Tchoudakova A 1997 Evolutionary and functional significance of two CYP19 genes differentially expressed in brain and ovary of goldfish. *Journal of Steroid Biochemistry and Molecular Biology* **61** 387–392.
- Chang XT, Nakamura M, Kobayashi T & Nagahama Y 1997a Immunocytochemical localization of steroidogenic enzymes in gonads during sex differentiation in tilapia. *Journal of Reproduction and Development* **43** 73–74.
- Chang XT, Kobayashi T, Kajiura H, Nakamura M & Nagahama Y 1997b Isolation and characterization of the cDNA encoding the tilapia (*Oreochromis niloticus*) cytochrome P450 aromatase (P450arom): changes in P450arom mRNA, protein and enzyme activity in ovarian follicles during oogenesis. *Journal of Molecular Endocrinology* **18** 57–66.
- Chardard D, Desvages G, Pieau C & Dournon C 1995 Aromatase activity in larval gonads of *Pleurodeles waltl* (urodele amphibia) during normal sex differentiation and during sex reversal by thermal treatment effect. *General and Comparative Endocrinology* **99** 100–107.
- Chirgwin JM, Przybyla AE, MacDonald RJ & Rutter WJ 1979 Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. *Biochemistry* **18** 5294–5299.
- Conover DO & Kynard BE 1981 Environmental sex determination: interaction of temperature and genotype in a fish. *Science* **213** 577–579.
- Corbin CJ, Graham-Lorence S, McPhaul M, Mason M, Mendelson CR & Simpson ER 1988 Isolation of a full-length cDNA insert encoding human aromatase system cytochrome P-450 and its expression in nonsteroidogenic cells. *Proceedings of the National Academy of Sciences of the USA* **85** 8948–8952.
- Desvages G & Pieau C 1992 Aromatase activity in gonads of turtle embryos as a function of the incubation temperature of eggs. *Journal of Steroid Biochemistry and Molecular Biology* **41** 851–853.
- Desvages G, Girondot M & Pieau C 1993 Sensitive stages for the effects of temperature on gonadal aromatase activity in embryos of the marine turtle *Dermochelys coriacea*. *General and Comparative Endocrinology* **92** 54–61.
- Feinberg AP & Vogelstein B 1983 A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Analytical Biochemistry* **132** 6–13.
- Frohman MA, Dush MK & Martin GR 1988 Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer. *Proceedings of the National Academy of Sciences of the USA* **85** 8998–9002.
- Gubler U & Hoffman BJ 1983 A simple and very efficient method for generating cDNA libraries. *Gene* **25** 263–269.
- Harada N 1988 Cloning of a complete cDNA encoding human aromatase: immunochemical identification and sequence analysis. *Biochemical and Biophysical Research Communications* **156** 725–732.
- Hickey GJ, Krasnow JS, Beattie WG & Richards JS 1990 Aromatase cytochrome P450 in rat ovarian granulosa cells before and after luteinization: adenosine 3',5'-monophosphate-dependent and independent regulation. Cloning and sequencing of rat aromatase cDNA and 5' genomic DNA. *Molecular Endocrinology* **4** 3–12.
- Hinshelwood MM, Corbin CJ, Tsang PCW & Simpson ER 1993 Isolation and characterization of a complementary deoxyribonucleic acid insert encoding bovine aromatase cytochrome P450. *Endocrinology* **133** 1971–1977.
- Hunter GA & Donaldson EM 1983 Hormonal sex control and its application to fish culture. In *Fish Physiology*, vol. IX, part B, pp 223–303. Eds WS Hoar, DJ Randall & EM Donaldson. New York: Academic Press.
- Jeyasuria P & Place AR 1997 Temperature-dependent aromatase expression in developing diamondback terrapin (*Malaclemys terrapin*) embryos. *Journal of Steroid Biochemistry and Molecular Biology* **61** 415–425.
- Kozak M 1986 Point mutations define a sequence flanking the AUG initiator codon that modulates translation by eukaryotic ribosomes. *Cell* **44** 283–292.
- Lephart ED, Peterson KG, Noble JF, George FW & McPhaul MJ 1990 The structure of cDNA clones encoding aromatase P-450 isolated from a rat Leydig cell tumor line demonstrates differential processing of aromatase mRNA in rat ovary and a neoplastic cell line. *Molecular and Cellular Endocrinology* **70** 31–40.
- McPhaul MJ, Noble JF, Simpson ER, Mendelson CR & Wilson JD 1988 The expression of a functional cDNA encoding the chicken cytochrome P-450arom (aromatase) that catalyzes the formation of estrogen from androgen. *Journal of Biological Chemistry* **263** 16358–16363.
- Matsumine H, Hirato K, Yanaihara T, Tamada T & Yoshida M 1986 Aromatization by skeletal muscle. *Journal of Clinical Endocrinology and Metabolism* **63** 717–720.
- Pasmanik M & Callard GV 1985 Aromatase and 5 alpha-reductase in the teleost brain, spinal cord, and pituitary gland. *General and Comparative Endocrinology* **60** 244–251.
- Piferrer F, Zanuy S, Carrillo M, Solar II, Devlin RH & Donaldson EM 1994 Brief treatment with an aromatase inhibitor during sex differentiation causes chromosomally female salmon to develop as normal, functional males. *Journal of Experimental Zoology* **270** 255–262.

- Price T, Aitken J & Simpson ER 1992 Relative expression of aromatase cytochrome P450 in human fetal tissues as determined by competitive polymerase chain reaction amplification. *Journal of Clinical Endocrinology and Metabolism* **74** 879–883.
- Rubin DA 1985 Effect of pH on sex ratio in cichlids and a poeciliid (Teleostei). *Copeia* **1985** 233–235.
- Sanger F, Air GM, Barrell BG, Brown NL, Coulson AR, Fiddes CA, Hutchison CA, Slocombe PM & Smith M 1977 Nucleotide sequence of bacteriophage phi X174 DNA. *Nature* **265** 687–695.
- Sasano H, Uzuki M, Sawai T, Nagura H, Matsunaga G, Kashimoto O & Harada N 1997 Aromatase in human bone tissue. *Journal of Bone and Mineral Research* **12** 1416–1423.
- Smith CA, Andrews JE & Sinclair AH 1997 Gonadal sex differentiation in chicken embryos: expression of estrogen receptor and aromatase genes. *Journal of Steroid Biochemistry and Molecular Biology* **60** 295–302.
- Sullivan JA & Schultz RJ 1986 Genetic and environmental basis of variable sex ratios in laboratory strains of *Poeciliopsis lucida*. *Evolution* **40** 152–158.
- Tabata K 1991 Induction of gynogenetic diploid males and presumption of sex determination mechanisms in the hirame *Paralichthys olivaceus*. *Bulletin of the Japanese Society of Scientific Fisheries* **57** 845–850.
- Tanaka H 1987 Gonadal sex differentiation in flounder, *Paralichthys olivaceus*. *Bulletin of National Research Institute of Aquaculture* **11** 7–19.
- Tanaka H 1988 Effects of estradiol-17 β on gonadal sex differentiation in flounder, *Paralichthys olivaceus*. *Bulletin of National Research Institute of Aquaculture* **13** 17–23.
- Tanaka M, Telecky TM, Fukada S, Adachi S, Chen S & Nagahama Y 1992 Cloning and sequence analysis of the cDNA encoding P-450 aromatase (P450arom) from a rainbow trout (*Oncorhynchus mykiss*) ovary; relationship between the amount of P450arom mRNA and the production of oestradiol-17 β in the ovary. *Journal of Molecular Endocrinology* **8** 53–61.
- Tanaka M, Fukada S, Matsuyama M & Nagahama Y 1995 Structure and promoter analysis of the cytochrome P-450 aromatase gene of the teleost fish, medaka (*Oryzias latipes*). *Journal of Biochemistry* **117** 719–725.
- Tchoudakova A & Callard GV 1998 Identification of multiple CYP19 genes encoding different cytochrome P450 aromatase isozymes in brain and ovary. *Endocrinology* **139** 2179–2189.
- Terashima M, Toda K, Kawamoto T, Kuribayashi I, Ogawa Y, Maeda T & Shizuta Y 1991 Isolation of a full-length cDNA encoding mouse aromatase P450. *Archives of Biochemistry and Biophysics* **285** 231–237.
- Trant JM 1994 Isolation and characterization of the cDNA encoding the channel catfish (*Ictalurus punctatus*) form of cytochrome P450arom. *General and Comparative Endocrinology* **95** 155–168.
- Trant JM, Lehrter J, Gregory T, Nunez S & Wunder J 1997 Expression of cytochrome P450 aromatase in the channel catfish, *Ictalurus punctatus*. *Journal of Steroid Biochemistry and Molecular Biology* **61** 393–397.
- Van Weemen BK, Bosch AM, Dawson EC & Schuurs AH 1979 Enzyme-immunoassay of steroids: possibilities and pitfalls. *Journal of Steroid Biochemistry* **11** 147–151.
- Wallace RA 1985 Vitellogenesis and oocyte growth in nonmammalian vertebrates. In *Developmental Biology*, pp 127–177. Ed LW Browder. New York: Plenum Press.
- Yamamoto E 1995 Studies on sex-manipulation and production of cloned populations in hirame flounder, *Paralichthys olivaceus* (Temminck et Schlegel). *Bulletin of the Tottori Prefectural Fisheries Experimental Station* **34** 1–145.
- Yamamoto E 1999 Studies on sex-manipulation and production of cloned populations in hirame, *Paralichthys olivaceus* (Temminck et Schlegel). *Aquaculture* **173** 235–246.
- Yamamoto E, Hirano R & Masutani R 1992 Progeny test of methyltestosterone-induced males of gynogenetic diploids and normal diploids in the hirame flounder, *Paralichthys olivaceus*, with evidence for the male heterogamety. *Bulletin of the Tottori Prefectural Fisheries Experimental Station* **33** 9–17.
- Yamazaki F 1983 Sex control and manipulation in fish. *Aquaculture* **33** 329–354.
- Yoshida K, Shimada K & Saito N 1996 Expression of P450_{17 α} hydroxylase and P450 aromatase genes in the chicken gonad before and after sexual differentiation. *General and Comparative Endocrinology* **102** 233–240.

REVISED MANUSCRIPT RECEIVED 21 April 1999