Analysis of the estrogen regulation of the zebrafish estrogen receptor (ER) reveals distinct effects of ER α , ER β 1 and ER β 2

A Menuet, Y Le Page, O Torres, L Kern, O Kah and F Pakdel

Endocrinologie Moléculaire de la Reproduction, UMR CNRS 6026, Université de Rennes 1, 35042 Rennes cedex, France

(Requests for offprints should be addressed to F Pakdel, Endocrinologie Moléculaire de la Reproduction, UMR CNRS 6026, Université de Rennes 1, Campus de Beaulieu, 35042 Rennes cedex, France; Email : farzad.pakdel@univ-rennes1.fr)

Abstract

We have previously cloned and characterized three estrogen receptors (ER) in the zebrafish (zfER α , zfER β 1 and zfER β 2). We have also shown that they are functional *in vitro* and exhibit a distinct expression pattern, although partially overlapping, in the brain of zebrafish. In this paper, we have shown that the hepatic expression of these zfER genes responds differently to estradiol (E2). In fact, a 48-h direct exposure of zebrafish to E2 resulted in a strong stimulation of zfER α gene expression while zfER β 1 gene expression was markedly reduced and zfER β 2 remained virtually unchanged. To establish the potential implication of each zfER in the E2 upregulation of the zfER α gene, the promoter region of this gene was isolated and characterized. Transfection experiments with promoter–luciferase reporter constructs together with different zfER expression vectors were carried out in different cell contexts. The data showed that *in vivo* E2 upregulation of the zfER α gene requires ER α itself and a conserved transcription unit sequence including at least an imperfect estrogen-responsive element (ERE) and an AP-1/ERE half site at the proximal transcription initiation site. Interestingly, although in the presence of E2 zfER α was the most potent at inducing the expression of its own gene, the effect of E2 mediated by zfER β 2 represented 50% of the zfER α activity. In contrast, zfER β 1 was unable to upregulate the zfER α gene whereas this receptor form was able to tightly bind E2 and activate a reporter plasmid containing a consensus ERE. Altogether, these results indicated that the two ER β forms recently characterized in teleost fish could have partially distinct and not redundant functions.

Journal of Molecular Endocrinology (2004) **32**, 975–986

Introduction

In all vertebrates, estradiol (E2) is involved in numerous physiological processes during development and adult life. These effects are mediated by members of the nuclear receptor family, the estrogen receptors ER α and ER β . These proteins classically regulate the expression of E2 target genes by direct binding with a specific palindromic DNA sequence called the estrogen-responsive element (ERE: AGGTCA_nnnTGACCT) which permits recruitment of cofactors necessary for transcription (Zilliacus *et al.* 1995, Klinge 2000). Moreover, an important number of E2-sensitive genes, which do not contain ERE but do contain other cis-elements such as AP-1 or Sp1, have been described. It was recently proved that, in this case,

ERs can regulate the transcription by direct protein–protein interactions with the AP-1 or Sp1 transcription factors (Paech *et al.* 1997, Webb *et al.* 1999, Saville *et al.* 2000). Although the mammalian ERs share the same modular organization, recent data showed that they may have different transcriptional capacities on E2 target genes (Kuiper *et al.* 1997, Tremblay *et al.* 1997). Differences between ER subtypes in relative ligand-binding affinity have been described. A range of natural and synthetic agonists or antagonists induced distinct conformational changes in the tertiary structure of the ERs which induced differential cofactor recruitment. In consequence, the effects of E2 at the transcriptional level depend on the promoter, the presence of cell-specific factors and the ER subtype, which

could explain in part the pleiotropic effects of E2 at the physiological level.

In addition to natural hormones or effectors, including growth factors, several classes of environmental contaminants could also interact with ERs and modulate the expression of E2 target genes. These latter molecules, usually referred to as endocrine disruptors, are susceptible to impairing the reproductive success in all classes of vertebrates (Petit *et al.* 1997, Tyler *et al.* 1998, Le Guevel & Pakdel 2001, McLachlan 2001, Jobling *et al.* 2003).

Although in mammals only two ER subtypes have been characterized (Green *et al.* 1986, Kuiper *et al.* 1996, Mosselman *et al.* 1996), we and others have recently reported the existence of three ER forms in fish species: ER α , ER β 1 and ER β 2 (Hawkins *et al.* 2000, Ma *et al.* 2000, Menuet *et al.* 2002). These receptors are generated from three distinct genes, are able to bind E2 with high affinity and activate, with approximately the same fold induction, a reporter gene under the control of a consensus ERE (Bardet *et al.* 2002, Lassiter *et al.* 2002, Menuet *et al.* 2002). These results clearly demonstrated that these new ER β forms were functional ligand-dependent transcription factors. However, *in vivo*, the majority of E2 target genes present imperfect ERE sequences or complex organizations with, notably, the presence of $\frac{1}{2}$ ERE close to the ERE. In consequence, it is necessary to investigate the respective transcriptional enhancement capacities of the ER β forms on an endogenous target gene.

In oviparous species, in addition to the hypothalamic–pituitary–gonad axis, the liver is an important E2 target organ, in which the synthesis of the yolk protein vitellogenin by hepatocyte cells is strictly under the control of E2. The vitellogenin production which is crucial for embryo development and reproduction success is tightly coupled to a substantial E2-dependent upregulation of ER α gene expression. In chicken, *Xenopus* and trout, E2 treatment increases the ER α mRNA and protein accumulation in the liver (Pakdel *et al.* 1991, Ninomiya *et al.* 1992, Lee *et al.* 1995). This induction occurs at the transcriptional and post-transcriptional levels (Flouriot *et al.* 1996a).

Here, we have shown that zebrafish (zf) ER β 1 and zfER β 2 mRNAs are co-expressed with zfER α in the liver and that E2 treatment differentially modifies the expression levels of these receptor

transcripts. To investigate whether or not zfER β 1 and zfER β 2 are involved in hepatic vitellogenesis and therefore in hepatic zfER α regulation, the zfER α promoter region was isolated and characterized. The data showed that, in contrast to zfER β 1, zfER β 2 was able to significantly stimulate the expression of the zfER α gene in the presence of E2. Moreover, expression of zfER β 1 was strongly downregulated by *in vivo* E2 treatment. This, added to the fact that *in vitro* data demonstrated that zfER β 1 is not directly implicated in the E2 upregulation of zfER α , suggests diverging functions for zfER β 1 and zfER β 2 in the liver.

Materials and methods

Animals

In this study, mature zebrafish (*Danio rerio*), raised in our facilities, were anesthetized on ice. Hormonal treatments consisted of exposure of fish for 48 h to 10^{-8} M 17 β -estradiol (E2) or to a solvent control (0.1% ethanol) before anesthesia. The liver was removed by dissection and total RNA was extracted using the Trizol method according to the manufacturer's instructions (Gibco-BRL, Eggenstein, Germany). The liver RNA samples were enriched in poly(A)⁺-RNA by the oligotex mRNA mini kit (Qiagen, Courtaboeuf, France).

Northern blot assay

Northern blot experiments were performed according to the previously published protocol (Thomas 1980). Poly(A)⁺-RNA (1.5 μ g) from E2-treated or untreated zebrafish liver was denatured at 65 °C for 10 min in formamide–formaldehyde solution, separated on agarose–formaldehyde gel and transferred onto a nylon membrane (Hybond-N; Amersham, Uppsala, Sweden). RNA were fixed by u.v. exposure (254 nm for 2 min) and by baking at 80 °C for 2 h. The membrane was prehybridized for 5–6 h, hybridized overnight with the probe labeled with [α -³²P]dCTP corresponding to the coding region of each zfER cDNA or to acidic ribosomal phosphoprotein (PO) cDNA as control. The membrane was then washed four times with $2 \times$ SSC, 0.1% SDS solution at room temperature for 5 min and three times with $0.2 \times$ SSC, 0.1% SDS solution at 55 °C for 30 min and exposed to biomax film.

After hybridization with the zfER α probe, the membrane was kept at -80°C for 1 month to decrease the radioactivity and stripped (incubation for 2 h at 65°C with 75% formamide, 10 mM NaH_2PO_4 solution) in order to rehybridize with the zfER β 2 probe. The same procedure was used for PO and zfER β 1 probes.

Primer extension analysis

The probe synthesis was carried out according to the protocol described previously (Flouriot *et al.* 1996b). Briefly, a biotinylated single-stranded DNA template was used to prepare a labeled probe by extension from a specific primer by the T7 DNA polymerase in the presence of [α - ^{32}P]dCTP. To generate the probe, the vector containing the zfER α cDNA was used to obtain biotinylated PCR product (346 bp) with a biotinylated primer 1 (5'-atgtaccctaaggaggagcacagcg-3') beginning at the ATG of the exon 2 and a non-biotinylated primer 2 (5'-cctgctgagaggacaccaca-3'). After purification, the biotinylated PCR product was bound to streptavidin-coated magnetic beads (Dyna, Great Neck, NY, USA) and non-biotinylated single-strand DNA was eluted in 0.1 M NaOH. The labeled probe (320 bp) was obtained by extension in the presence of [α - ^{32}P]dCTP from internal primer 3 of exon 3 (5'-tggctcagatcgaggacag-3'). The probes were then eluted with an alkaline solution and purified on a 4% denaturing polyacrylamide-urea gel. Approximately 2×10^5 c.p.m. of single-strand probe were coprecipitated with 30 μg RNA (RNA from E2-treated zebrafish liver or yeast total RNA as control) and resuspended in 20 μl hybridization buffer. The templates were denatured at 80°C for 10 min and incubated at 50°C overnight. After an ethanol precipitation of RNA/probe hybrids, reverse transcription was carried out at 42°C for 1 h with 50 U Expand reverse transcriptase (Boehringer Mannheim, Mannheim, Germany). RNA matrix was digested at 37°C for 30 min with RNase A and EDTA (0.1 M). After purification and denaturation, the samples were separated through a 4% denaturing polyacrylamide-urea gel.

Cloning of zfER α promoter

To clone the zfER α promoter, we generated a genomic DNA fragment corresponding to exon 1 (+15), intron 1 and exon 2 (+426). This zebrafish

genomic DNA fragment (1181 bp) was amplified by PCR with a forward primer within exon 1 (5'-AGTCAGAGATACATCAGTGAGAG-3') and a reverse primer within exon 2 (5'-GCGCTGTGCTCCTCCTTAG-3'). This fragment was subcloned and sequenced.

The zebrafish genomic library inserted in $\lambda\text{gt}10$ EMBL 3 SP6/T7 vector was obtained from Clontech (Palo Alto, CA, USA). Recombinants phages (5×10^5) were screened with a ^{32}P -labeled probe corresponding to exon 1, intron 1 and exon 2 (corresponding to the 1181 bp genomic DNA fragment). A positive clone, $\lambda 2021$, containing a 7 kb genomic fragment was purified after two to three rounds of screening. The fragment was subcloned into the BamHI site of Bluescribe vector (Stratagene, La Jolla, CA, USA) and the promoter region restricted to 1.8 kb upstream of the initiation site was inserted into the KpnI/BglII site of pGL2-basic vector containing the luciferase reporter gene. This construct, named PA-1.8 kb, was sequenced on both strands by the PRISM ready reaction big dye terminator cycle sequencing protocol (PE Biosystems, Courtaboeuf, France) and the putative transcription factor binding sites were analysed by the MatInspector program (Quandt *et al.* 1995).

Plasmid constructions and site-directed mutagenesis

PA-0.3 kb construct was obtained from the PA-1.8 kb vector by deletion of the 1500 bp upstream of the AP4 box using the NcoI restriction site. The PA-0.23 kb, PA-ERE, PA-EREm1 and PA-AP-1:½ERE constructs, corresponding to various lengths of the 5' flanking sequence of the zfER α gene, were obtained by PCR from the PA-1.8 kb vector. The forward primers (A, B, C, D) and the reverse primer (Rev) contained the BamHI or SmaI restriction site and are described in Table 1. PCR products were subcloned in SmaI/BglII sites of the pGL2-basic vector (Promega, Madison, WI, USA). The QuickChange site-directed mutagenesis kit from Stratagene was used for ½ERE and 3' ERE half-site mutations of the PA-0.23 kb and PA-EREm2 constructs. The primers used are also described in Table 1. Finally, each construct was sequenced on both strands by the PRISM ready reaction big dye terminator cycle sequencing protocol.

Table 1 Oligonucleotides used for plasmid constructions and site-directed mutagenesis

	Constructs	Sequences
Primers		
A	PA-0.23 kb	TCCCCCGGGATCAAGCGGTGACCTCCTAT
B	PA-ERE	TCCCCCGGGCTGGTTGCCATGACCTGCT
C	PA-EREm1	TCCCCCGGGCT aa TTGCCATGACCTGCTC
D	PA-AP1:½ERE	TCCCCCGGGGCTCTGAGAAGTGACCGTCAG
Rev	All	CGCGGATCCGTTCACTCCTCTGATGTTTTAC
QC-0.23up	PA-0.23 kbm	GGGATCAAGCGGTGA aa TCCTATCTCTTGTTTACCTGG
QC-0.23do	PA-0.23 kbm	CAAGAGATAGG Att TCACCGCTTGATCCCGGGGG
QC-EREup	PA-EREm2	GGGCTGGTTGCCATGA aa TGCTCTGAGAAGTGACC
QC-EREdo	PA-EREm2	CTCAGAGCAT tt CATGGCAACCAGCCCCGGGG

Italic letters correspond to enzyme restriction site *Sma*I and *Bam*HI. Responsive elements are underlined. Nucleotide mutations are noted by small bold letters.

Cell culture and transfection experiments

CHO, Hela, and HepG2 cells were maintained at 37 °C in a 5% CO₂ atmosphere in Dulbecco's modified Eagle's medium (DMEM; Sigma, St Louis, MO, USA) supplemented with 5% fetal calf serum (FCS; Life Technologies, Carlsbad, CA, USA), 100 U/ml penicillin, 100 mg/ml streptomycin and 25 mg/ml amphotericin (Sigma). CHO and Hela cells were seeded in 24-well plates (5 × 10⁵ cells/well) or in six-well plates (30 × 10⁴ cells/well) for HepG2 cells. After 24 h, the medium was replaced by fresh phenol red-free DMEM-F12 containing 2.5% charcoal/dextran FCS and ethanol (0.1%) with or without E2 (10^{−8} M). Cells were transfected with plasmid using FuGENE 6 reagent as indicated by the manufacturer (Boehringer Mannheim). The DNA templates for CHO and Hela transfections contained 25 ng expression vector (with or without the coding region of each *z*fER), 150 ng luciferase reporter gene vector and 50 ng internal β-galactosidase control vector. For HepG2 transfection assays, the DNA templates contained four times more vector. The luciferase activities were assayed 36 h later using a luciferase assay system (Promega). The β-galactosidase activity was used to normalize transfection efficiency in all experiments.

Results

E2 upregulates *z*fERα mRNA expression but not *z*fERβ forms

Twenty-four adult zebrafish were divided into two groups and treated for 48 h either with 10^{−8} M E2

or the vehicle (ethanol 0.1%). The livers from each group were mixed and poly(A)⁺-RNA was prepared. Figure 1 shows the results of Northern blot experiments using each *z*fER probe. Hybridization with *z*fERα probe showed a single band of 4.8 kb which was clearly upregulated after E2 treatment. The *z*fERβ1 probe revealed a strong signal band at 4 kb in the liver of control animals. Interestingly, this mRNA was markedly downregulated after E2

Figure 1 Northern blot analysis of ER mRNAs in the liver of untreated and E2-treated zebrafish. Poly(A)⁺-RNA (1.5 µg), from the liver of zebrafish exposed for 48 h to ethanol (−) or 10 nM E2 (+), was separated on a denaturing formaldehyde–agarose gel and transferred to a nylon membrane. This membrane was successively hybridized under stringent conditions with ³²P-labeled *z*fERα, *z*fERβ1, *z*fERβ2 and PO probes prior to autoradiography. The size of *z*fER mRNAs was determined from standard markers and the position of ribosomal RNA stained by methylene blue is indicated on the right.

treatment. Hybridization with the zfER β 2 probe showed a more complex expression pattern. A major signal was detected at 7.5 kb and two lower bands were also observed at 2.8 kb and 1.4 kb. The amounts of these mRNAs were not modified after E2 treatment.

Characterization of the promoter region and transcription initiation start site of the zfER α gene

To investigate the molecular mechanisms involved in the E2 upregulation of zfER α mRNA, we cloned the promoter region of the zfER α gene. A genomic DNA library was screened using a probe spanning exon 1 and exon 2. A sequence of 1.8 kb at the 5' end of the zfER α gene was isolated and sequenced. First, to identify the transcription initiation site, a primer extension experiment on total RNA from liver was carried out (Fig. 2). The size of the major product of the primer extension assay was determined using the accompanying sequence ladder and was localized 139 bp upstream of the ATG of exon 2 (Fig. 2). Staining of other lower extension fragments was also detected and corresponded to minor initiation start sites.

In a second step, the presence of presumptive transcription factor DNA-binding sites was determined using computer analysis (Fig. 3). The identified promoter sequence was composed of 1745 bp. The major initiation start site is represented by an arrow. The promoter has no TATA or CAAT box, but presents an Inr sequence at -50 bp (Javahery *et al.* 1994). Two C-Ets-1 sites at -1625 and +8bp and two C/EBP- β sites at -1280 and -150 bp were found. Four GATA potential binding sites were also located (three GATA-1 sites at -1744, -785 and +17 bp; GATA-2 or 3 at -1321 bp). Brn2 and SREBP potential binding sites were identified at -1708 and -1265 bp respectively. Three AP-1 potential binding sites, which could include a $\frac{1}{2}$ ERE, were located at -850, -129 and -80 bp. In the proximal part of the promoter region, an imperfect ERE was found at -104 bp. AP-1 and AP-4 sites were also located at -605 and at -200 bp.

E2-dependent upregulation of the zfER α gene uses an imperfect ERE

To determine if the transcription rate of the zfER α gene was altered directly by ERs under E2

Figure 2 Identification of the transcription initiation start sites of the zfER α gene.

Labeled probe, extending from exon 2 to exon 3, was hybridized with 30 μ g total RNA (tRNA) from E2-treated zebrafish liver or with yeast tRNA as control. The major extension product signal is noted by a thicker arrow and the lesser stainings by the other arrows. The extension product sizes were determined using the accompanying sequence ladder. The differently initiated mRNA are schematically represented on the left.

treatment, the zfER α gene promoter (PA-1.8 kb) was used as the reporter gene and was cotransfected with the zfER α expression vector into CHO cells. After 36 h of E2 treatment (10^{-8} M), a significant E2-dependent induction (approximately 10-fold) of the PA-1.8 kb reporter was observed (Fig. 4). Because this result showed that the zfER α gene was induced by zfER α itself, we next examined whether DNA potential sites, particularly the imperfect ERE described above at -104 bp by computer analysis, could be involved. To address

-1745 CTCGAGGATA TTACGAGCTC AGGGCTCTCT CCCGGGAG CATGCCAAAT ACGCTTTATT GATTATCAGC TAAGTG TGAA
GATA-1 *Brn2*
 -1665 CTCTTGAAAG CTTATTAATG AGCATGAAAA CAGGCCATAG TCAGGACGAT GTGTA CTGCGTCTAT
C-Ets-1
 -1585 TTTGAAAATT GAATAAAATA TGGGAAGATT TCTTTTGTTT TTTTTTTTTA ACTTAATTCC ACTGCTTATT TCTGTATAAA
 -1505 TCTATTAATA CATTTTCTCT TCCCAAATCA TACAAATAAC AAATTGTCAA AGTTTCTCTG ATGCATTTTA TAGGATTTTT
 -1425 TTTTAGTTCT GTGTACACAC ACACACACAC ACACACACAC GTATGTATTT ATGTATATAT CTATGTATTT GCAAGGAGCT
 -1345 TATTGTTCTA TCTGTTCTAG AGAGTAAGAG ATAAGAGTAA AGAACCCCA AAAGCACAAA TCATTAATGT TCTGCAATTA
GATA-2 and 3 *C/EBP*
 -1265 TCACGTGAAC GCTATCTTTG TGTGTTTGGT GTCAGTTGAG GACTCTGGCC AAGCTGTTTT TTTTGGAAA TCAAGGCCAT
SREBP
 -1185 GTTTTTGGTC TTTGAACGTG ACTAGTTCTA GCTCTGTTCT GGTTAAGCGT AGCTCAAGCA TACACTAAA ATGCACGTGT
 -1105 TTAATAAAT AATAGTGTG TATATCTAAA ATGTGTTTTT TAAGAACCTT GCAGGAGCAG ATTGAGACTT AATTGAGCCA
 -1005 AAAGGACACA GACATCTGCT TGCATAAAGG CTATTAATAC AAAATGCTCT ATTTAGGAAA AGAACATTAC TATTTTAGAT
 -945 TTGTAAGCTG CAAACATGA ATACACACAT CAAATGAGC CATTAGCAGT AATTATATTT GAGGAAAGAA TTGTGTTTTT
 -865 TTTTATTAAC ACGGCTAATG ACCAATAAAA AAAAATTGT GATCATTAGA AATTCTGTTT TTACCACCAC TATGTTCAAT
1/2ERE AP-1
 -785 ATATGATATA TATTTTAAA TGATTACAA TATATTTTAT CTGTATTTAT TTGACAGTAA AAACCAATTA AATATATTTA
GATA-1
 -705 AAACCATTTA ATGTGAAACA GTATTAAAT GTAAATGTTT TTCTATTTTG ATTTTTTTTC AAATATTTTA TTAGTCACAA
 -625 GAATATATAT TAATATGCAT ACTGACAGC AACGTAAAG TCAATAATT ATCAATGTCT ACTCAGTTAT TTGAAATATT
AP-1
 -545 TATTTCAATG CAAACGTTTT CACTGCCTTT TCTAATCAAA TCAATTCTTT TACTCAAATC TAATCAACTC TTGCTGAAAA
 -465 AAAGTGTCAG TTTATTTTAT TGTTCACAA GACTGTATA TTTAAATCCA TAGTCAGTCT AATGAAGTTT CCCTAATCAA
 -385 ACAAAAAGCA CAACACCCAC ACATGTTCCA TTCTGTGTA AAATAGCTGC CATTACAGAG AAATGATTAT AATCAACACG
 -305 ATTTGTTTTG TCCCTAGAGA GCTGAACAGC GTCTCCTTCC CTTTTAACC TCCTGCAAGC ATCTTCCAAC GAATTTAATT
 -225 ACTCCTTTGA CATATAATTT CCCATGGCAG CAGCAATGTAA AGTGGTTTCA CAGCGCATCA CCTGTAAAC TCAAGGTTT
AP-4
 -145 TGGCAAGTGA ATCAAGCGT GACCTCCTAT CTCTTGTTTA CCGGTTGCC ATGACCTGCT CTGAGAGCTG ACCGTCAGTA
C/EBP *1/2ERE AP-1* *ERE* *1/2ERE AP-1*
 -65 TCGACCTGGC TGTCTTTAAT TTSTTTACCT CCTCTTTTTT TAAGCCAGAG ACACAGACAG AGAAAGTAAG GACACAGGAA
hr *+1* *C-Ets-1*
 +15 GTCAGAGATA CATCAGTGAG AGAGAGAAAG CATCCAGCCT GTAATGGGAC TCAAGTAAAA CATCAGAGGA GTGAACATTT
GATA-1
 +95 TGGAAGGAT GCTGGAAAGA ATTCAGCAAC CAATTTTTC GCTTTTGTTT TTGTAGTAAT GAAATTCAT GTGTCTTTCA
 CTCTAGCGTT TTGTAATCGA GCATTGTTTC TATTATTAGT GCTTTCAGTT GAGTTTCCGT TCCTGTCCGT TTCTGTGTTT
 ATAGAAAAGC ACACCTTTCA GTTGGGTTTA TCCTTTCTTA CACTGCAAT AATAATGCTT TTTGTCTGT GTCTAGTCCA
 AATACCTAAC AATTATCTAA AAAAATGATG TAAAAATACT GTTTTGTGT TGTGTTGTCC CTTAAAAATA ATCTGCCAAT
 GAAGTGGGTA AATAAATCTT AATTTCCGGT TGAAATAAGA TCATTTTAAA ATCATTTTTG TTGTTTGACA TAAATCTCC
 CTCAATTTTG ACATTATTTT TGTTCAAAG TAAACAAAA CAATTTTTT TACTGTGCTA GAAATGCTT CCTGAATGCT
 CTTTCTTCC AAAAAAAAAA AAAAAAAAAAG CAGCTTCAAG AAGTCAGAAA TGCAATTTCTT TGCAGTGTAA ATGTAATGCA
 TAAGACATTT GGGGAAGAA GAACAATGAG GGGGAAAAGA TGGGATAAAA GAAAGGCAAA AGAATCAAAA TGAAATGTG
 TTTACAGAGG GAAAGACAGG AAGAAAGAGA AAGAAAAGCA GTCTCAGCTA TCTTAGAAAA ACATCATACT TTGTCCTCTC
 AGCCCAAGGG GGACTGTAGG AGAGGAAGGT AGTTTTTTT TAGGAGGAGT AGGCAGAGCA GCGAGAGAGT GAAGCGGAAG
 +137 CGTAGCAGAA CAAGGGGGAG AGAGTTTGAG AAAGACTGTG GGAGGTGAA AAAAAGAAGC CTGAGCTTAA AAAAGACAGA
 +207 AACAGTCTTT TGCTTATGAA CAGATAGTAC CCTAAGTCAT GCTTGCTGCG TGTGTGTTTC CCATGGTGAT GTCTGGAGGG
 +287 CAGACCAGCG GAGAGGCTGT TGGTGCCAGG CAGCGACGCA GGACCAGCCC GAGTCCTGAG AGAGAAGCCC TGGAGGAACC
 +367 TTCTCTGCCA CCTGCCGCC ACAAACCTCT ACCCATGTAC CCTAAGGAGG AGCAGAGCGC
 *

Figure 3 Sequence analysis of the 5' end region of the zER α gene. The putative DNA-binding sites for several transcription factors were determined with the MatInspector program (Quandt *et al.* 1995) and are boxed. The major initiation start site is noted by an arrow. Exon 1 is identified by bold letters and the italic letters correspond to intron 1. The underlined sequence corresponds to exon 2. The potential ATG codon is noted by an asterisk.

Figure 4 A complex unit containing 1/2ERE-ERE is involved in the E2-dependent upregulation of the zfERα gene. E2-treated or untreated CHO cells were cotransfected with an expression vector (empty or containing the coding region of zfERα) and several constructs of the zfERα gene promoter described at the left. Mutations on the half site and imperfect ERE are indicated by asterisks. On the right, data are expressed as percentage of activity versus the E2 induction (±S.E.M.) of PA-1.8 kb. Each experiment was repeated at least twice in triplicate.

this question, we compared E2 induction of several luciferase constructs versus the E2 induction of PA-1.8 kb.

Figure 4 shows that, compared with the PA-1.8 kb construct, the PA-0.3 kb vector was less inducible by zfERα (65% of E2-PA-1.8 kb) indicating that several binding sites, notably the CEBP-β and AP-1:1/2ERE, located between -1280 and -850 bp, could be involved in the E2-dependent zfERα gene induction. Curiously, E2 induction of the PA-0.23 kb construct was higher (85% of E2-PA-1.8 kb) than PA-0.3 kb (65% of E2-PA-1.8 kb), suggesting that a silencer element, unidentified by computer analysis, could exist between -205 and -135 bp.

Compared with PA-0.23 kb, E2 induction was significantly diminished when the AP-1:1/2ERE located at -129 bp was mutated (PA-0.23 kbm) or entirely deleted (PA-ERE), demonstrating that this

half site is involved in the E2-dependent induction of the zfERα gene (Fig. 4). To investigate the role of the imperfect ERE (-104 bp), two different mutations were performed. Figure 4 shows that the stimulatory effect of E2 on the PA-ERE promoter was reduced from 50% to 25% of E2-PA-1.8 kb activity by mutation of the ERE in the 5' half site (PA-EREm1). Interestingly, this induction was completely abolished with the mutation in the 3' half site (PA-EREm2). Moreover, the deletion of this ERE, PA-API:1/2ERE construct, also showed no obvious induction. These data confirmed the importance of this imperfect ERE for the positive regulation of the zfERα gene. The comparison of the zfERα promoter region with the proximal promoter of several estrogen-sensitive fish genes highlights a complex unit containing a 1/2ERE close to an ERE presenting the same structural organization (Fig. 5).

Figure 5 Alignment of the proximal promoters from some estrogen-sensitive fish genes. Proximal promoters of the ER α and brain aromatase genes from several fish species (zf, zebrafish; gf, goldfish; rt, rainbow trout) were aligned with the GeneJockey II program. Putative EREs, as well as Inr sequences, are shaded. bp Positions were calculated from the transcriptional start site when it had been determined. The relative position of the three 1/2 EREs on the DNA double helix is schematically illustrated below.

zfER β 2 but not zfER β 1 upregulates the zfER α gene

In order to investigate the E2-dependent transcriptional capacity of the two zfER β s on the zfER α gene, each expression vector (empty or containing the different coding region of each zfER) was cotransfected with PA-1.8 kb into CHO (Fig. 6A), Hela (Fig. 6B) and HepG2 cells (Fig. 6C). E2 stimulation of the reporter gene mediated by zfER α reached 10-fold in CHO, 6.5-fold in Hela and 5-fold in HepG2. When using the zfER β 2 expression vector, the E2 stimulation of the reporter gene was 50–60% lower with 3- to 4-fold induction in CHO and in Hela cells and 2.5-fold in HepG2 cells. Surprisingly, irrespective of the cell line used, no obvious E2 induction of the zfER α promoter was observed when using the zfER β 1

expression vector. Nevertheless, this form clearly induced the ERE-TK-luc construct under the same experimental conditions (data not shown) (Menuet *et al.* 2002).

Discussion

The egg yolk protein vitellogenin can be activated *de novo* from a totally silent state within hepatocytes by estrogens. In oviparous species, this process of hepatic vitellogenin is tightly coupled to a clear E2-dependent upregulation of ER α gene expression (Pakdel *et al.* 1991, Flouriot *et al.* 1996a). We have recently cloned and characterized two zfER β forms which co-express with zfER α in the liver of zebrafish (Menuet *et al.* 2002). In this study, we have investigated the responsiveness of these zfER

Figure 6 zfER α and zfER β 2, but not zfER β 1, induce the zfER α gene promoter. (A) CHO, (B) Hela and (C) HepG2 cells were cotransfected with different expression vectors (control (ctrl), zfER α , zfER β 1 or zfER β 2) and the PA-1.8 kb reporter gene. Cells were treated with 10 nM E2 (solid bars) or 0.1% ethanol as control (open bars). The histograms show the mean of fold induction versus ethanol (etoh) ($\pm$ S.E.M.) for each reporter gene. Each experiment was repeated at least twice in triplicate.

genes to *in vivo* E2 treatment and studied the potential effects of these zfER β forms on zfER α gene regulation.

Nothern blot analysis of zebrafish liver revealed one single major band of 4.8 kb for zfER α , one band of 4 kb for zfER β 1 and three bands of 7 kb, 2.8 kb and 1.4 kb for zfER β 2. Since the entire coding region of zfER β 2 extends over 1.8 kb (Menuet *et al.* 2002), one can assume that the 1.4 kb transcript would generate a truncated zfER β 2 protein isoform. Similarly, small transcripts for ER β s were also found in the liver of other teleost fish (Xia *et al.* 2000, Socorro *et al.* 2000). It would therefore be interesting to investigate whether or not these transcripts could generate truncated protein and to seek their potential roles.

Interestingly, exposure of zebrafish to E2 showed a differential response of zfER genes. Our data confirmed that in the zebrafish, as in other oviparous species, the expression of the zfER α gene is robustly stimulated by E2 treatment *in vivo*. In contrast and surprisingly, the expression level of zfER β 1 was strongly reduced whereas the level of zfER β 2 mRNA was either not affected or very slightly affected. At the present time, the transcriptional and/or post-transcriptional effects of E2 treatment on these messengers remain to be determined. However, the present results may indicate that, during the reproductive cycle, when E2 levels fluctuate, the zfER β forms could have different patterns of expression, suggesting a distinct implication on hepatic functions, notably on vitellogenesis.

To investigate the effect of zfER β s on this function, we analysed *in vitro* their potential actions on the regulation of the zfER α gene expression known to be essential for vitellogenin production. First, a zebrafish genomic DNA library was screened and a 7 kb DNA fragment was isolated. This fragment contained 1.8 kb of the promoter region and the first exons and introns of the zfER α gene. Interestingly, this genomic fragment showed a similar organization and structure to that of the corresponding fragment in the rainbow trout ER α gene. In fact, two classes of ER α mRNAs could be generated from the trout ER α gene by an alternative splicing and promoter usage (Leroux *et al.* 1993, Pakdel *et al.* 2000). These transcripts encode two functional ER α isoforms with different estrogen dependencies. The major hepatic isoform, rtER α short, was initiated from the ATG located in

exon 2, while a minor isoform, rtER α long, was raised from an in-frame ATG initiator codon located in an intronic sequence of intron 1 (called exon 2a) which can be differentially spliced (Pakdel *et al.* 2000). As expected, our primer extension analysis showed that the major zfER α isoform corresponded with the classical short isoform issued from transcripts containing exon 1/exon 2. However, the presence of an additional in-frame ATG initiator codon in intron 1 (or exon 2a) located at the same position as in the rtER α gene suggested that a zfER α long isoform also exists in zebrafish.

From the genomic DNA, we have isolated 1.8 kb of the 5' end of zfER α gene and linked it to the luciferase reporter gene. Cotransfection analysis of this promoter construct with the zfER α expression vector in different cell lines showed a clear induction of the reporter gene in the presence of E2. These results confirmed the E2 stimulation of the zebrafish ER α gene *in vivo* and demonstrated that E2 upregulation of the ER α gene by ER α itself is likely widespread among oviparous vertebrates (Pakdel *et al.* 1991, Ninomiya *et al.* 1992, Le Drean *et al.* 1995, Lee *et al.* 1995). Mutations and deletions of this promoter region showed clearly that the $\frac{1}{2}$ ERE (−127 bp) and the imperfect ERE (−104 bp) were largely involved in the E2-dependent induction of the zfER α gene. The comparison of the proximal promoter region of some fish estrogen-sensitive genes (ER α and brain aromatase) showed the conserved presence of $\frac{1}{2}$ ERE and ERE, separated by 21–23 bp (center-to-center). This number corresponds to two helix turns, indicating that these two elements are on the same side of the DNA (Fig. 5). This organization could stabilize receptor binding and enhance E2 induction. However, for the zfER α gene, we cannot exclude that the $\frac{1}{2}$ ERE located at −850 bp and −80 bp could also contribute to the E2 responsiveness. Moreover, computer analysis revealed that the $\frac{1}{2}$ EREs are included in an AP-1-like site. Numerous studies showed that ERs are able to modulate promoter activity by interacting with different DNA-bound transcription factors such as Jun and Fos which bind specifically to the AP-1 element (Paech *et al.* 1997, Webb *et al.* 1999). In this regard, more investigations will be necessary to determine if AP-1-like elements are also involved in the E2-dependent induction of the zfER α gene.

To investigate the ER β effects on the zfER α promoter, transfection experiments were carried out in several cell lines including CHO, Hela and HepG2. Curiously, the two ER β forms were characterized by differential impacts on the E2-dependent induction of the zfER α construct. In fact, although clear inductions were obtained with zfER α , zfER β 2 seemed to be less effective (about 40–50% of zfER α activity). In contrast, whatever the cell line used, zfER β 1 showed very low, if any, transcriptional activity on the zfER α promoter. The molecular reasons for this low transcriptional activity are still unknown. However, these data contrast with previous results showing that zfER β 1 is able to induce reporter genes under the control of a consensus ERE in the same way as zfER α and zfER β 2 (Bardet *et al.* 2002, Menuet *et al.* 2002). Besides the fact that the ERE of the zfER α gene is imperfect with three mutations, it is possible that the inefficiency of zfER β 1 to induce zfER α promoter is due to its incapacity to recognize the zfER α ERE with high affinity. Gel mobility shift assays carried out with labeled oligonucleotides containing this imperfect ERE sequence did not allow us to obtain information about the *in vitro* DNA-binding capacity of these different zfERs. We were limited by the low sensitivity of this technique due to the high degree of mutation in this ERE sequence. Nevertheless, our assays performed in parallel with a consensus ERE showed low, but detectable, specific complexes, with higher intensity for zfER α than both zfER β s (data not shown). Similar to our finding, previous studies have reported that mammalian ER β s have lower transcriptional activity than ER α (Tremblay *et al.* 1997, Cowley & Parker 1999). At the present time, it is difficult to know which one of the zfER β s functions like the mammalian ER β . This will necessitate a full comparison using a series of endogenous E2-sensitive genes. Moreover, these experiments may provide information about potential EREs that could favor ER β transcriptional activity preferentially.

Although numerous studies have demonstrated that in oviparous species ER α gene regulation involves several nuclear factors including ER α itself, COUP-TFI (Lazennec *et al.* 1997, Petit *et al.* 1999, Métivier *et al.* 2002), C/EBP β and the glucocorticoid receptor (Lethimonier *et al.* 2000, 2002), the present study has revealed that the zfER β forms, and more probably zfER β 2,

could also be largely involved. Interestingly, zfER β 2 has been characterized by a better affinity for E2 than the other forms (Menuet *et al.* 2002). Furthermore, the present data revealed that zfER β 2 expression is unmodified by E2 and that it is able to induce zfER α promoter activity significantly. Consequently, it is tempting to speculate that, *in vivo*, zfER β 2 could be involved in the maintenance of zfER α gene expression when the E2 levels are low during the reproductive cycle. Moreover, the fact that, in contrast with zfER α , zfER β 1 expression is downregulated by E2 treatment and unable to significantly induce zfER α promoter activity *in vitro*, suggests that this zfER form is likely not involved *in vivo* in zfER α gene regulation and in the vitellogenesis process.

In conclusion, this study has shown for the first time that the two ER β forms that have emerged in teleost species are differentially regulated by E2 in the liver and characterized by distinct transcriptional activity on an endogenous gene such as zfER α . These data may also suggest that each ER β form has different physiological functions that remain to be elucidated. Moreover, the isolation of the zfER α promoter and the demonstration of its sensitivity to estrogen provide new molecular tools in the zebrafish, a species commonly used to analyze the impact of endocrine disrupters in a whole vertebrate model organism.

Acknowledgements

This work was supported by the European Community (project QLK4-CT-2002-00603-EDEN), the French Ministry for Environment (Programme Environnement et Santé, subvention no. 01108), ACI (Biologie du Développement et Physiologie Intégrative), CNRS and fellowships from the French Ministry for Education, Research and Technology to A M.

References

- Bardet PL, Horard B, Robinson-Rechavi M, Laudet V & Vanacker JM 2002 Characterization of oestrogen receptors in zebrafish (*Danio rerio*). *Journal of Molecular Endocrinology* **28** 153–163.
- Cowley SM & Parker MG 1999 A comparison of transcriptional activation by ER alpha and ER beta. *Journal of Steroid Biochemistry and Molecular Biology* **69** 165–175.

- Flouriot G, Pakdel F & Valotaire Y 1996a Transcriptional and post-transcriptional regulation of rainbow trout estrogen receptor and vitellogenin gene expression. *Molecular and Cellular Endocrinology* **124** 173–183.
- Flouriot G, Nestor P, Kenealy MR, Pope C & Gannon F 1996b An S1 nuclease mapping method for detection of low abundance transcripts. *Analytical Biochemistry* **237** 159–161.
- Green S, Walter P, Kumar V, Krust A, Bornert JM, Argos P & Chambon P 1986 Human oestrogen receptor cDNA: sequence, expression and homology to v-erb-A. *Nature* **320** 134–139.
- Hawkins MB, Thornton JW, Crews D, Skipper JK, Dotte A & Thomas P 2000 Identification of a third distinct estrogen receptor and reclassification of estrogen receptors in teleosts. *PNAS* **97** 10751–10756.
- Javahery R, Khachi A, Lo K, Zenzie-Gregory B & Smale ST 1994 DNA sequence requirements for transcriptional initiator activity in mammalian cells. *Molecular and Cellular Biology* **14** 116–127.
- Jobling S, Casey D, Rodgers-Gray T, Oehlmann J, Schulte-Oehlmann U, Pawlowski S, Baunbeck T, Turner AP & Tyler CR 2003 Comparative responses of molluscs and fish to environmental estrogens and an estrogenic effluent. *Aquatic Toxicology* **65** 205–220.
- Klinge CM 2000 Estrogen receptor interaction with co-activators and co-repressors. *Steroids* **65** 227–251.
- Kuiper GG, Enmark E, Pelto-Huikko M, Nilsson S & Gustafsson JA 1996 Cloning of a novel receptor expressed in rat prostate and ovary. *PNAS* **93** 5925–5930.
- Kuiper GG, Carlsson B, Grandien K, Enmark E, Haggblad J, Nilsson S & Gustafsson JA 1997 Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology* **138** 863–870.
- Lassiter CS, Kelley B & Linney E 2002 Genomic structure and embryonic expression of estrogen receptor beta (ERbeta a) in zebrafish (*Danio rerio*). *Gene* **299** 141–151.
- Lazennec G, Kern L, Valotaire Y & Salbert G 1997 The nuclear orphan receptors COUP-TF and ARP-1 positively regulate the trout estrogen receptor gene through enhancing autoregulation. *Molecular and Cellular Biology* **17** 5053–5066.
- Le Drean Y, Lazennec G, Kern L, Saligaut D, Pakdel F & Valotaire Y 1995 Characterization of an estrogen-responsive element implicated in regulation of the rainbow trout estrogen receptor gene. *Journal of Molecular Endocrinology* **115** 37–47.
- Lee JH, Kim J & Shapiro DJ 1995 Regulation of *Xenopus laevis* estrogen receptor gene expression is mediated by an estrogen response element in the protein coding region. *DNA Cell Biology* **14** 419–430.
- Le Guevel R & Pakdel F 2001 Assessment of oestrogenic potency of chemicals used as growth promoter by *in vitro* methods. *Human Reproduction* **16** 1030–1036.
- Le Roux MG, Theze N, Wolff J & Le Pennec JP 1993 Organization of a rainbow trout estrogen receptor gene. *Biochimica et Biophysica Acta* **1172** 226–230.
- Lethimonier C, Flouriot G, Valotaire Y, Kah O & Ducouret B 2000 Transcriptional interference between glucocorticoid receptor and estradiol receptor mediates the inhibitory effect of cortisol on fish vitellogenesis. *Biology of Reproduction* **62** 1763–1771.
- Lethimonier C, Flouriot G, Kah O & Ducouret B 2002 The glucocorticoid receptor represses the positive autoregulation of the trout estrogen receptor gene by preventing the enhancer effect of a C/EBPbeta-like protein. *Endocrinology* **143** 2961–2974.
- Ma CH, Dong KW & Yu KL 2000 cDNA cloning and expression of a novel estrogen receptor beta-subtype in goldfish (*Carassius auratus*). *Biochimica et Biophysica Acta* **1490** 145–152.
- McLachlan JA 2001 Environmental signaling: what embryos and evolution teach us about endocrine disrupting chemicals. *Endocrine Reviews* **22** 319–341.
- Menuet A, Pellegrini E, Anglade I, Blaise O, Laudet V, Kah O & Pakdel F 2002 Molecular characterization of three estrogen receptor forms in zebrafish: binding characteristics, transactivation properties, and tissue distributions. *Biology of Reproduction* **66** 1881–1892.
- Métivier R, Gay FA, Hubner MR, Flouriot G, Salbert G, Gannon F, Kah O & Pakdel F 2002 Formation of an hER alpha-COUP-TFI complex enhances hER alpha AF-1 through Ser118 phosphorylation by MAPK. *EMBO Journal* **21** 3443–3453.
- Mosselman S, Polman J & Dijkema R 1996 ER beta: identification and characterization of a novel human estrogen receptor. *FEBS Letters* **392** 49–53.
- Ninomiya Y, Mochii M, Eguchi G, Hasegawa T, Masushige S & Kato S 1992 Tissue-specific response of estrogen receptor gene expression to estrogen in chick. *Biochemical and Biophysical Research Communications* **187** 1374–1380.
- Paech K, Webb P, Kuiper GG, Nilsson S, Gustafsson J, Kushner PJ & Scanlan TS 1997 Differential ligand activation of estrogen receptors ERalpha and ERbeta at AP1 sites. *Science* **277** 1508–1510.
- Pakdel F, Feon S, Le Gac F, Le Menn F & Valotaire Y 1991 *In vivo* induction of hepatic estrogen receptor mRNA and correlation with vitellogenin mRNA in rainbow trout. *Molecular and Cellular Endocrinology* **75** 205–212.
- Pakdel F, Metivier R, Flouriot G & Valotaire Y 2000 Two estrogen receptor (ER) isoforms with different estrogen dependencies are generated from the trout ER gene. *Endocrinology* **141** 571–580.
- Petit F, Le Goff P, Cravédi JP, Valotaire Y & Pakdel F 1997 Two complementary bioassays for screening the estrogenic potency of xenobiotics: recombinant yeast for trout estrogen receptor and trout hepatocyte cultures. *Journal of Molecular Endocrinology* **19** 321–335.
- Petit F, Métivier R, Valotaire Y & Pakdel F 1999 Synergism between a half site estrogen-responsive-element (ERE) and an imperfect ERE and cooperation with COUP-TFI are required for estrogen receptor (ER) to achieve a maximal estrogen-stimulation of rainbow trout ER gene. *European Journal of Biochemistry* **259** 385–395.
- Quandt K, Frech K, Karas H, Wingender E & Werner T 1995 MatInd and MatInspector: new fast and versatile tools for detection of consensus matches in nucleotide sequence data. *Nucleic Acids* **23** 4878–4884.
- Saville B, Wormke M, Wang F, Nguyen T, Enmark E, Kuiper G, Gustafsson JA & Safe S 2000 Ligand-, cell-, and estrogen receptor subtype (alpha/beta)-dependent activation at GC-rich (Sp1) promoter elements. *Journal of Biological Chemistry* **275** 5379–5387.
- Socorro S, Power DM, Olsson PE & Canario AV 2000 Two estrogen receptors expressed in the teleost fish, *Sparus aurata*: cDNA cloning, characterization and tissue distribution. *Journal of Endocrinology* **166** 293–306.
- Thomas PS 1980 Hybridization of denatured RNA and small DNA fragments transferred to nitrocellulose. *PNAS* **77** 5201–5205.
- Tremblay GB, Tremblay A, Copeland NG, Gilbert DJ, Jenkins NA, Labrie F & Giguere V 1997 Cloning, chromosomal localization, and functional analysis of the murine estrogen receptor beta. *Molecular Endocrinology* **11** 353–365.
- Tyler CR, Jobling S & Sumpter JP 1998 Endocrine disruption in wildlife: a critical review of the evidence. *Critical Reviews in Toxicology* **28** 319–361.
- Webb P, Nguyen P, Valentine C, Lopez GN, Kwok GR, McInerney E, Katzenellenbogen BS, Enmark E, Gustafsson JA, Nilsson S & Kushner PJ 1999 The estrogen receptor enhances AP-1 activity by two distinct mechanisms with different requirements for receptor transactivation functions. *Molecular Endocrinology* **13** 1672–1685.
- Xia Z, Gale WL, Chang X, Langenau D, Patino R, Maule AG & Densmore LD 2000 Phylogenetic sequence analysis, recombinant

expression, and tissue distribution of a channel catfish estrogen receptor beta. *General and Comparative Endocrinology* **118** 139–149.

Zilliacus J, Wright AP, Carlstedt-Duke J & Gustafsson JA 1995 Structural determinants of DNA-binding specificity by steroid receptors. *Molecular Endocrinology* **9** 389–400.

Received in final form 8 January 2004
Accepted 19 March 2004