Urocortin induces interleukin-6 release from rat cardiomyocytes through p38 MAP kinase, ERK and NF- κ B activation

Man Huang^{1,2}, Duraisamy Kempuraj², Nikoletta Papadopoulou^{1,3}, Taxiarchis Kourelis², Jill Donelan^{1,4}, Akriki Manola¹ and Theoharis C Theoharides^{1,5,6}

¹Molecular Immunopharmacology and Drug Discovery Laboratory, Department of Pharmacology and Experimental Therapeutics and ²Bowles Center for Alcohol Studies, University of North Carolina at Chapel Hill, CB# 7178, Thurston Bowles Building, Chapel Hill, North Carolina 27599-7178, USA

³Medical Faculty, Institute of Medical Microbiology, Immunology, and Hygiene, University of Cologne, Goldenfelsstraße 19-21, D-50935 Köln, Germany

⁴Millennium Pharmaceuticals, 40 Landsdowne Street, Cambridge, Massachusetts 02139, USA

⁵Departments of Internal Medicine and ⁶Biochemistry, Tufts University School of Medicine, Tufts Medical Center, 136 Harrison Avenue, Boston, Massachusetts 02111, USA

(Correspondence should be addressed to T C Theoharides; Email: theoharis.theoharides@tufts.edu)

Abstract

CRH and its structurally related peptide urocortin (Ucn) are released under stress. Ucn is a potent agonist for CRH-receptor 2 (CRH-R2), which is strongly expressed in rodent heart. Stress induces Ucn mRNA expression in the heart, where it may be cardioprotective. However, increasing evidence indicates that Ucn may also have pro-inflammatory actions. Here, we show that neonatal rat cardiomyocytes express CRH-R2 by western blot analysis and Ucn induces interleukin-6 (IL-6) release in a time- and dose-dependent fashion. Ucn stimulates activation of ERK and p38 MAP kinases, while both MEK1 and p38 inhibitor block Ucn-induced IL-6 release. Ucn also activates nuclear factor kappa B (NF- κ B) and a NF- κ B inhibitor blocks Ucn-induced IL-6 release. Finally, the CRH-R antagonists α -helical (9–41) CRH and astressin-2B completely inhibit Ucn-induced IL-6 release, as well as activation of ERK, p38, and NF- κ B. These findings indicate that Ucn induces IL-6 synthesis and release from neonatal rat cardiomyocytes. Our findings suggest that even though Ucn may confirm some protection on cardiomyocyte survival, it can also release IL-6, which is an independent risk factor for acute coronary syndrome. The precise role of cardiac Ucn *in vivo* remains to be elucidated.

Journal of Molecular Endocrinology (2009) **42**, 397–405

Introduction

The stress response is regulated by CRH, which activates the hypothalamic–pituitary–adrenal (HPA) axis typically leading to anti-inflammatory actions (Chrousos 1995). There are two main types of the G protein-coupled CRH receptors, CRH-receptor 1 (CRH-R1), and CRH-R2, which mediate the effects of the CRH family of structurally related peptides that includes CRH, urotensin, sauvagine, and the urocortins (Ucn). Ucn binding to CRH-R2 is about 40 times more potent than for CRH itself. CRH-R1 is mainly expressed in the central nervous system (CNS) and anterior pituitary corticotrophs, activation of which leads to release of adrenocorticotrophic hormone (Chrousos 1995). CRH-R2 has three different spliced forms, α , β , and γ , of which CRH-R2 α is mainly in the CNS, while CRH-R2 β is predominantly found in the periphery, especially in the heart (Nishikimi *et al.* 2000, Kimura *et al.* 2002) including cardiomyocytes (Okosi *et al.* 1998). Ucn mRNA is also expressed in the heart (Nishikimi *et al.* 2000). Species differences have been reported in the

distribution of CRH-R2 in the cardiovascular system, showing 10 $\times$ higher expression in mice than rats or humans cardiac muscle (Waser *et al.* 2006).

Ucn has been generally considered to be cardioprotective, especially in ischemia-reperfusion (IR) injury (Bale *et al.* 2003, Latchman 2003, Townsend *et al.* 2007), but is also increased (Nishikimi *et al.* 2000, Ikeda *et al.* 2003) with parallel decrease in cardiac CRH-R2 β expression in cardiac hypertrophy (Nishikimi *et al.* 2000, Ikeda *et al.* 2003). On the other hand, acute stress, CRH, and Ucn have been implicated in the pathophysiology of some neuroinflammatory disorders (Chrousos 1995, Karalis *et al.* 1997, Theoharides & Cochrane 2004) and myocardial ischemia (MI; Jiang *et al.* 1996, Krantz *et al.* 2000). Acute stress elevates plasma interleukin-6 (IL-6) levels in rodents (Ando *et al.* 1998, Nukina *et al.* 2001) and such levels are higher in apolipoprotein E (Apo E) knockout mice that develop atherosclerosis (Huang *et al.* 2003).

Local inflammation is now recognized as a key component of coronary artery disease (CAD; Packard & Libby 2008). The proinflammatory cytokine IL-6 is

thought to contribute to the development of CAD (Huber *et al.* 1999), cardiomyopathy, congestive heart failure (Deliargyris *et al.* 2000) and MI (Miyao *et al.* 1993). IL-6 is a major inducer of C reactive protein (CRP) and both are elevated in and are associated with increased risk of CAD. The Health ABC study showed that plasma IL-6 levels had a stronger association with CAD than CRP (Cesari *et al.* 2003), while the PRIME study showed that only IL-6 remained significantly associated with MI (Luc *et al.* 2003). In another study, the incidence of future acute coronary events and mortality of patients with stable CAD or healed MI was strongly correlated with serum IL-6 levels over a 6-year observation (Fisman *et al.* 2006). In patients with acute coronary syndrome (ACS), plasma IL-6 was increased and appeared to derive primarily from a cardiac source (Deliargyris *et al.* 2000), but the exact cell type remains known. Cardiomyocytes are capable of releasing IL-6 in response to hypoxia and to cytokines (Yamauchi-Takahara *et al.* 1995, Coste *et al.* 2001).

Here, we show for the first time that neonatal rat cardiomyocytes express CRH-R2 protein and secrete IL-6 in response to Ucn through the activation of CRH-R2 and subsequently MAP kinases and nuclear factor kappa B (NF- κ B).

Materials and methods

Culture of neonatal rat cardiomyocytes

This protocol was approved by the Institutional Animal Care Committee. Ventricular cardiomyocytes were isolated from the hearts of 1–3-day old neonatal male Sprague/Dawley rats (Charles River Laboratories, Inc., Wilmington, MA, USA) using the Neonatal Cardiomyocyte Isolation System (Worthington Biochemical Corporation, Lakewood, NJ, USA). Cardiomyocytes were enriched by preplating for 60 min to allow the non-myocytes to adhere to the plate. Non-adherent cardiomyocytes were then removed by aspiration and plated in DMEM/F-12 (Gibco) supplemented with 5% fetal bovine serum (Hyclone, Logan, UT, USA), 10% horse serum (Sigma), 1% penicillin/streptomycin (Gibco), 20 μ M vitamin B₁₂ (Sigma) at 37 °C in 95% air 5% CO₂. For the first 3 days of culture, 0.1 mM 5-bromo-2'-deoxyuridine (BrdU, Sigma) was added to the medium to prevent proliferation of non-myocytes. The culture medium was changed 40–48 h following seeding, to serum depleted medium: DMEM/F-12 supplemented with 1 $\times$ insulin-transferrin-selenium (Gibco), 0.5% penicillin/streptomycin, 1% fetal bovine serum and 20 μ M vitamin B₁₂; cardiomyocytes remained in this medium for 24 h before further experiments. All cells in culture were seen beating spontaneously, indicative of cardiomyocytes. Ucn is expressed in cells

of the immune system (Karalis *et al.* 1997) including mast cells (Kempuraj *et al.* 2004).

We stained the cardiomyocytes with 1% toluidine blue solution (pH <2) for mast cells and with Giemsa stain for lymphocytes in the final cardiomyocyte preparation. Electron microscopy confirmed the absence of any fibroblasts or endothelial cells (data not shown).

IL-6 measurements

Time-course and dose-response studies of Ucn-stimulated IL-6 release from cardiomyocytes were conducted. Cells were either treated with 100 nM Ucn for different periods of time (0–72 h), or treated with different concentrations of Ucn (0–10 μ M) for 48 h; following stimulation, the culture medium was collected and was assayed for IL-6 using the quantitative sandwich enzyme immunoassay technique (R&D).

Cyclic AMP assays

Cardiomyocytes were incubated in serum-free culture medium containing 100 nM Ucn for the designated times and cAMP was measured with an ELISA kit (R&D) as previously described (Heldwein *et al.* 1996).

Western blot analysis

For the detection of CRH-R2, heart tissue and isolated cardiomyocytes were disrupted in ice-cold lysis buffer (50 mM Tris pH 8.0, 150 mM NaCl, 0.1% SDS, 0.02% sodium azide, 1% NP-40, 0.5% sodium deoxycholate, 100 μ g/ml PMSF, and 1 μ g/ml aprotinin) and were then centrifuged at 15 000 *g* for 15 min at 4 °C. For the detection of phosphorylation of MAP kinases, cardiomyocytes were exposed to 100 nM Ucn in the presence or absence of inhibitors for the indicated time, after which the cells were lysed in lysis buffer, vortexed, and centrifuged at 15 000 *g* for 15 min at 4 °C. Aliquots of tissue and cell lysates were kept for protein estimation (Bio-Rad). An equal amount of protein from each sample was mixed with an equal volume of 3X SDS sample buffer. The samples were boiled for 3 min and proteins were subjected to electrophoresis on a 10% SDS-polyacrylamide gel, subsequently transferred onto the PVDF membrane (Immobilon-P) by the semi-dry method (BioRad). The membrane was then blocked with 5% dry milk in PBS containing 0.05% Tween-20 (PBS-T) and probed for 2 h at room temperature with two different rabbit polyclonal anti-CRH-R2 antibodies (Santa Cruz Biotechnology, Santa Cruz, CA, USA) and (Imgenex, San Diego, CA, USA) or an antibody that detects phospho-ERK and phospho-p38 (Cell Signaling Danvers, MA, USA). For the detection of total ERK

and p38, the membrane was probed for 1 h at room temperature with either an anti-ERK1 antibody or an anti-p38 antibody. The membrane was washed in PBS-T and incubated with a HRP-conjugated secondary antibody (Santa Cruz) for 1 h. The immunoreactive bands were visualized by enhanced chemiluminescence (ECLJ).

Electromobility gel shift assays

The nuclear extract of cardiomyocytes was collected as previously described (Li *et al.* 2001). The consensus NF- κ B oligonucleotide (AGTTGAGGGGACTTCC-CAGGC, Santa Cruz) was radiolabeled by mixing 50 ng oligonucleotide, 70 μ Ci [γ - 32 P] ATP, 1 μ l T4 polynucleotide kinase, 1.5 μ l 10 $\times$ T4 polynucleotide buffer and double-distilled water in a 15 μ l reaction volume. The nuclear extract protein (10 μ g) was then mixed with 0.5 ng [γ - 32 P] ATP-labeled NF- κ B oligonucleotide, 20 μ g BSA, 2 μ g pdI-dC, 2 μ l Buffer D+ (20 mM Hepes pH 7.9, 20% glycerol, 100 mM KCl, 0.5 mM EDTA, 0.25% NP40), 4 μ l Buffer F (20% FICOLL 400, 100 mM Hepes pH 7.9, 300 mM KCl), and DTT (2 mM) in 20 μ l total volume at room temperature for 20 min. At the end of the reaction, the mixture was loaded on a non-denaturing 4% polyacrylamide gel using a running buffer containing 50 mM Tris pH 7.5, 0.38 M glycine, and 2 mM EDTA, after which the gel was dried and exposed to Kodak X-ray film for autoradiography at -80°C .

Drug pretreatment

Cardiomyocytes were pretreated with the p38 MAP kinase inhibitor SB203580 (10 μ M) or the MEK1 inhibitor PD98059 (20 μ M) 1 h before Ucn treatment. The NF- κ B inhibitor pyrrolidine dithiocarbamate (PDTC) was added at 50 μ M to the culture medium 2 h before Ucn treatment. The peptide CRH-R antagonists α -helical (9–41) CRH or astressin-2B (10 μ M) were added 1 h before Ucn treatment. The optimal concentrations and the duration of pretreatment for all the inhibitors used were predetermined by time- and dose-response experiments (not shown). All concentrations of the different inhibitors used in these experiments have been previously shown not to be toxic to cardiomyocytes; moreover, the viability of the cardiomyocytes was intact, as they were still beating and Trypan blue exclusion was minimal at 48 h.

Statistical analysis

Each experiment was performed at least four times unless stated otherwise. The differences between unstimulated and stimulated cells were compared

using both the two-tailed Student's *t*-test and the non-parametric Mann–Whitney U-test; comparison between different treatment groups was performed by ANOVA using one-way ANOVA. Results are presented as mean $\pm$ S.E.M. For all analyses, $P < 0.05$ was considered to indicate statistical significance.

Results

Time- and dose-dependent increase of IL-6 release by urocortin

Ucn increased IL-6 release from isolated cardiomyocytes in a time- and dose-dependent fashion. Significant increase of IL-6 release was observed as early as 4 h of Ucn (100 nM) exposure. Ucn produced 15.5 ± 0.5 pg/ml of IL-6 as compared with 7.1 ± 1.4 pg/ml from unstimulated cells (Fig. 1A, $n = 6$, $P < 0.05$). The Ucn-induced IL-6 release increased with time and maximal release was observed at 48 h of stimulation when Ucn treated cells produced 120.2 ± 9.5 pg/ml as

Figure 1 Time- and dose-dependent effect of Ucn on IL-6 release from cardiomyocytes. (A) Cells were stimulated with 100 nM Ucn in serum-free medium for the designated length of time (0–72 h). (B) Cells were treated with different concentrations of Ucn (0–10 μ M) for 48 h ($n = 6$; $*P < 0.05$).

compared with 59.1 ± 7.2 pg/ml of IL-6 from untreated cells (Fig. 1A, $n=6$, $P<0.05$). Unstimulated cells also released more IL-6 over time, but the difference between them and the treatment group remained significant at every time point (Fig. 1A). Ucn-induced IL-6 release from cardiomyocytes was also dose-dependent. When cells were treated with different concentrations of Ucn for 48 h, 10 nM was the lowest concentration of Ucn that could stimulate significant IL-6 increase of 83.5 ± 5.5 pg/ml versus 59.1 ± 7.2 pg/ml in controls (Fig. 1B, $n=6$, $P<0.05$). Maximal IL-6 release of 148.3 ± 8.1 pg/ml was induced by 1 μ M of Ucn (Fig. 1B, $n=6$, $P<0.05$); higher Ucn concentrations (10 μ M) did not increase IL-6 release any further (152.0 ± 6.1 pg/ml, Fig. 1B, $n=6$).

Presence of CRH-R2 in cardiomyocytes

Western blot analysis using anti-CRH-R2 antibodies from two different sources demonstrated the presence of a strong immunoreactive band of about 49 kDa in mouse and rat heart, as well as in neonatal rat cardiomyocytes (positive controls included rat cortex and cerebellum; Fig. 2A–C). Use of a blocking peptide for CRH-R2 abolished immunoreactivity, while one for CRH-R1 had no effect (data not shown). There was no CRH-R1 expression noted (data not shown). In order to investigate whether the CRH-R2 on cardiomyocytes is functional, cellular cAMP levels were measured following Ucn stimulation, since CRH and related peptides typically bind to CRH-R and activate adenylate cyclase. Production of cAMP increased significantly by 3.4 ± 0.2 -fold above basal levels after 2 min of stimulation with 100 nM Ucn, reaching a maximum of 6.9 ± 0.2 -fold above basal level ($n=6$, $P<0.05$) at 5 min, and then decreased slowly over time, but still remained 5.3 ± 0.1 -fold above basal values at 20 min (results not shown).

Urocortin activates ERK and p38 MAP kinases

In order to study the possible mechanism(s) by which Ucn stimulates IL-6 release, we investigated the effect of Ucn on the phosphorylation of MAP kinases, including ERK and p38. Cardiomyocytes responded to 100 nM Ucn with the activation of both ERK and p38. Phosphorylation of ERK (p44/42) was observed at 2 min and was maximized at 5 min after which it decreased over time, but remained above basal level up to 1 h (Fig. 3A).

The peptide CRH-R inhibitor astressin completely reversed the effect of Ucn (100 nM for 15 min) on ERK activation (Fig. 3A). Pretreatment with PD98059 (200 nM for 1 h), the specific inhibitor of MEK1, which is the upstream stimulator of ERK, also

Figure 2 Western blot analysis of CRH-R2 in rodent hearts and isolated neonatal rat cardiomyocytes. (A) Mouse heart with rat cortex homogenates included as a positive control. (B) Mouse heart, rat heart homogenates, and rat cardiomyocytes (A & B using anti-CRH-R2 from Santa Cruz, CA, USA). (C) Rat cardiomyocytes with rat cortex and cerebellum as positive controls using a rabbit polyclonal anti-CRH-R2 antibody from Imgenex (San Diego). Representative gels ($n=3$). The amount of protein loaded for this gel was about 1/5th of gels shown in Fig. 2A and 2B due to limited availability of cardiomyocyte material.

completely inhibited the phosphorylation of ERK. The specific inhibitor of p38 MAP kinase SB203580 (100 nM) had no apparent effect on ERK activation (Fig. 3A). There was no difference in total ERK between control samples, the positive control stimulated by IL-1 β and Ucn (Fig. 3A, lower panel).

Phosphorylation of p38 MAP kinase occurred later than that of ERK (Fig. 3A and B); there was a moderate increase of p38 phosphorylation observed at 15 min of Ucn exposure (Fig. 3B). Both CRH-R antagonists, α -helical (9–41) CRH and astressin completely

Figure 3 Effect of Ucn on phosphorylation of MAP kinases by western blot analysis. (A) Phosphorylation of ERK and effects of the peptide CRH-R antagonist astressin (10 μ M), as well as the specific inhibitor of p38 MAP kinase SB203580 (10 μ M) and the specific inhibitor of MEK1 PD98059 (20 μ M) for 1 h (upper panel); total ERK was detected using an antibody to ERK1 and remained at the same level in unstimulated cells, those treated with IL-1 β (10 ng/ml for 10 min) or Ucn-treated cells. (B) Phosphorylation of p38 MAP kinase and effects of α -helical (9–41) CRH and astressin, as well as SB203580 and PD98059 (upper panel); total p38 remained at the same level (lower panel) in both unstimulated cells and Ucn-treated cells (representative gel from $n=5$; $*P<0.05$).

inhibited Ucn (100 nM for 15 min)-induced phosphorylation of p38. Since SB203580 inhibits the downstream signaling pathway of p38, it did not inhibit the phosphorylation of p38 itself (Fig. 3B); as expected, the MEK1 inhibitor PD98059 did not inhibit p38 phosphorylation either (Fig. 3B). The total p38 remained the same in Ucn-stimulated and unstimulated cells (Fig. 3B lower panel).

Urocortin activates NF- κ B

There was moderate activation of NF- κ B after Ucn treatment (100 nM, 15 min to 6 h duration), as shown by gel-shift assay (Fig. 4A). Both α -helical CRH (9–41)

and SB203580 completely inhibited Ucn (100 nM for 15 min)-induced NF- κ B activation (Fig. 4A), suggesting that the activation of NF- κ B in cardiomyocytes is mediated by CRH-R and is dependent on p38 phosphorylation. Pretreatment with PD98059 (20 μ M for 1 h) unexpectedly did not have any effect on NF- κ B activation. Supershift assay using an antibody against the p50 subunit of NF- κ B confirmed that the Ucn-stimulated increase in binding resulted from NF- κ B activity in the nuclear extracts of Ucn-treated cells (Fig. 4B).

Urocortin increases IL-6 release by a CRH-R2-induced MAP kinase- and NF- κ B-dependent mechanism

In order to determine whether Ucn-induced IL-6 release from cardiomyocytes involves MAP kinases or NF- κ B activation, cardiomyocytes were pretreated for 1 h before Ucn (100 nM) stimulation with either the p38 MAP kinase inhibitor SB203580 (10 μ M) or the MEK1 inhibitor PD98059 (20 μ M). Otherwise, cardiomyocytes were pretreated with the NF- κ B inhibitor PDTC (50 μ M) for 2 h before Ucn addition. As shown in Fig. 5A, both SB203580 and PD98059 completely inhibited Ucn-induced IL-6 release, indicating that both p38 and ERK are involved in Ucn-stimulated IL-6 release from cardiomyocytes. PDTC (50 μ M for 1 h) also completely inhibited IL-6 release stimulated by Ucn, further indicating the involvement of NF- κ B. Both α -helical CRH (9–41) and astressin (100 nM) completely inhibited Ucn-induced IL-6 release (Fig. 5B), confirming that Ucn exerts its effect on IL-6 release through the activation of CRH-R.

Discussion

Here, we show for the first time that Ucn stimulates IL-6 release from neonatal rat cardiomyocytes. CRH-R2 appears to be involved, since it is identified in cardiac tissue and in isolated cardiomyocytes by immunoblotting, and the release of IL-6 is blocked by astressin-2B. CRH-R2 was detected by antibodies from two different sources, using rat cortex and cerebellum as positive controls. The same Santa-Cruz antibody used here was previously reported to be specific (Bishop *et al.* 2006), and also identified a band around 49 kDa. Moreover, the western blot with the second anti-CRH-R2 antibody from Imgenex also generated a band around 49 kDa, as shown in the product specification. A similar molecular weight band is also recognized by one more anti-CRH-R2 antibody (data not shown) by Novus Biologicals (Littleton, CO, USA; www.novusbio.com/data_sheet/index/NB100-56485). Some authors have reported the molecular weight for the full length CRH-R2 to be about 70 kDa in the pituitary, but lower in

Figure 4 Effect of Ucn on NF-κB in cardiomyocytes. (A) NF-κB activation demonstrated by binding of NF-κB to the radiolabeled oligonucleotide shown by EMSA and effects of α -helical (9–41) CRH, SB203580 (10 μ M) and PD98059 (20 μ M). (B) Supershift assay using antibody against the p50 subunit of NF-κB confirmed that the Ucn-stimulated increase in binding resulted from NF-κB activity in the nuclear extracts of Ucn-treated cells (representative gel from $n=5$; * $P<0.05$).

the brain (Grigoriadis & De Souza 1988); the 49 kDa band may represent a CRH-R2 form that may be truncated, with less glycosylation, or a tissue specific variant.

Ucn and CRH-R2 α mRNA expression has been detected in cardiac tissue with CRH-R2 β predominantly expressed in the left ventricle (Kimura *et al.* 2002). CRH-R2 protein expression was shown by

autoradiography using a specific radiolabeled anti-CRH-R2 ligand to be very high in rodent heart, with very little CRH-R1 detected (Waser *et al.* 2006). CRH-R2 β mRNA expression was depressed in left ventricular hypertrophy, possibly through downregulation by Ucn (Nishikimi *et al.* 2000). Restraint stress upregulated Ucn mRNA, but decreased CRH-R2 β mRNA levels in the rat heart and aorta (Pournajafi *et al.* 2003). CRH-R2 mRNA expression was also downregulated by IL-1 and TNF in the mouse heart (Coste *et al.* 2001), possibly acting to limit the inotropic and chronotropic effects of Ucn.

Ucn II was shown to stimulate rat neonatal cardiomyocytes to release atrial natriuretic peptide in a CRH-R2 and cAMP-dependant manner (Ikeda *et al.* 2005). Ucn mediates stress-induced IL-6 release *in vivo* and administration of Ucn causes elevation of plasma IL-6 in rats (Ando *et al.* 1998). Ucn also stimulates IL-6 secretion from human peripheral mononuclear cells *in vitro* (Kohno *et al.* 2001), as well as an increase in IL-6 mRNA levels through CRH-R2 in rat aortic smooth muscle cells (Kageyama *et al.* 2007).

Our study demonstrates that IL-6 release by Ucn involves p38 MAP kinase, ERK and NF-κB activation. Our results are consistent with a previous study that showed p38 MAP kinase activation by Ucn, but not CRH, in cultured human pregnant myometrial cells (Grammatopoulos *et al.* 2000), as well as NF-κB involvement in TNF-induced IL-6 expression and release from cardiomyocytes (Craig *et al.* 2000) and upregulation of inflammatory IL-6 expression (TNF- α , IL-1 and IL-6) in cardiomyocyte hypertrophy (Purcell *et al.* 2001). Surprisingly, our results show that while Ucn-induced IL-6 release requires NF-κB activation, Ucn-induced NF-κB activation does not appear to require ERK. This discrepancy may indicate that ERK1/2 contributes to Ucn stimulation of IL-6 production independent of NF-κB.

Figure 5 Ucn increases IL-6 release in cardiomyocytes through CRHR MAP kinase and NF-κB. (A) Pretreatment with SB203580 at 10 μ M, PD98059 at 20 μ M for 1 h before Ucn stimulation (100 nM, 48 h) or the NF-κB inhibitor pyrrolidine dithiocarbamate (PDTC) at 50 μ M 2 h before Ucn stimulation (100 nM, 48 h) ($n=6$; * $P<0.05$ versus control; # $P<0.05$ versus Ucn 100 nM). (B) Pretreatment with α -helical (9–41) CRH or astressin-2B at 10 μ M for 1 h before Ucn stimulation (100 nM, 48 h) ($n=5$; * $P<0.05$ versus control; # $P<0.05$ versus Ucn 100 nM).

The role of Ucn in the heart is far from clear. Ucn protects the rat heart from IR injury *in vitro* and *in vivo* through upregulation of the p42/p44 MAPK pathway (Schulman *et al.* 2003). Ucn-II and Ucn-III are also cardioprotective against IR injury (Brar *et al.* 2004, Woodcock 2004). Stimulation of CRH-R2 β by Ucn-II and III resulted in cAMP-independent ERK1/2 phosphorylation during IR and reduced infarct size (Brar *et al.* 2004). Both Ucn (Brar *et al.* 2000) and IL-6 (Craig *et al.* 2000) may protect neonatal rat cardiomyocytes from apoptosis induced by hypoxia. However, while i.v. administration of Ucn-II had inotropic and chronotropic effects in mice and humans, it decreased blood pressure, effects absent in CRH-R2 knockout mice (Bale *et al.* 2004, Davis *et al.* 2007). Ucn-II also decreased mean arterial blood pressure in the rat, which was blocked by the CRH-R2 antagonist anti-sauvagine without any effect on normal cardiac function (Mackay *et al.* 2003). Ucn is released before and independently of cardiomyocyte death (Knight *et al.* 2008) and could have proinflammatory actions (Chrousos 1995) through a mechanism that involves mast cells (Theoharides *et al.* 2004) as in allergic asthma (Wu *et al.* 2006), and in intestinal inflammation (Kokkotou *et al.* 2006, Wu *et al.* 2006) in rats.

Cardiomyocyte-derived IL-6 in response to Ucn may even have dual and opposing actions. It might originally lead to inhibition of apoptosis, but eventually to hypertrophy alone or following angiotensin II (Sano *et al.* 2000). In macrophages, both CRH-R1 and CRH-R2 agonists have an early anti-inflammatory, but delayed pro-inflammatory effect on TNF- α release (Tsatsanis *et al.* 2007). The particular effects of Ucn may depend on the stage of maturation of the cardiomyocytes and/or activation of specific CRH-R2 isoforms, documented in keratinocytes (Pisarchik & Slominski 2004) and mast cells (Cao *et al.* 2005). For instance, a soluble CRH-R2 α isoform was shown to neutralize the effect of CRH-R agonists (Chen *et al.* 2005), while Ucn mRNA antisense transcripts have been identified in rat tissue (Shi *et al.* 2000).

Acute stress is implicated in cardiovascular pathology (Theoharides & Cochrane 2004), especially in eliciting MI in patients with CAD (Jiang *et al.* 1996). Acute stress elevates plasma IL-6 levels in rodents (Ando *et al.* 1998, Nukina *et al.* 2001) and such levels are higher in ApoE knockout mice that develop atherosclerosis (Huang *et al.* 2003). IL-6 plays a crucial role in MI (Miyao *et al.* 1993), CAD (Huber *et al.* 1999), cardiac hypertrophy (Sano *et al.* 2000), and congestive heart failure (Deliargyris *et al.* 2000). IL-6 expression is also increased in coronary arteries of aged rats that are more prone to atherosclerosis and CAD (Cesari *et al.* 2003). The importance of inflammation including IL-6 and its stimulation of CRP production in atherosclerosis was reviewed recently (Packard & Libby 2008).

Elevated intracoronary levels of IL-6 (Deliargyris *et al.* 2000) shown in patients with acute CAD may also be involved in coronary hypersensitivity leading to Kounis syndrome (Kounis *et al.* 2007). Consequently, any suggestion that Ucn peptides could be used therapeutically for heart conditions may be too premature.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported in part by Theta Biomedical Consulting and Development Co. Inc., (Brookline, MA) to TCT.

Acknowledgements

We thank Mrs Jessica Christian for her word processing skills.

References

- Ando T, Rivier J, Yanaihara H & Arimura A 1998 Peripheral corticotropin-releasing factor mediates the elevation of plasma IL-6 by immobilization stress in rats. *American Journal of Physiology* **275** R1461–R1467.
- Bale TL, Giordano FJ & Vale WW 2003 A new role for corticotropin-releasing factor receptor-2: suppression of vascularization. *Trends in Cardiovascular Medicine* **13** 68–71.
- Bale TL, Hoshijima M, Gu Y, Dalton N, Anderson KR, Lee KF, Rivier J, Chien KR, Vale WW & Peterson KL 2004 The cardiovascular physiologic actions of urocortin II: acute effects in murine heart failure. *PNAS* **101** 3697–3702.
- Bishop GA, Tian JB, Stanke JJ, Fischer AJ & King JS 2006 Evidence for the presence of the type 2 corticotropin releasing factor receptor in the rodent cerebellum. *Journal of Neuroscience Research* **84** 1255–1269.
- Brar BK, Jonassen AK, Stephanou A, Santilli G, Railson J, Knight RA, Yellon DM & Latchman DS 2000 Urocortin protects against ischemic and reperfusion injury via a MAPK-dependent pathway. *Journal of Biological Chemistry* **275** 8508–8514.
- Brar BK, Jonassen AK, Egorina EM, Chen A, Negro A, Perrin MH, Mjos OD, Latchman DS, Lee KF & Vale W 2004 Urocortin-II and urocortin-III are cardioprotective against ischemia reperfusion injury: an essential endogenous cardioprotective role for corticotropin releasing factor receptor type 2 in the murine heart. *Endocrinology* **145** 24–35.
- Cao J, Papadopoulou N, Kempuraj D, Boucher WS, Sugimoto K, Cetrulo CL & Theoharides TC 2005 Human mast cells express corticotropin-releasing hormone (CRH) receptors and CRH leads to selective secretion of vascular endothelial growth factor. *Journal of Immunology* **174** 7665–7675.
- Cesari M, Penninx BW, Newman AB, Kritchevsky SB, Nicklas BJ, Sutton-Tyrrell K, Tracy RP, Rubin SM, Harris TB & Pahor M 2003 Inflammatory markers and cardiovascular disease (the health, aging and body composition (health ABC) study). *American Journal of Cardiology* **92** 522–528.
- Chen AM, Perrin MH, Digruccio MR, Vaughan JM, Brar BK, Arias CM, Lewis KA, Rivier JE, Sawchenko PE & Vale WW 2005 A soluble mouse brain splice variant of type 2 alpha corticotropin-releasing factor (CRF) receptor binds ligands and modulates their activity. *PNAS* **102** 2620–2625.

- Chrousos GP 1995 The hypothalamic-pituitary-adrenal axis and immune-mediated inflammation. *New England Journal of Medicine* **332** 1351–1362.
- Coste SC, Heldwein KA, Stevens SL, Tobar-Dupres E & Stenzel-Poore M 2001 IL-1 alpha and TNF alpha down-regulate CRH receptor-2 mRNA expression in the mouse heart. *Endocrinology* **142** 3537–3545.
- Craig R, Larkin A, Mingo AM, Thuerauf DJ, Andrews C, McDonough PM & Glembotski CC 2000 p38 MAPK and NF-kappa B collaborate to induce interleukin-6 gene expression and release, evidence for a cytoprotective autocrine signaling pathway in a cardiac myocyte model system. *Journal of Biological Chemistry* **275** 23814–23824.
- Davis ME, Pemberton CJ, Yandle TG, Fisher SF, Lainchbury JG, Frampton CM, Rademaker MT & Richards AM 2007 Urocortin 2 infusion in healthy humans: hemodynamic, neurohormonal, and renal responses. *Journal of the American College of Cardiology* **49** 461–471.
- Deliaargyris EN, Raymond RJ, Theoharides TC, Boucher WS, Tate DA & Dehmer GJ 2000 Sites of interleukin-6 release in patients with acute coronary syndromes and in patients with congestive heart failure. *American Journal of Cardiology* **86** 913–918.
- Fisman EZ, Benderly M, Esper RJ, Behar S, Boyko V, Adler Y, Tanne D, Matas Z & Tenenbaum A 2006 Interleukin-6 and the risk of future cardiovascular events in patients with angina pectoris and/or healed myocardial infarction. *American Journal of Cardiology* **98** 14–18.
- Grammatopoulos DK, Randeava HS, Levine MA, Katsanou ES & Hillhouse EW 2000 Urocortin, but not corticotropin-releasing hormone (CRH), activates the mitogen-activated protein kinase signal transduction pathway in human pregnant myometrium: an effect mediated via R1alpha and R2beta CRH receptor subtypes and stimulation of Gq-proteins. *Molecular Endocrinology* **14** 2076–2091.
- Grigoriadis DE & De Souza EB 1988 The brain corticotropin-releasing factor (CRF) receptor is of lower apparent molecular weight than the CRF receptor in anterior pituitary, evidence from chemical cross-linking studies. *Journal of Biological Chemistry* **263** 10927–10931.
- Heldwein KA, Redick DL, Rittenberg MB, Claycomb WC & Stenzel-Poore MP 1996 Corticotropin-releasing hormone receptor expression and functional coupling in neonatal cardiac myocytes and AT-1 cells. *Endocrinology* **137** 3631–3639.
- Huang M, Pang X, Karalis K & Theoharides TC 2003 Stress-induced interleukin-6 release in mice is mast cell-dependent and more pronounced in apolipoprotein E knockout mice. *Cardiovascular Research* **59** 241–249.
- Huber SA, Sakkinen P, Conze D, Hardin N & Tracy R 1999 Interleukin-6 exacerbates early atherosclerosis in mice. *Arteriosclerosis and Thrombosis* **19** 2364–2367.
- Ikeda K, Tojo K, Tokudome G, Ohta M, Sugimoto K, Tamura T, Tajima N, Mochizuki S, Kawakami M & Hosoya T 2003 Cardiac expression of urocortin (Ucn) in diseased heart; preliminary results on possible involvement of Ucn in pathophysiology of cardiac diseases. *Molecular and Cellular Biochemistry* **252** 25–32.
- Ikeda K, Tojo K, Otsubo C, Udagawa T, Hosoya T, Tajima N, Nakao K & Kawamura M 2005 Effects of urocortin II on neonatal rat cardiac myocytes and non-myocytes. *Peptides* **26** 2473–2481.
- Jiang W, Babyak M, Krantz DS, Waugh RA, Coleman E, Hanson MM, Frid DJ, McNulty S, Morris JJ, O'Connor CM *et al.* 1996 Mental stress-induced myocardial ischemia and cardiac events. *JAMA* **275** 1651–1656.
- Kageyama K, Hanada K, Nigawara T, Furukawa K, Terui K, Ogura E, Motomura S & Suda T 2007 Inhibitory effects of glucocorticoids on urocortin-mediated increases in interleukin-6 gene expression in rat aortic smooth muscle cells. *Peptides* **28** 1059–1067.
- Karalis K, Louis JM, Bae D, Hilderbrand H & Majzoub JA 1997 CRH and the immune system. *Journal of Neuroimmunology* **72** 131–136.
- Kempuraj D, Papadopoulou NG, Lytinas M, Huang M, Kandere-Grzybowska K, Madhappan B, Boucher W, Christodoulou S, Athanassiou A & Theoharides TC 2004 Corticotropin-releasing hormone and its structurally related urocortin are synthesized and secreted by human mast cells. *Endocrinology* **145** 43–48.
- Kimura Y, Takahashi K, Totsune K, Muramatsu Y, Kaneko C, Darnel AD, Suzuki T, Ebina M, Nukiwa T & Sasano H 2002 Expression of urocortin and corticotropin-releasing factor receptor subtypes in the human heart. *Journal of Clinical Endocrinology and Metabolism* **87** 340–346.
- Knight RA, Chen-Scarabelli C, Yuan Z, McCauley RB, Di RJ, Scarabelli GM, Townsend PA, Latchman D, Saravolatz L, Faggian G *et al.* 2008 Cardiac release of urocortin precedes the occurrence of irreversible myocardial damage in the rat heart exposed to ischemia/reperfusion injury. *FEBS Letter* **582** 984–990.
- Kohno M, Kawahito Y, Tsubouchi Y, Hashiramoto A, Yamada R, Inoue KI, Kusaka Y, Kubo T, Elenkov IJ, Chrousos GP *et al.* 2001 Urocortin expression in synovium of patients with rheumatoid arthritis and osteoarthritis: relation to inflammatory activity. *Journal of Clinical Endocrinology and Metabolism* **86** 4344–4352.
- Kokkotou E, Torres D, Moss AC, O'Brien M, Grigoriadis DE, Karalis K & Pothoulakis C 2006 Corticotropin-releasing hormone receptor 2-deficient mice have reduced intestinal inflammatory responses. *Journal of Immunology* **177** 3355–3361.
- Kounis NG, Hahalis G & Theoharides TC 2007 Coronary stents, hypersensitivity reactions and the Kounis syndrome. *Journal of Invasive Cardiology* **20** 314–323.
- Krantz DS, Sheps DS, Carney RM & Natelson BH 2000 Effects of mental stress in patients with coronary artery disease: evidence and clinical implications. *Journal of the American Medical Association* **283** 1800–1802.
- Latchman DS 2003 Urocortin protects against ischemic injury via a MAPK-dependent pathway. *Trends in Cardiovascular Medicine* **11** 169.
- Li RCX, Ping P, Zhang J, Wead WB, Cao X, Gao J, Zheng Y, Huang S, Han J & Bolli R 2001 PKCε modulates NF-κB and AP-1 via mitogen-activated protein kinases in adult rabbit cardiomyocytes. *American Journal of Physiology. Heart and Circulatory Physiology* **279** 1679–1689.
- Luc G, Bard JM, Juhan-Vague I, Ferrieres J, Evans A, Amouyel P, Arveiler D, Fruchart JC & Ducimetiere P 2003 C-reactive protein, interleukin-6, and fibrinogen as predictors of coronary heart disease: the PRIME Study. *Arteriosclerosis, Thrombosis and Vascular Biology* **23** 1255–1261.
- Mackay KB, Stiefel TH, Ling N & Foster AC 2003 Effects of a selective agonist and antagonist of CRF2 receptors on cardiovascular function in the rat. *European Journal of Pharmacology* **469** 111–115.
- Miyao Y, Yasue H, Ogawa H, Misumi I, Masuda T, Sakamoto T & Morita E 1993 Elevated plasma interleukin-6 levels in patients with acute myocardial infarction. *American Heart Journal* **126** 1304.
- Nishikimi T, Miyata A, Horio T, Yoshihara F, Nagaya N, Takishita S, Yutani C, Matsuo H, Matsuoaka H & Kangawa K 2000 Urocortin, a member of the corticotropin-releasing factor family, in normal and diseased heart. *American Journal of Physiology. Heart and Circulatory Physiology* **279** H3031–H3039.
- Nukina H, Sudo N, Aiba Y, Oyama N, Koga Y & Kubo C 2001 Restraint stress elevates the plasma interleukin-6 levels in germ-free mice. *Journal of Neuroimmunology* **115** 46–52.
- Okosi A, Brar BK, Chan M, D'Souza L, Smith E, Stephanou A, Latchman DS, Chowdrey HS & Knight RA 1998 Expression and protective effects of urocortin in cardiac myocytes. *Neuropeptides* **32** 167–171.
- Packard RR & Libby P 2008 Inflammation in atherosclerosis: from vascular biology to biomarker discovery and risk prediction. *Clinical Chemistry* **54** 24–38.
- Pisarchik A & Slominski A 2004 Molecular and functional characterization of novel CRFR1 isoforms from the skin. *European Journal of Biochemistry* **271** 2821–2830.
- Pournajafi NH, Tanaka Y, Dorobantu M & Hashimoto K 2003 Modulation of corticotropin-releasing hormone receptor type 2 mRNA expression by CRH deficiency or stress in the mouse heart. *Regulatory Peptides* **115** 131–138.

- Purcell NH, Tang G, Yu C, Mercurio F, DiDonato JA & Lin A 2001 Activation of NF-kappa B is required for hypertrophic growth of primary rat neonatal ventricular cardiomyocytes. *PNAS* **98** 6668–6673.
- Sano M, Fukuda K, Kodama H, Pan J, Saito M, Matsuzaki J, Takahashi T, Makino S, Kato T & Ogawa S 2000 Interleukin-6 family of cytokines mediate angiotensin II-induced cardiac hypertrophy in rodent cardiomyocytes. *Journal of Biological Chemistry* **275** 29717–29723.
- Schulman D, Latchman DS & Yellon DM 2003 Urocortin protects the heart from reperfusion injury via upregulation of p42/p44 MAPK signaling pathway. *American Journal of Physiology. Heart and Circulatory Physiology* **283** H1481–H1488.
- Shi M, Yan X, Ryan DH & Harris RB 2000 Identification of urocortin mRNA antisense transcripts in rat tissue. *Brain Research Bulletin* **53** 317–324.
- Theoharides TC & Cochrane DE 2004 Critical role of mast cells in inflammatory diseases and the effect of acute stress. *Journal of Neuroimmunology* **146** 1–12.
- Theoharides TC, Donelan JM, Papadopoulou N, Cao J, Kempuraj D & Conti P 2004 Mast cells as targets of corticotropin-releasing factor and related peptides. *Trends in Pharmacological Sciences* **25** 563–568.
- Townsend PA, Davidson SM, Clarke SJ, Khaliulin I, Carroll CJ, Scarabelli TM, Knight RA, Stephanou A, Latchman DS & Halestrap AP 2007 Urocortin prevents mitochondrial permeability transition in response to reperfusion injury indirectly by reducing oxidative stress. *American Journal of Physiology. Heart and Circulatory Physiology* **293** 928–938.
- Tsatsanis C, Androulidaki A, Dermizaki E, Gravanis A & Margioris AN 2007 Corticotropin releasing factor receptor 1 (CRF1) and CRF2 agonists exert an anti-inflammatory effect during the early phase of inflammation suppressing LPS-induced TNF-alpha release from macrophages via induction of COX-2 and PGE2. *Journal of Cell Physiology* **210** 774–783.
- Waser B, Rehmann R, Rivier J, Vale W & Reubi JC 2006 CRF receptors in the rodent and human cardiovascular systems: species differences. *Peptides* **27** 3029–3038.
- Woodcock EA 2004 Ucn-II and Ucn-III are cardioprotective against ischemia reperfusion injury: an essential endogenous cardioprotective role for CRFR2 in the murine heart. *Endocrinology* **145** 21–23.
- Wu Y, Zhou H, Xu Y & Li S 2006 Enhanced expression of urocortin in lung tissues of rats with allergic asthma. *Biochemical and Biophysical Research Communications* **341** 532–540.
- Yamauchi-Takahara K, Ihara Y, Ogata A, Yoshizaki K, Azuma J & Kishimoto T 1995 Hypoxic stress induces cardiac myocyte-derived interleukin-6. *Circulation* **91** 1520–1524.

Received in final form 18 January 2009

Accepted 11 February 2009

Made available online as an Accepted Preprint 11 February 2009