Aromatase promoter I.f is regulated by progesterone receptor in mouse hypothalamic neuronal cell lines

M Bertan Yilmaz^{1,2}, Andrew Wolfe³, Hong Zhao¹, David C Brooks¹ and Serdar E Bulun¹

¹Division of Reproductive Biology Research, Department of Obstetrics and Gynecology, Northwestern University Feinberg School of Medicine, 303 E. Superior Street, Suite 4-123, Chicago, Illinois 60611, USA

²Division of Medical Sciences, Department of Medical Biology and Genetics, School of Medicine, Cukurova University, Adana 01330, Turkey

³Division of Endocrinology, Department of Pediatrics, Johns Hopkins University, Baltimore, Maryland 21287, USA

(Correspondence should be addressed to S E Bulun; Email: s-bulun@northwestern.edu)

Abstract

Aromatase catalyzes the conversion of C₁₉ steroids to estrogens. Aromatase and progesterone, both of which function at different steps of steroidogenesis, are crucial for the sexually dimorphic development of the fetal brain and the regulation of gonadotropin secretion and sexual interest in adults. The aromatase gene (*Cyp19a1*) is selectively expressed in distinct neurons of the mouse hypothalamus through a distal brain-specific promoter, I.f, located ~40 kb upstream of the coding region. However, the regulation of aromatase expression in the brain is not well understood. In this study, we investigated a short feedback effect of progesterone analogues on aromatase mRNA expression and enzyme activity in estrogen receptor α (Esr1)-positive or -negative mouse embryonic hypothalamic neuronal cell lines that express aromatase via promoter I.f. In a hypothalamic neuronal cell line that highly expresses aromatase, progesterone receptor (Pgr), and Esr1, a progesterone agonist, R5020, inhibited aromatase mRNA level and enzyme activity. The inhibitory effect of R5020 was reversed by its antagonist, RU486. Deletion mutants of promoter I.f suggested that inhibition of aromatase expression by progesterone is conferred by the nt -1000/-500 region, and R5020 enhanced binding of Pgr to the nt -800/-600 region of promoter I.f. Small interfering RNA knockdown of *Pgr* eliminated progesterone-dependent inhibition of aromatase mRNA and enzyme activity. Taken together, progesterone enhances recruitment of Pgr to specific regions of the promoter I.f of *Cyp19a1* and regulates aromatase expression in hypothalamic neurons.

Journal of Molecular Endocrinology (2011) **47**, 69–80

Introduction

Sex steroids influence most aspects of cellular organization and mammalian development. They play important roles in shaping neural functions and reproductive behavior throughout all stages of life. One of the major sex steroids is progesterone, which has a wide spectrum of functions in many organs, including the brain. Progesterone is primarily synthesized by the corpus luteum in females and by the testes and adrenal cortex in males, but it can also be locally synthesized by other tissues such as the brain (Keefe 2002, Wagner 2006).

Besides its well-established organizational and activation effects on reproductive neuroendocrine function and recovery after brain injury, progesterone also exerts a wide variety of actions on regions of the developing and adult brain that influence higher cognitive functions, pain mechanisms, fine motor skills, susceptibility to seizures, mood, temperature regulation, and sleep (Levine *et al.* 2001, Pluchino *et al.* 2006, Wagner 2006). Estrogen biosynthesis is dependent on the entry of cholesterol into mitochondria, followed by six enzymatic steps. Aromatase is the key enzyme that

catalyzes the final step, the conversion of C₁₉ steroids to biologically active estrogen, estradiol (E₂; Simpson & Davis 2001, Bulun *et al.* 2005).

A single-copy gene (*Cyp19a1*) encodes aromatase, the inhibition of which effectively eliminates estrogen production in the entire body (Simpson & Davis 2001). The mouse *Cyp19a1* gene, which spans a 30 kb coding region and a 75 kb regulatory region (~105 kb in total length), is located on the long arm of chromosome 9 (Zhao *et al.* 2009). The downstream 30 kb coding region comprises nine coding exons (II–X; Golovine *et al.* 2003). The upstream 75 kb portion of the gene contains multiple promoters that direct transcription of alternative first exons, giving rise to aromatase mRNA species with unique 5'-untranslated regions (Chow *et al.* 2009). The upstream regulatory region contains two major promoters regulated in a tissue-specific manner: the proximal gonad-specific promoter and the distal brain-specific promoter, which lies ~40 kb upstream of the ATG translational start site in coding exon II (Honda *et al.* 1996; <http://www.ncbi.nlm.nih.gov/BLAST/>).

In the vertebrate brain, aromatase is primarily expressed in the hypothalamus, hippocampus, and

amygdala via a highly conserved promoter, I.f (Lephart 1997, Honda *et al.* 1999). Aromatase expression in the hypothalamus is primarily localized in the medial preoptic area and the ventromedial nucleus of the hypothalamus, which are the centers that govern reproductive functions of both sexes of different species (Lephart 1996, Sharma *et al.* 2004, Peterson *et al.* 2005, Voigt *et al.* 2007, Zhao *et al.* 2007).

The physiological roles of aromatase in the brain became obvious after examining mice with a disrupted aromatase gene (aromatase knockout (ArKO)) and men with inactivating mutations of the aromatase gene (Fisher *et al.* 1998, Jones *et al.* 2006). In both cases, aromatase deficiency was associated with increased testosterone and gonadotropin levels, indicating that aromatase and E₂ were essential for regulating gonadotropin secretion. In addition, libido was significantly decreased in aromatase-deficient male mice and men and could be restored by E₂ treatment, thus revealing essential roles for aromatase and E₂ in regulating sexual behavior (Fisher *et al.* 1998, Bakker *et al.* 2004a,b, Simpson 2004b).

The signaling pathways or molecular mechanisms that regulate the brain-specific aromatase promoter I.f are not well understood. Several groups of investigators have found that protein kinases A and C and cAMP regulate aromatase expression and activity in the brain (Balthazart *et al.* 2001, Lavaque *et al.* 2006). Other groups showed that testosterone also upregulates aromatase mRNA and enzyme activity in the brain (Abdelgadir *et al.* 1994, Roselli *et al.* 1997, Negri-Cesi *et al.* 2001). Moreover, E₂ was found to upregulate and downregulate hypothalamic aromatase mRNA and enzyme activity under both *in vivo* and *in vitro* circumstances (Iivonen *et al.* 2006, Zhao *et al.* 2007). However, progesterone has been thought to suppress aromatase mRNA levels and enzyme activity in some tissues (Fortune & Vincent 1983) but induce it in others, implying that it may have divergent effects on aromatase (Lephart *et al.* 1998, Trainor *et al.* 2003, Pluchino *et al.* 2006, Wagner 2006). However, none of these studies provided a connection between aromatase expression and the regulation of promoter I.f in brain tissue. One group proposed that the transcription factor stress response element regulates the aromatase promoter via binding to its upstream region (Honda *et al.* 1999). Progesterone, a steroid hormone primarily secreted by the ovary and also locally synthesized in the brain, exerts its functions mostly via progesterone receptor (Pgr). Previously, progesterone has been proposed to regulate aromatase expression in the brain (Rhoda *et al.* 1987, Ing & Tornesi 1997, Prange-Kiel *et al.* 2001). In this regard, we conducted this study to elucidate the mechanisms regulating hypothalamic aromatase activity by progesterone via the brain-specific aromatase promoter I.f in mouse hypothalamic neuronal cell lines.

Materials and methods

Cell culture

Mouse embryonic hypothalamic neuronal cell lines were purchased from Cellutions Biosystems, Inc. (Toronto, Canada) and were cultured in DMEM (Gibco) supplemented with 10% fetal bovine serum (FBS; Gibco) and 1% penicillin–streptomycin (Gibco) at 37 °C in 95% air and 5% CO₂. The cells were studied at passages 15–16. After 24 h of serum starvation, cells were treated with various doses of the progesterone agonist, R5020 (Promegestone, Sigma–Aldrich), for varying lengths of time.

RNA isolation and real-time RT-PCR

Hypothalamic neurons were treated with either vehicle (ethanol) or 10^{−7}, 10^{−8}, or 10^{−9} M R5020, and total RNA was extracted after 6, 12, and 24 h of treatment using TRI reagent (Sigma) as per the manufacturer's instructions. On-column DNase digestion was carried out using a DNase I kit (Qiagen). The integrity of the isolated RNA was verified by running 5 µg total RNA on a 1% formaldehyde gel. Total RNA (5 µg) from vehicle, R5020 or progesterone antagonist, RU486 (mifepristone)-treated hypothalamic neurons was used for reverse transcription in a final volume of 20 µl using the Superscript III First Strand RT synthesis kit (Invitrogen) according to the manufacturer's protocol. cDNA (5 µl) was used for real-time PCR (the Taqman real-time primers and probe for aromatase were designed using ABI primer express Software 3.0) on an Applied Biosystems 7000 Sequence Detection System (Applied Biosystems, Foster City, CA, USA). The reactions were carried out using the ABI Taqman assay system for aromatase promoter I.f and power SYBR green for the aromatase coding region according to the manufacturer's protocol. The primers and probes used are as follows: Taqman assay primers for aromatase promoter I.f; forward 5'-AACTCACCATCTTCAAGAGTCCA-3', reverse 5'-GAGTGGCATGGCACTGACAGT-3'; probe 5'-AGGTCCGGTTTA-3'. Each contained a 6-carboxy-fluorescein phosphoramidite (FAM dye) label at the 5'-end and a minor groove binder and nonfluorescent quencher at the 3'-end and were designed to hybridize to the junction between promoter I.f and exon II. SYBR green assay primers for aromatase are as follows: forward 5'-TGTGTTGACCCATCATGAGACA-3', reverse 5'-CTTGACGGATCGTTCACTTTC-3'; for estrogen receptor α (*Esr1*): forward 5'-ATGAAAGGCGGCATACGGAAG-3', reverse 5'-CACCCATTTCATTTCCGGCCTTC-3'; and for glyceraldehyde-3-phosphate dehydrogenase (*Gapdh*): forward 5'-ACCACAGTCCATGCCATCAC-3', reverse 5'-TCCACACCCTGTTGCTGTA-3'.

In order to elucidate which *Pgr* isoform is predominantly expressed in hypothalamic neuronal cell lines, we amplified all three isoforms employing real-time PCR. For *Pgr-B*: forward 5'-GTGGAGGGCGCTT-TCTCTG-3', reverse 5'-TCTGCCTCCCTCCCTATGAGT-3'; *Pgr-AB*: forward 5'-CTGGAGACCGAGGGCTCT-3', reverse 5'-CCAGTGCTCGAGGTTTGCTG-3'; *Pgr-ABC*: forward 5'-GACACTGGCTGTGGAATTTCC-3', reverse 5'-CCAGGATCTTGGGCAACTG-3'. Conventional PCR primers for *Esr1*: forward 5'-TCCTTCTAGACCCTTCAGTGAAGCC-3', reverse 5'-ACATGTCAAAGATCTC-CACCATGCC-3'.

PCR was carried out in triplicate in a 50 µl reaction volume using SYBR green PCR mix. The reactions were incubated at 50 °C for 2 min and 95 °C for 10 min, followed by 40 cycles of 95 °C for 15 s and 60 °C for 1 min. The threshold cycle (C_T) was defined as the fractional cycle number at which the fluorescence passes the fixed threshold. TaqMan C_T values were converted into absolute copy numbers. All RNA samples were normalized to the expression levels of *Gapdh* endogenous control. 'No template' and 'no RT' controls were used to ensure the absence of genomic DNA and reaction specificity. Totally, three independent experiments were performed to demonstrate reproducibility. Real-time RT-PCR product specificity was confirmed by melt curve analysis, gel electrophoresis, and product sequencing.

Aromatase enzyme activity assay

Aromatase enzyme activity within hypothalamic neurons was measured by a [^3H] water release assay, which is used routinely in our laboratory (Shozu *et al.* 2003). In each well, 60 pmol [^3H]androstenedione (PerkinElmer Life Sciences, Waltham, MA, USA) and 240 pmol cold androstenedione (Sigma) were added to 2 ml serum-free DMEM covering hypothalamic neurons in culture dishes. In order to study the effect of R5020, hypothalamic neurons at 80–90% confluency were serum starved for 24 h and then treated with vehicle or R5020 for 6, 12, and 24 h at 37 °C in 95% air and 5% CO_2 . Each treatment was performed in triplicate. A mixture of labeled and cold androstenedione was added to each well 4 h before the end of each time point and incubated until the completion of each treatment period. [^3H]Androstenedione conversion to [^3H]E $_2$ was stopped by adding 10% (weight/volume) trichloroacetic acid. Steroidal compounds containing unconverted [^3H]androstenedione were removed from the mixture by first mixing with 4 ml chloroform followed by centrifugation at 900 g. The upper aqueous layer was removed and mixed with dextran-coated charcoal (1% weight/volume). Charcoal was precipitated by centrifugation. From each tube, 2 ml clear solution was placed into a scintillation vial containing

10 ml scintillation fluid and counted in a scintillation counter (LS 6500, Beckman Coulter, Inc., Fullerton, CA, USA).

Immunoblotting

Hypothalamic neurons were cultured in 150 mm dishes in DMEM media containing 10% FBS until reaching 80% confluency. Total protein was extracted from neurons using the M-PER mammalian protein extraction reagent (Pierce, Rockford, IL, USA). Protein concentration was determined using the BCA protein assay kit (Pierce). Cell lysate (50 µg) was mixed with 2×Laemmli Sample Buffer (Bio-Rad Laboratories) and fractionated in 4–15% SDS-PAGE. Proteins from the gel were transferred to a nitrocellulose membrane. The membranes were then incubated with anti-Esr1 or anti-Pgr antibody (1:1000 dilution; Upstate, Charlottesville, VA, USA) in 5% milk solution overnight at 4 °C. The membrane was washed for 30 min and then incubated with anti-rabbit IgG-peroxidase conjugate (Upstate) as a secondary antibody (1:3000 dilution). Incubation with the secondary antibody was performed at room temperature for 1 h. The signal was detected using SuperSignal West Femto maximum sensitivity substrate chemiluminescence kit (Pierce) according to the manufacturer's instructions followed by exposure to BioMax ML X-ray film (Eastman Kodak) for 1–5 min. Then, the membranes were stripped of the first antibody and probed with anti-actin followed by the same signal detection procedure.

Transient transfection and luciferase assays

Transient transfection of N42 hypothalamic neurons was carried out in 12-well plates with Fugene HD transfection reagent (Roche Diagnostics) with the following plasmids: 1) 1 µg pGL3-basic luciferase reporter plasmid containing the promoter I.f nt –1000/–1 fragment or progressively truncated promoter I.f fragments (–700, –500, –200, and –50) and 2) 50 ng pRL-TK plasmid as an internal control (Promega). Neurons were serum deprived 1 day before transfection. After 24 h of transfection, N42 hypothalamic neurons were treated either with vehicle (ethanol) or with 10^{-7} M R5020 for 24 h. At the end of the treatment period, transfected neurons were washed twice in PBS and lysed in 250 µl lysis buffer (0.1 M potassium phosphate (pH 7.8), 1% Triton X-100, 1 mM dithiothreitol, and 2 mM EDTA). Luciferase assays were performed with 20 µl cell lysate employing a Dual-Luciferase Reporter Assay System kit (Promega). Luminescence was measured with a LUMAT LB9507 luminometer (EG&G Berthold, Bad Wildbad, Germany). Results are presented as the average of

luciferase activity from triplicate experiments and expressed as the ratio to the internal standard Renilla luciferase. Plasmids used in transfection experiments were purified using an EndoFree Plasmid Isolation Kit (Qiagen), and their purity was verified by spectrophotometry and agarose gel electrophoresis. All transfection assays were performed using equimolar amounts of plasmids.

Chromatin immunoprecipitation assay

Chromatin immunoprecipitation (ChIP) assays were performed with the Acetyl-Histone H4 Immunoprecipitation Assay Kit (Upstate) following the manufacturer's instructions with minor modifications. Briefly, after reaching 80–90% confluency in 150 mm culture dishes ($\sim 2 \times 10^7$ cells), hypothalamic neurons were serum starved for 24 h, then treated with either vehicle (ethanol) or 10^{-7} M R5020 for 6 h. Neurons were then cross-linked with 1% formaldehyde for 10 min at room temperature on an orbital shaker followed by the addition of 1 M glycine for 5 min. Neurons were washed with cold PBS containing protease inhibitors and were lysed using a lysis buffer (50 mM Tris-HCl, pH 8.1, containing 1% SDS and 10 mM EDTA). The cells were sheared using a Sonic Dismembrator Model 100 (Fisher Scientific, Hampton, NH, USA) at 70% maximum amplitude (15 s on, 60 s off) for five pulses to obtain DNA fragments in the range of 200–1000 bp. The sonicated cell supernatant was diluted tenfold in ChIP dilution buffer and 40 μ l was used as input DNA (positive control). Pre-cleared chromatin was used for immunoprecipitation reactions with 5 μ g of the Pgr antibody or nonspecific (NS) rabbit IgG (Upstate). The reactions were incubated on a rotator overnight at 4 °C. The immune complexes were collected using protein A-agarose/DNA beads and washed five times using low salt, high salt, and lithium chloride and twice with Tris-EDTA buffers. The absorbed immune complexes were recovered by incubation with elution buffer (1% SDS and 0.1 M NaHCO₃). Part of the eluate was saved and checked for targeted protein immunoprecipitation by immunoblotting. After reverse cross-linking at 65 °C for 4 h, the genomic DNA was purified using a DNA purification kit (Qiagen). The promoter I.f region was scanned for Pgr recruitment by PCR amplification using seven different pairs of primer sets representing the nt –1000/–1 region in ~ 150 bp intervals. PCRs were optimized according to each primer set, with the best amplification conditions for all primer sets being the following: 35 cycles (30 s denaturation at 94 °C, 1 min annealing at 57.5 °C, and 1 min elongation at 72 °C). ChIP primers were designed using the Primer3 Software (<http://frodo.wi.mit.edu/primer3/input.htm>) and were as follows: forward 5'-GGCTTCTCTTGCTACGCTGA-3' and reverse

5'-TTGTTGCTAAGAGATCAGTTGCTT-3', which amplified the nt –190/–40 region of promoter I.f to yield a 150 bp amplicon; forward 5'-ACCACAGAGAGTGAAAGTTTGAG-3' and reverse 5'-GCGTACCAAGAGAAGC-CAAT-3', which amplified the nt –334/–173 region of promoter I.f to yield a 162 bp amplicon; forward 5'-GCATTCAAGTTTGCTCAGAGG-3' and reverse 5'-TTCTTTTGATGGGGTTGCAC-3', which amplified the nt –431/–281 region of promoter I.f to yield a 151 bp amplicon; forward 5'-CCTGCCTAAAGGCTAAGATCC-3' and reverse 5'-ACACACATACACCTCTGAGCAAA-3', which amplified the nt –549/–399 region of promoter I.f to yield a 151 bp amplicon; forward 5'-CCCAAACCTTATCAACTTAGCC-3' and reverse 5'-GGATCTTAGCCTTAGGCAGGT-3', which amplified the nt –692/–528 region of promoter I.f to yield a 165 bp amplicon; forward 5'-TGAGATGTAAACATATATGTGTGTGCT-3' and reverse 5'-GGTTTGGGTTT-AGGGGAAT-3', which amplified the nt –793/–685 region of promoter I.f to yield a 109 bp amplicon; and forward 5'-AGCAGAGATGGCTTGTGGTT-3' and reverse 5'-GCACACACATAATGTTTACATCTCA-3', which amplified the nt –925/–768 region of promoter I.f to yield a 158 bp amplicon. All ChIP buffers used in this protocol contained a 1x protease inhibitor cocktail (Sigma).

Small interfering RNA

NS small interfering RNA (siRNA) and *Pgr* siRNA were purchased from Dharmacon (Chicago, IL, USA). *Pgr* knockdown was verified by real-time RT-PCR and immunoblotting. Before 1 day of transfection, hypothalamic neurons were plated in 6-well plates to achieve 50% confluency at the time of transfection and were then transfected with 100 nM NS siRNA or *Pgr* siRNA in triplicate using Lipofectamine RNAiMAX reagent (Invitrogen Life Technologies, Inc.) according to the manufacturer's protocol. After 48 h of transfection, neurons were serum-starved for 24 h, followed by treatment with 10^{-7} M R5020 for 6 h. Total mRNA was extracted for real-time RT-PCR assay, and total protein was isolated using M-PER reagent for immunoblot analysis.

Statistical analysis

Statistical analyses were performed by Welch's paired *t* test and one-way analysis of variance followed by a Tukey multiple comparisons test using the StatView 5.0 Statistical Software package (SAS Institute, Cary, NC, USA). Significance was determined at $\alpha=0.05$ and $\beta=0.20$. The values for mRNA and aromatase activity were provided as mean $\pm$ S.E.M.

Results

Aromatase, Esr1, and Pgr expression levels in hypothalamic neuronal cell lines

To verify estrogen production and estrogen and progesterone responsiveness of the study cell lines, conventional RT-PCR, real-time RT-PCR, and immunoblot analyses were performed to measure aromatase, *Esr1*, and *Pgr* mRNA or protein in N38 and N42 hypothalamic neurons. Promoter I.f-driven aromatase expression (Fig. 1A), *Pgr* expression (Fig. 1B and C), and *Esr1* expression (Fig. 1D and E) were lower in N38 hypothalamic neurons compared with N42 hypothalamic neurons. These results were reproducibly observed in three independent experiments.

Effect of the progesterone agonist, R5020, on aromatase mRNA expression and enzyme activity in N38 and N42 cell lines

Real-time RT-PCR and aromatase enzyme activity assays were performed to determine whether the progesterone agonist R5020 had an effect on aromatase mRNA expression and enzyme activity. We treated N38 and N42 hypothalamic neurons with 10^{-7} M R5020 for 6, 12, and 24 h. We observed that R5020 suppressed aromatase mRNA expression (Fig. 2A) and enzyme activity (Fig. 2B) at the 6 h time point but not at other time points in N42 hypothalamic neurons. However, aromatase RNA expression and enzyme activity at the 6 h time point in N38 hypothalamic neurons were induced by R5020 treatment (data not shown). Aromatase enzyme activity in each neuronal line treated with the aromatase enzyme inhibitor letrozole (LET) was undetectable.

The progesterone antagonist, RU486, inhibits R5020-dependent suppression of aromatase mRNA expression and enzyme activity in N42 hypothalamic neuronal cell lines

To clarify whether the effect of R5020 on aromatase mRNA expression and enzyme activity was *Pgr* dependent, we treated N42 hypothalamic neurons with the progesterone antagonist, RU486 (mifepristone). RU486 inhibited R5020-dependent suppression of aromatase mRNA expression and enzyme activity as assessed by real-time RT-PCR and aromatase enzyme activity assays (Fig. 3A and B). Aromatase activity in N42 neuronal line treated with the aromatase enzyme inhibitor LET was undetectable.

The nt -1000/-500 region of aromatase promoter I.f confers responsiveness to a progestin

To identify progestin-responsive cis-regulatory elements within the nt -1000/-1 region of aromatase

Figure 1 Promoter I.f-driven aromatase expression and *Pgr* and *Esr1* expression in N38 and N42 hypothalamic neurons. (A) Real-time RT-PCR was performed with a probe complementary to the promoter I.f to exon II junction to measure promoter I.f-driven aromatase mRNA expression. Relative units are shown as the ratio of aromatase mRNA to *Gapdh* mRNA. Results are expressed as mean $\pm$ S.E.M. of three independent experiments (** $P < 0.01$, paired t test). (B) Conventional RT-PCR and (C) immunoblotting were performed to measure *Pgr* expression in brain tissue (positive control) and N38 or N42 hypothalamic neurons. (D) Conventional RT-PCR and (E) immunoblotting were performed to measure *Esr1* expression in brain tissue (positive control) and N38 or N42 hypothalamic neurons. *Gapdh* and actin were used as loading controls. Figures represent one of the three independently performed experiments.

promoter I.f, we generated a series of truncated reporter constructs and transiently transfected them into N38 and N42 hypothalamic neurons. In N42 hypothalamic neurons, the nt -1000/-1 and the nt -700/-1 promoter-luciferase reporter constructs were responsive to R5020 (threefold repression; Fig. 4).

Pgr interacts with aromatase promoter I.f

We performed ChIP assays to demonstrate that *Pgr* is recruited to aromatase promoter I.f in N42 hypothalamic neurons. We used seven different sets of overlapping primer pairs representing the nt -1000/-1 region of promoter I.f in ~150 bp

Figure 2 The progesterone agonist, R5020, regulates aromatase mRNA expression and enzyme activity in N42, but not N38, hypothalamic neurons. (A) Real-time RT-PCR was performed to measure aromatase mRNA expression after 6, 12, and 24 h of R5020 (10^{-7} M) treatment. Aromatase mRNA levels were normalized to *Gapdh* mRNA levels. Results are expressed as mean $\pm$ S.E.M. from three independent experiments (** $P < 0.01$, paired *t* test). Aromatase activity assays were performed in (B) N42 hypothalamic neurons after 6, 12, and 24 h of R5020 (10^{-7} M) treatment. The results are expressed as mean $\pm$ S.E.M. from three independent experiments (** $P < 0.01$, paired *t* test).

intervals. The most consistent and intense *Pgr* recruitment occurred in the nt $-800/-600$ region (Fig. 5A). Under serum-starved conditions, we did not observe *Pgr* recruitment to the nt $-800/-600$ region of promoter I.f in N42 hypothalamic neurons; however, recruitment to this region was enhanced by R5020 treatment (10^{-7} M, 6 h; Fig. 5B). We applied densitometry to PCR gels to calculate fold enrichment as a ratio of recruitment of *Pgr* antibody-bound chromatin to that of nonspecific IgG-bound chromatin. We demonstrated that R5020 induced recruitment of *Pgr* by 2.1-fold to this progestin-responsive region (Fig. 5B).

***Pgr* mediates R5020-dependent suppression of aromatase mRNA expression and enzyme activity**

siRNA was used to determine whether endogenous *Pgr* in N42 hypothalamic neurons mediates R5020-dependent

suppression of aromatase mRNA expression and enzyme activity. As shown in Fig. 6A and B, siRNA knockdown of *Pgr* in N42 hypothalamic neurons abolished suppression of aromatase mRNA levels and enzyme activity by R5020 (10^{-7} M) treatment. The knockdown of *Pgr* was confirmed by real-time RT-PCR and immunoblotting (Fig. 6C and D).

Relative *Pgr* and *Esr1* expression is critical for E_2 - and progestin-dependent regulation of aromatase mRNA expression and enzyme activity

We have previously shown that E_2 induces aromatase mRNA expression and enzyme activity in N42 but not in N38 hypothalamic neurons, which express lower *Esr1* and *Pgr* levels compared with N42 hypothalamic neurons (Fig. 1C and D). To determine whether different levels of *Esr1* or *Pgr* expression affect estrogen or progestin regulation of aromatase mRNA levels and enzyme activity, we treated N38 and N42 hypothalamic neurons with 10^{-7} M R5020, 10^{-7} M E_2 , or both for 6 h. R5020 treatment induced aromatase mRNA expression (Fig. 7A) and enzyme activity (Fig. 7B) in N38 hypothalamic neurons, but suppressed them in N42 hypothalamic neurons. In addition, E_2 treatment did not change aromatase mRNA expression and enzyme activity in N38 hypothalamic neurons but induced both

Figure 3 The progesterone antagonist, RU486, reverses R5020-dependent suppression of aromatase mRNA expression and enzyme activity in N42 hypothalamic neurons. (A) Real-time RT-PCR was performed in the presence or absence of 10^{-7} M R5020 plus or minus 10^{-5} M RU486 for 6 h. Aromatase mRNA levels were normalized to *Gapdh* mRNA levels. The results are expressed as mean $\pm$ S.E.M. from three independent experiments (** $P < 0.01$, ANOVA). (B) Aromatase activity assays were performed to measure enzyme activity in the presence or absence of 10^{-7} M R5020 plus or minus 10^{-5} M RU486 for 6 h. The results are expressed as mean $\pm$ S.E.M. ($n = 3$; ** $P < 0.01$, ANOVA).

Figure 4 Cis-regulatory elements within the nt $-1000/-500$ region of aromatase promoter I.f are essential for R5020-dependent suppression of aromatase expression in N42 hypothalamic neurons. Serial deletion analysis was performed to localize the R5020-responsive regions of aromatase promoter I.f. The promoter I.f-luciferase reporter constructs were named according to the nucleotide positions of their 5'-termini. All constructs were transiently transfected into N42 hypothalamic neurons. Luciferase assays were performed a minimum of three times in the presence or absence of 10^{-7} M R5020. Mean luciferase activity of each construct is given relative to the nt $-1000/-1$ promoter-luciferase reporter construct. Summary data for three independent experiments are shown. Results are expressed as mean $\pm$ S.E.M. (** $P < 0.01$, paired t test).

aromatase expression and activity in N42 hypothalamic neurons. Moreover, when treated with R5020 and E_2 together, aromatase mRNA expression and enzyme activity were induced in both hypothalamic neuronal lines (Fig. 7A and B). Aromatase enzyme activity in each neuronal line treated with the aromatase enzyme inhibitor LET was undetectable.

Discussion

Aromatase deficiency in humans disrupts normal gonadotropin regulation and libido (Jones *et al.* 2006). Likewise, ArKO mice exhibit disrupted reproductive behavior, including defective lordosis (females), low frequency of copulatory behavior (males; Bakker *et al.* 2003), reduced sexual interest in the opposite sex, and deficiency in pup care (Bakker *et al.* 2002, 2003, Matsumoto *et al.* 2003a,b). The gonadotropin secretion profile is also disrupted in ArKO mice, which have increased LH, FSH, testosterone, and prolactin levels, and decreased E_2 , but unchanged dihydrotestosterone and progesterone (Fisher *et al.* 1998, Jones *et al.* 2006).

Aromatase is primarily expressed in gonads and the brain (Simpson 2004a, Bulun *et al.* 2005). Loss of local E_2 synthesis in the brain likely leads to the hormonal and behavioral phenotypes observed in aromatase-deficient humans and mice. In fact, alterations in brain aromatase expression have been directly linked to changes in sexual behavior in birds, rodents, sheep, and humans (Balthazart *et al.* 2003, Matsumoto *et al.* 2003a,b, Jones *et al.* 2006, Perkins & Roselli 2007).

The role of progesterone in females appears similar in a variety of species, particularly for maternal and reproductive behavior. However, progesterone is capable of protecting females from the masculinizing and defeminizing effects of testosterone (Sodersten & Hansen 1977, Hansen & Sodersten 1978). However, progesterone treatment suppresses mounting and intromissive behavior in males (Michael *et al.* 1968, DeBold *et al.* 1978, Witt *et al.* 1995).

Gonadectomy reduces aromatase expression and enzyme activity in the brain in parallel with a recession in gender-specific sexual behavior of both sexes (Jakab *et al.* 1993, Iivonen *et al.* 2006). However, treatment of gonadectomized animals with progesterone markedly restores female-specific but not male-specific sexual behavior (Sodersten & Hansen 1977, Hansen & Sodersten 1978, Witt *et al.* 1995). Moreover, treatments of intact animals with the progesterone antagonist, RU486, or antisense oligonucleotides to *Pgr* increased the number of mounts and intromissions in males but

Figure 5 ChIP analysis reveals Pgr recruitment to aromatase promoter I.f in N42 hypothalamic neurons. We performed ChIP assays using N42 hypothalamic neuron extracts treated with vehicle or 10^{-7} M R5020 for 6 h employing different sets of primers; the most consistent recruitment was observed with the primers (underlined) in (A). Sonicated cell supernatant was used as input DNA (positive control). Pre-cleared chromatin was used for immunoprecipitation reactions with a rabbit polyclonal antibody directed against human PGR and normal rabbit IgG. The images in (B) are from one of the five independently performed experiments. The bars represent densitometric measurements of the ratio of enrichment of DNA precipitated using an antibody against Pgr versus nonspecific IgG. The bars were generated from repeated experiments.

Figure 6 Pgr mediates R5020-dependent suppression of aromatase mRNA expression in N42 hypothalamic neurons. N42 hypothalamic neurons were transfected with *Pgr* siRNA or with non-specific siRNA as a negative control (NS) and cultured for an additional 48 h. (A) Aromatase mRNA levels were analyzed by real-time RT-PCR in the presence or absence of R5020 (10^{-7} M, 6 h). Aromatase mRNA levels were normalized to *Gapdh* mRNA levels. The results are expressed as mean $\pm$ S.E.M. from three independent experiments (** $P < 0.01$, ANOVA). (B) Aromatase activity assays were performed in the presence or absence of R5020 (10^{-7} M, 6 h). The results are expressed as mean $\pm$ S.E.M. from three independent experiments (** $P < 0.01$, ANOVA). Aromatase enzyme activity in N42 neuronal line treated with the aromatase enzyme inhibitor letrozole was undetectable (data not shown). (C) Real-time RT-PCR and (D) immunoblotting were performed to measure *Pgr* mRNA and protein levels respectively. Actin was used as a loading control. The immunoblot represents one of the three independently performed experiments. Real-time RT-PCR results are expressed as mean $\pm$ S.E.M. from three independent experiments (** $P < 0.01$, ANOVA) and *Gapdh* was used as a loading control. Representative results from three independent experiments are shown.

decreased the frequency of lordosis in females (Pollio *et al.* 1993, Mani *et al.* 1994). Therefore, the actions of progesterone are necessary for the proceptive components of female sexual behavior, although the mechanism of progesterone action is not fully understood.

The reproductive phenotypes manifested in ArKO male mice are opposite to those observed in progesterone receptor knockout (PRKO) male mice, in which most measures of male sexual behavior are enhanced, suggesting that aromatase and *Pgr* are important in mediating opposite aspects of reproductive behavior controlled by the brain, in particular, by the hypothalamus in males (Matsumoto *et al.* 2003a, Schneider *et al.* 2005). However, PRKO female mice are sterile with impaired maternal and reproductive behavior and neuroendocrine and gonadotropin regulation, suggesting a prominent role of *Pgr* in feminization rather than masculinization process (Conneely *et al.* 2001).

Taken together, the roles that are attributed to progesterone in gender-specific sexual behavior mostly favors the feminization process both in males and in females, which is in contrast to what has been observed in male mice with inactivated *Esr1* or *Cyp19a1* (aromatase), suggesting that *Pgr* and *Esr1*/aromatase may regulate opposite aspects of reproductive behavior, particularly in males.

In the hypothalamic tissue of neonatal males, progesterone does not affect the accumulation of testosterone, but completely prevents the formation of E_2 , suggesting that progesterone may inhibit hypothalamic aromatase expression and/or enzyme activity in the neonatal brain (Rhoda *et al.* 1987). Consistent with these findings, we found that progesterone treatment suppressed aromatase mRNA expression and enzyme activity (Fig. 2).

The varying intracellular concentration of progesterone and possible alterations in the composition of the transcriptional complex at promoter I.f over time may explain the mechanism of biphasic response observed in this study. These considerations will be investigated as future directions.

Progesterone can also inhibit *Esr1* expression and E_2 can induce the expression of both *Pgr* isoforms (Evans & Leavitt 1980, Smanik *et al.* 1983). We observed a slight decrease in *Esr1*, but not *Esr2* (also known as $ER\beta$) expression levels upon R5020 treatment in hypothalamic cells (data not shown). This may represent an alternative mechanism whereby progesterone regulates aromatase expression in hypothalamic neurons.

In parallel, it is known that E_2 induces *Pgr* expression and inhibition of aromatase suppresses *Pgr* expression in the brain, suggesting a negative feedback mechanism between progesterone and aromatase (Ing & Tornesi

Figure 7 Combinatory effects of E_2 and R5020 on aromatase are cell-type specific. (A) Real-time RT-PCR was performed to measure aromatase mRNA expression after 6 h of R5020 (10^{-7} M) or E_2 (10^{-7} M) or R5020 (10^{-7} M) + E_2 (10^{-7} M) treatment in both N38 and N42 hypothalamic neurons. Aromatase mRNA levels were normalized to *Gapdh* mRNA levels. Results are expressed as mean $\pm$ S.E.M. from three independent experiments (** $P < 0.01$, paired *t* test). (B) Aromatase activity assays were performed in both N38 and N42 hypothalamic neurons following the described treatment procedure in (A). The results are expressed as mean $\pm$ S.E.M. from three independent experiments (** $P < 0.01$, paired *t* test).

1997, Prange-Kiel *et al.* 2001). Consistent with this hypothesis, we found that Pgr interacts with the aromatase I.f promoter in the presence but not in the absence of R5020, with the highest intensity at the nt $-800/-600$ region of promoter I.f (Fig. 4).

The perinatal rodent brain expresses all the enzymes required for the *de novo* synthesis of progesterone from cholesterol, potentially producing locally high concentrations of this steroid. E_2 concentration in certain brain areas, such as the hypothalamus, preoptic area, and the hippocampus, is higher than the plasma E_2 levels (Hojo *et al.* 2004, Rune *et al.* 2006). Even though the total amount of E_2 synthesized by these brain sites seems to be low, the achieved local tissue concentrations are probably quite high and exert significant biologic influence locally (Simpson *et al.* 1999, Prange-Kiel *et al.* 2003). The same phenomenon could also be true for progesterone adding another layer of complexity to the phenomenally complex mechanism involving estrogen and progesterone production and action in various regions.

Various studies investigating the effects of progesterone on reproductive behavior used supraphysiological levels of progesterone that would greatly exceed its levels accounted for by ovarian secretion (Erpino 1975, Connolly & Resko 1989). However, the endogenous levels of progesterone can be locally regulated in specific tissues or regions of the brain (Le Goascogne *et al.* 2000, Tagawa *et al.* 2006). In our experiments, aromatase regulation by 10^{-7} M R5020, which may represent the cellular progesterone concentration in a particular brain region, cannot be generalized to the effects of this steroid to the entire brain tissue. As a further twist, both the Pgr and the *Esr1* levels in N42 neurons are higher than those in N38 hypothalamic neurons, which are possibly accounted for differential responsiveness of these cells to progesterone. Differential responsiveness of N42 and N38 cells to R5020 with respect to aromatase expression was a function of Pgr levels and consistent across various experiments because knockdown of Pgr in N42 cells converted an inhibitory response to a stimulatory one (Figs 1, 6 and 7). We speculate that, *in vivo*, the varying levels of Pgr and *Esr1* in specific brain cells would be key determinants of progesterone responsiveness.

The developmental and tissue-, region-, and neuron-specific expression of aromatase in the brain has been reported to be transcriptionally regulated through promoter I.f (Honda *et al.* 1999, Nausch *et al.* 2007). There are several potential cis-regulatory elements, which may be important for promoter regulation, residing in the 5'-flanking region of promoter I.f including PRE half sites. In our experiments, R5020 significantly suppressed luciferase activity of the nt $-1000/-1$ and nt $-700/-1$ promoter-reporter constructs but not other promoter-reporter luciferase constructs in N42 hypothalamic neurons (Fig. 4). We did not observe suppression of luciferase activity of the promoter-luciferase reporter constructs on R5020 treatment in N38 hypothalamic neurons, due possibly

to higher Pgr expression in N42 hypothalamic neurons compared with N38 hypothalamic neurons.

Aromatase in the fetal brain is also proposed to be regulated via sex hormones, specifically E₂ that has both stimulatory and inhibitory effects on aromatase expression and enzyme activity in the brain (Iivonen *et al.* 2006). Our group has recently showed that Esr1 mediates E₂ actions on aromatase expression and enzyme activity through its brain-specific promoter in embryonic hypothalamic neuronal cell lines (Yilmaz *et al.* 2009). Progesterone has also been implicated in the regulation aromatase expression and enzyme activity (Rhoda *et al.* 1987, Ing & Tornesi 1997, Prange-Kiel *et al.* 2001, Trainor *et al.* 2003). Interestingly, Brock *et al.* (2010) has recently showed that Pgr and aromatase expression is conversely related in the mouse hypothalamus, suggesting a feedback loop between progesterone and aromatase. In this study, we showed that progesterone enhances recruitment of Pgr to specific regions of the promoter I.f of *Cyp19a1* and regulates aromatase expression in hypothalamic neurons.

We identified several progesterone response elements (half sites) in both nt -1000/-700 and nt -700/-500, with nt -800/-600 having the highest intensity of Pgr recruitment sites as assessed by ChIP. Furthermore, the effect of E₂ and progesterone on aromatase mRNA expression and enzyme activity is Esr1/Pgr expression dependent (Fig. 7), with higher receptor expression favoring induction of aromatase on E₂ treatment and suppression on R5020 treatment, and with lower receptor expression favoring induction on R5020 treatment and no alteration on E₂ treatment.

Taken together, we demonstrated the mechanism of a Pgr action in mediating progesterone-dependent regulation of aromatase expression and enzyme activity in hypothalamic neurons in a time-, concentration-, and Esr1/Pgr expression-dependent manner. Our future directions include verification of these findings using *in vivo* models. We also plan to further investigate the *in vitro* and *in vivo* effects of other hormones and metabolites on aromatase expression and activity to elucidate the molecular mechanisms mediating local E₂ synthesis in the hypothalamus and other regions of the brain.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was funded, in part, by grants from the NIH (HD38691, HD57877) and Friends of Prentice.

Author contribution statement

Dr M B Y performed the key experiments and wrote the draft manuscript. Dr H Z and C B performed other important experiments. Drs A W and S E B significantly contributed to the experimental design.

Acknowledgements

We acknowledge Dr Zhihong Lin and Scott Reierstad for their technical contributions to the ChIP experiments.

References

- Abdelgadir SE, Resko JA, Ojeda SR, Lephart ED, McPhaul MJ & Roselli CE 1994 Androgens regulate aromatase cytochrome P450 messenger ribonucleic acid in rat brain. *Endocrinology* **135** 395–401. (doi:10.1210/en.135.1.395)
- Bakker J, Honda S, Harada N & Balthazart J 2002 The aromatase knockout mouse provides new evidence that estradiol is required during development in the female for the expression of sociosexual behaviors in adulthood. *Journal of Neuroscience* **22** 9104–9112.
- Bakker J, Honda S, Harada N & Balthazart J 2003 The aromatase knockout (ArKO) mouse provides new evidence that estrogens are required for the development of the female brain. *Annals of the New York Academy of Sciences* **1007** 251–262. (doi:10.1196/annals.1286.024)
- Bakker J, Baillien M, Honda S, Harada N & Balthazart J 2004a Relationships between aromatase activity in the brain and gonads and behavioural deficits in homozygous and heterozygous aromatase knockout mice. *Journal of Neuroendocrinology* **16** 483–490. (doi:10.1111/j.1365-2826.2004.01191.x)
- Bakker J, Honda S, Harada N & Balthazart J 2004b Restoration of male sexual behavior by adult exogenous estrogens in male aromatase knockout mice. *Hormones and Behavior* **46** 1–10. (doi:10.1016/j.yhbeh.2004.02.003)
- Balthazart J, Baillien M & Ball GF 2001 Phosphorylation processes mediate rapid changes of brain aromatase activity. *Journal of Steroid Biochemistry and Molecular Biology* **79** 261–277. (doi:10.1016/S0960-0760(01)00143-1)
- Balthazart J, Baillien M, Charlier TD, Cornil CA & Ball GF 2003 Multiple mechanisms control brain aromatase activity at the genomic and non-genomic level. *Journal of Steroid Biochemistry and Molecular Biology* **86** 367–379. (doi:10.1016/S0960-0760(03)00346-7)
- Brock O, Douhard Q, Baum MJ & Bakker J 2010 Reduced prepubertal expression of progesterone receptor in the hypothalamus of female aromatase knockout mice. *Endocrinology* **151** 1814–1821. (doi:10.1210/en.2009.1379)
- Bulun SE, Lin Z, Imir G, Amin S, Demura M, Yilmaz B, Martin R, Utsunomiya H, Thung S, Gurates B *et al.* 2005 Regulation of aromatase expression in estrogen-responsive breast and uterine disease: from bench to treatment. *Pharmacological Reviews* **57** 359–383. (doi:10.1124/pr.57.3.6)
- Chow JD, Simpson ER & Boon WC 2009 Alternative 5'-untranslated first exons of the mouse *Cyp19A1* (aromatase) gene. *Journal of Steroid Biochemistry and Molecular Biology* **115** 115–125. (doi:10.1016/j.jsbmb.2009.03.010)
- Conneely OM, Mulac-Jericevic B, Lydon JP & De Mayo FJ 2001 Reproductive functions of the progesterone receptor isoforms: lessons from knock-out mice. *Molecular and Cellular Endocrinology* **179** 97–103. (doi:10.1016/S0303-7207(01)00465-8)
- Connolly PB & Resko JA 1989 Progestins affect reproductive behavior and androgen receptor dynamics in male guinea pig brain. *Brain Research* **503** 312–316. (doi:10.1016/0006-8993(89)91681-8)

- DeBold JF, Morris JL & Clemens LG 1978 The inhibitory actions of progesterone: effects on male and female sexual behavior of the hamster. *Hormones and Behavior* **11** 28–41. (doi:10.1016/0018-506X(78)90056-9)
- Erpino MJ 1975 Androgen-induced aggression in neonatally androgenized female mice: inhibition by progesterone. *Hormones and Behavior* **6** 149–157. (doi:10.1016/0018-506X(75)90030-6)
- Evans RW & Leavitt WW 1980 Progesterone inhibition of uterine nuclear estrogen receptor: dependence on RNA and protein synthesis. *PNAS* **77** 5856–5860. (doi:10.1073/pnas.77.10.5856)
- Fisher CR, Graves KH, Parlow AF & Simpson ER 1998 Characterization of mice deficient in aromatase (ArKO) because of targeted disruption of the cyp19 gene. *PNAS* **95** 6965–6970. (doi:10.1073/pnas.95.12.6965)
- Fortune JE & Vincent SE 1983 Progesterone inhibits the induction of aromatase activity in rat granulosa cells *in vitro*. *Biology of Reproduction* **28** 1078–1089. (doi:10.1095/biolreprod28.5.1078)
- Golovine K, Schwerin M & Vanselow J 2003 Three different promoters control expression of the aromatase cytochrome p450 gene (cyp19) in mouse gonads and brain. *Biology of Reproduction* **68** 978–984. (doi:10.1095/biolreprod.102.008037)
- Hansen S & Sodersten P 1978 Effects of subcutaneous implants of progesterone on the induction and duration of sexual receptivity in ovariectomized rats. *Journal of Endocrinology* **77** 373–379. (doi:10.1677/joe.0.0770373)
- Hoyo Y, Hattori TA, Enami T, Furukawa A, Suzuki K, Ishii HT, Mukai H, Morrison JH, Janssen WG, Kominami S *et al.* 2004 Adult male rat hippocampus synthesizes estradiol from pregnenolone by cytochromes P45017alpha and P450 aromatase localized in neurons. *PNAS* **101** 865–870. (doi:10.1073/pnas.2630225100)
- Honda S, Harada N & Takagi Y 1996 The alternative exons 1 of the mouse aromatase cytochrome P-450 gene. *Biochimica et Biophysica Acta* **1305** 145–150. (doi:10.1016/0167-4781(95)00200-6)
- Honda S, Harada N, Abe-Dohmae S & Takagi Y 1999 Identification of cis-acting elements in the proximal promoter region for brain-specific exon 1 of the mouse aromatase gene. *Brain Research. Molecular Brain Research* **66** 122–132. (doi:10.1016/S0169-328X(99)00017-0)
- Iivonen S, Heikkinen T, Puolivali J, Helisalmi S, Hiltunen M, Soininen H & Tanila H 2006 Effects of estradiol on spatial learning, hippocampal cytochrome P450 19, and estrogen alpha and beta mRNA levels in ovariectomized female mice. *Neuroscience* **137** 1143–1152. (doi:10.1016/j.neuroscience.2005.10.023)
- Ing NH & Tornesi MB 1997 Estradiol up-regulates estrogen receptor and progesterone receptor gene expression in specific ovine uterine cells. *Biology of Reproduction* **56** 1205–1215. (doi:10.1095/biolreprod56.5.1205)
- Jakab RL, Horvath TL, Leranath C, Harada N & Naftolin F 1993 Aromatase immunoreactivity in the rat brain: gonadectomy-sensitive hypothalamic neurons and an unresponsive "limbic ring" of the lateral septum-bed nucleus-amygdala complex. *Journal of Steroid Biochemistry and Molecular Biology* **44** 481–498. (doi:10.1016/0960-0760(93)90253-S)
- Jones ME, Boon WC, Proietto J & Simpson ER 2006 Of mice and men: the evolving phenotype of aromatase deficiency. *Trends in Endocrinology and Metabolism* **17** 55–64. (doi:10.1016/j.tem.2006.01.004)
- Keefe DL 2002 Sex hormones and neural mechanisms. *Archives of Sexual Behavior* **31** 401–403. (doi:10.1023/A:1019883923346)
- Lavaque E, Mayen A, Azcoitia I, Tena-Sempere M & Garcia-Segura LM 2006 Sex differences, developmental changes, response to injury and cAMP regulation of the mRNA levels of steroidogenic acute regulatory protein, cytochrome p450sc, and aromatase in the olivocerebellar system. *Journal of Neurobiology* **66** 308–318. (doi:10.1002/neu.20221)
- Le Goascogne C, Eychenne B, Tonon MC, Lachapelle F, Baumann N & Robel P 2000 Neurosteroid progesterone is up-regulated in the brain of jimpy and shiverer mice. *Glia* **29** 14–24. (doi:10.1002/(SICI)1098-1136(20000101)29:1<14::AID-GLIA2>3.0.CO;2-N)
- Lephart ED 1996 A review of brain aromatase cytochrome P450. *Brain Research. Brain Research Reviews* **22** 1–26. (doi:10.1016/0165-0173(96)00002-1)
- Lephart ED 1997 Molecular aspects of brain aromatase cytochrome P450. *Journal of Steroid Biochemistry and Molecular Biology* **61** 375–380. (doi:10.1016/S0960-0760(97)80035-0)
- Lephart ED, Butler PC, Mills RH, Jacobson NA, Ladle DR & Bloch GJ 1998 Effects of testosterone and progesterone on brain 5alpha-reductase and aromatase in Long-Evans males and comparison of aromatase in Long-Evans vs. Sprague-Dawley rats. *Brain Research* **789** 327–330. (doi:10.1016/S0006-8993(97)01553-9)
- Levine JE, Chappell PE, Schneider JS, Sleiter NC & Szabo M 2001 Progesterone receptors as neuroendocrine integrators. *Frontiers in Neuroendocrinology* **22** 69–106. (doi:10.1006/frnc.2001.0210)
- Mani SK, Blaustein JD, Allen JM, Law SW, O'Malley BW & Clark JH 1994 Inhibition of rat sexual behavior by antisense oligonucleotides to the progesterone receptor. *Endocrinology* **135** 1409–1414. (doi:10.1210/en.135.4.1409)
- Matsumoto T, Honda S & Harada N 2003a Alteration in sex-specific behaviors in male mice lacking the aromatase gene. *Neuroendocrinology* **77** 416–424. (doi:10.1159/000071313)
- Matsumoto T, Honda S & Harada N 2003b Neurological effects of aromatase deficiency in the mouse. *Journal of Steroid Biochemistry and Molecular Biology* **86** 357–365. (doi:10.1016/S0960-0760(03)00345-5)
- Michael RP, Saayman GS & Zump D 1968 The suppression of mounting behaviour and ejaculation in male rhesus monkeys (*Macaca mulatta*) by administration of progesterone to their female partners. *Journal of Endocrinology* **41** 421–431. (doi:10.1677/joe.0.0410421)
- Nausch N, Manteuffel G & Vanselow J 2007 0.2 kb promoter sequence of the murine Cyp19 gene target beta-galactosidase expression to specific brain areas of transgenic mice. *Journal of Steroid Biochemistry and Molecular Biology* **103** 119–128. (doi:10.1016/j.jsbmb.2006.08.007)
- Negri-Cesi P, Colciago A, Motta M, Martini L & Celotti F 2001 Aromatase expression and activity in male and female cultured rat hypothalamic neurons: effect of androgens. *Molecular and Cellular Endocrinology* **178** 1–10. (doi:10.1016/S0303-7207(01)00442-7)
- Perkins A & Roselli CE 2007 The ram as a model for behavioral neuroendocrinology. *Hormones and Behavior* **52** 70–77. (doi:10.1016/j.yhbeh.2007.03.016)
- Peterson RS, Yarram L, Schlenger BA & Saldanha CJ 2005 Aromatase is pre-synaptic and sexually dimorphic in the adult zebra finch brain. *Proceedings. Biological Sciences/The Royal Society* **272** 2089–2096. (doi:10.1098/rspb.2005.3181)
- Pluchino N, Luisi M, Lenzi E, Centofanti M, Begliuomini S, Freschi L, Ninni F & Genazzani AR 2006 Progesterone and progestins: effects on brain, allopregnanolone and beta-endorphin. *Journal of Steroid Biochemistry and Molecular Biology* **102** 205–213. (doi:10.1016/j.jsbmb.2006.09.023)
- Pollio G, Xue P, Zanisi M, Nicolin A & Maggi A 1993 Antisense oligonucleotide blocks progesterone-induced lordosis behavior in ovariectomized rats. *Brain Research. Molecular Brain Research* **19** 135–139. (doi:10.1016/0169-328X(93)90158-L)
- Prange-Kiel J, Rune GM, Zwirner M, Wallwiener D & Kiesel L 2001 Regulation of estrogen receptor alpha and progesterone receptor (isoform A and B) expression in cultured human endometrial cells. *Experimental and Clinical Endocrinology and Diabetes* **109** 231–237. (doi:10.1055/s-2001-15111)
- Prange-Kiel J, Wehrenberg U, Jarry H & Rune GM 2003 Para/auto-crine regulation of estrogen receptors in hippocampal neurons. *Hippocampus* **13** 226–234. (doi:10.1002/hipo.10075)

- Rhoda J, Valens M, Edwards DA & Roffi J 1987 Effect of progesterone on the testosterone and estradiol levels in the hypothalamus of neonatal rats. *Biology of the Neonate* **51** 255–259. (doi:10.1159/000242661)
- Roselli CE, Abdelgadir SE & Resko JA 1997 Regulation of aromatase gene expression in the adult rat brain. *Brain Research Bulletin* **44** 351–357. (doi:10.1016/S0361-9230(97)00214-1)
- Rune GM, Lohse C, Prange-Kiel J, Fester L & Frotscher M 2006 Synaptic plasticity in the hippocampus: effects of estrogen from the gonads or hippocampus? *Neurochemical Research* **31** 145–155. (doi:10.1007/s11064-005-9004-8)
- Schneider JS, Burgess C, Sleiter NC, DonCarlos LL, Lydon JP, O'Malley B & Levine JE 2005 Enhanced sexual behaviors and androgen receptor immunoreactivity in the male progesterone receptor knockout mouse. *Endocrinology* **146** 4340–4348. (doi:10.1210/en.2005-0490)
- Sharma TP, Blache D, Roselli CE & Martin GB 2004 Distribution of aromatase activity in brain and peripheral tissues of male sheep: effect of nutrition. *Reproduction, Fertility, and Development* **16** 709–715. (doi:10.1071/RD04018)
- Shozu M, Sebastian S, Takayama K, Hsu WT, Schultz RA, Neely K, Bryant M & Bulun SE 2003 Estrogen excess associated with novel gain-of-function mutations affecting the aromatase gene. *New England Journal of Medicine* **348** 1855–1865. (doi:10.1056/NEJMoa021559)
- Simpson ER 2004a Aromatase: biologic relevance of tissue-specific expression. *Seminars in Reproductive Medicine* **22** 11–23. (doi:10.1055/s-2004-823023)
- Simpson ER 2004b Models of aromatase insufficiency. *Seminars in Reproductive Medicine* **22** 25–30. (doi:10.1055/s-2004-823024)
- Simpson ER & Davis SR 2001 Minireview: aromatase and the regulation of estrogen biosynthesis – some new perspectives. *Endocrinology* **142** 4589–4594. (doi:10.1210/en.142.11.4589)
- Simpson E, Rubin G, Clyne C, Robertson K, O'Donnell L, Davis S & Jones M 1999 Local estrogen biosynthesis in males and females. *Endocrine-Related Cancer* **6** 131–137. (doi:10.1677/erc.0.0060131)
- Smanik EJ, Young HK, Muldoon TG & Mahesh VB 1983 Analysis of the effect of progesterone *in vivo* on estrogen receptor distribution in the rat anterior pituitary and hypothalamus. *Endocrinology* **113** 15–22. (doi:10.1210/endo-113-1-15)
- Sodersten P & Hansen S 1977 Effects of oestradiol and progesterone on the induction and duration of sexual receptivity in cyclic female rats. *Journal of Endocrinology* **74** 477–485. (doi:10.1677/joe.0.0740477)
- Tagawa N, Sugimoto Y, Yamada J & Kobayashi Y 2006 Strain differences of neurosteroid levels in mouse brain. *Steroids* **71** 776–784. (doi:10.1016/j.steroids.2006.05.008)
- Trainor BC, Bird IM, Alday NA, Schlinger BA & Marler CA 2003 Variation in aromatase activity in the medial preoptic area and plasma progesterone is associated with the onset of paternal behavior. *Neuroendocrinology* **78** 36–44. (doi:10.1159/000071704)
- Voigt C, Ball GF & Balthazart J 2007 Neuroanatomical specificity of sex differences in expression of aromatase mRNA in the quail brain. *Journal of Chemical Neuroanatomy* **33** 75–86. (doi:10.1016/j.jchem-neu.2006.12.004)
- Wagner CK 2006 The many faces of progesterone: a role in adult and developing male brain. *Frontiers in Neuroendocrinology* **27** 340–359. (doi:10.1016/j.yfrne.2006.07.003)
- Witt DM, Young LJ & Crews D 1995 Progesterone modulation of androgen-dependent sexual behavior in male rats. *Physiology and Behavior* **57** 307–313. (doi:10.1016/0031-9384(94)00247-3)
- Yilmaz MB, Wolfe A, Cheng YH, Glidewell-Kenney C, Jameson JL & Bulun SE 2009 Aromatase promoter 1f is regulated by estrogen receptor alpha (ESR1) in mouse hypothalamic neuronal cell lines. *Biology of Reproduction* **81** 956–965. (doi:10.1095/biolreprod.109.077206)
- Zhao C, Fujinaga R, Tanaka M, Yanai A, Nakahama K & Shinoda K 2007 Region-specific expression and sex-steroidal regulation on aromatase and its mRNA in the male rat brain: immunohistochemical and *in situ* hybridization analyses. *Journal of Comparative Neurology* **500** 557–573. (doi:10.1002/cne.21193)
- Zhao H, Innes J, Brooks DC, Reierstad S, Yilmaz MB, Lin Z & Bulun SE 2009 A novel promoter controls Cyp19a1 gene expression in mouse adipose tissue. *Reproductive Biology and Endocrinology* **24** 7–37. (doi:10.1186/1477-7827-7-37)

Received in final form 2 May 2011

Accepted 27 May 2011

Made available online as an Accepted Preprint 31 May 2011