The emerging role of the mitochondrial-derived peptide humanin in stress resistance

Kelvin Yen, Changan Lee, Hemal Mehta and Pinchas Cohen

USC Davis School of Gerontology, University of Southern California, Los Angeles, California 90089-0191, USA

Correspondence should be addressed to P Cohen
Email
hassy@usc.edu

Abstract

The discovery of humanin, a novel, mitochondrial-derived peptide, has created a potentially new category of biologically active peptide. As more research unravels the endogenous role of humanin as well as its potential pharmacological use, its role in stress resistance has become clearer. Humanin protects cells from oxidative stress, serum starvation, hypoxia, and other insults *in vitro* and also improves cardiovascular disease as well as Alzheimer's disease *in vivo*. In this review, we discuss the emerging role of humanin in stress resistance and its proposed mechanism of action.

Key Words

- ▶ Apoptosis
- ▶ Cardiovascular
- ▶ Diabetes (all)
- ▶ Signal transduction

Journal of Molecular Endocrinology (2013) 50, R11–R19

An overview of mitochondrial function

Mitochondria are generally considered the powerhouse of the cell as the majority of cellular ATP production occurs there. It is within the mitochondria that the Krebs cycle occurs, setting up an electrochemical gradient that will be used to produce ATP (Salway 1994). In addition to its role in energy production and metabolism, mitochondria play a major role in apoptosis, oxidative stress, and calcium sequestration/signaling.

The central role that mitochondria play in apoptosis has been enumerated on several occasions (reviewed in Wang (2001) and Kantari & Walczak (2011)). Mitochondria play a critical role in the intrinsic apoptotic pathway that is triggered by cellular damage of which some cells are dependent on the mitochondria for apoptosis (type 2), while others are not (type 1) (Kantari & Walczak 2011). Furthermore, many nucleus-encoded mitochondrial proteins directly activate apoptosis when they are released from the mitochondria into the cytosol or nucleus. In addition to apoptosis,

mitochondria are also believed to be the largest generator of free radicals in the cell (reviewed in Balaban *et al.* (2005)). Free radicals are the major source of oxidative stress within a cell and these free radicals can oxidize and damage proteins, lipids, and DNA. Besides the physiological role that certain free radicals play, increased oxidative stress has been associated with a number of different diseases, although a beneficial effect of transient oxidative stress has been reported as well (Ristow *et al.* 2009, Ristow & Zarse 2010).

Mitochondria possess an elaborate system for facilitating the transport of Ca^{2+} in and out of their inner membrane, which plays an important role in cytoplasmic Ca^{2+} signaling, ATP production, and hormone metabolism. Deregulation of the mitochondrial Ca^{2+} homeostasis can trigger cell death and disrupt the function of intra-mitochondrial enzymes. Mitochondrial Ca^{2+} uptake and release differ from other membrane-bound organelles in that Ca^{2+} uptake is not dependent on ATP and Ca^{2+} release uses gated channels and Na^{+} or $\text{H}^{+}/\text{Ca}^{2+}$ exchangers (reviewed in Drago *et al.* (2011)).

Communication from the cell to the mitochondria is well established, but how mitochondria signal to the cell is relatively less understood. In addition to the chemical signaling via Ca^{2+} and reactive oxygen species (ROS), in simpler organisms like yeast, mitochondria communicate with the cell via the retrograde signaling (RTG) pathway. In animal cells, many of the genes of the RTG pathway do not exist, suggesting that animal mitochondria have an alternative method for signaling the cell. Two recent papers suggest that the mitochondrial unfolded protein response (UPR) may be an alternative way that mitochondria signal to the cell. The first paper used *Caenorhabditis elegans* to demonstrate that tissue-specific mitochondrial dysfunction lead to a life-increasing, endocrine response and that this increase was dependent on the mitochondrial UPR (Durieux *et al.* 2011). Another paper showed that mitochondria undergoing temperature stress release degraded protein peptides that trigger mitochondrial UPR (Haynes *et al.* 2010). Humanin, the topic of this review, is another peptide that possibly acts as a mitochondrial autocrine, paracrine, and endocrine signal.

The discovery of humanin

Approximately 10 years ago, the Nishimoto Lab (Hashimoto *et al.* 2001b) discovered humanin (MTRNR2) while searching for survival factors in the unaffected brain section of an Alzheimer's patient. Using a cDNA library, they searched for genes that could resist apoptosis induced by a mutant form of amyloid precursor protein (APP V642I), which is a gene that is mutated in one form of familial Alzheimer's disease (AD). They further showed that humanin could protect against other familial AD genes but could not protect against a Huntington's disease model or amyotrophic lateral sclerosis model (Hashimoto *et al.* 2001a,b). This suggested that humanin conferred specific protection to AD models and did not merely protect cells from all apoptotic events.

Humanin (HN/MTRNR2) has also been cloned as a BAX partner that could suppress apoptosis caused by staurosporine, serum deprivation, or u.v. irradiation (Guo *et al.* 2003). Using co-immunoprecipitation experiments, humanin specifically bound to BAX, tBID, and BimEL (BCL2L11), but not other BCL2 family proteins such as BCL2 and BCL-B (BCL2L10) (Luciano *et al.* 2005, Zhai *et al.* 2005). Furthermore, the anti-apoptotic effect of humanin was specific to BAX-dependent apoptosis, as apoptosis by a BAX-independent stimulus, such as the use of tumor necrosis factor, was not suppressed.

Humanin also interacts with insulin-like growth factor-binding protein 3 (IGFBP3), where it can block IGFBP3-induced apoptosis in glial cell lines but not in neuronal cell lines or primary neurons (Ikonen *et al.* 2003). Although humanin can block the apoptotic effects of IGFBP3 in glial cells, unexpectedly, IGFBP3 and humanin have a positive, synergistic effect on protecting neurons from amyloid- β -induced apoptosis (Ikonen *et al.* 2003).

The discovery of humanin is interesting for many reasons. It is a small, secreted, 24 or 21 amino acid peptide, depending on cytoplasmic or mitochondrial translation respectively, with biological activity that does not have a signal peptide for secretion but may act as a signal peptide itself (Yamagishi *et al.* 2003). More interesting is the fact that it is the first new peptide discovered within the mitochondrial genome since its complete sequencing in 1981 (discussed in further detail below) (Anderson *et al.* 1981). The mitochondrial genome itself is extremely small and compact (16 569 nucleotides in humans), with no known introns and very few non-coding nucleotides between adjacent genes; traditionally, it has been known to code for only 2 rRNAs, 13 mRNAs, and 22 tRNAs. Humanin is an open reading frame (ORF) found within the 16s rRNA gene (Fig. 1). Like a Russian nesting doll, humanin is a gene within a gene within a genome of an organelle within a cell and may represent the first peptide of a new class of mitochondrial-derived peptides (MDP).

Humanin action in stress

Alzheimer's disease

As mentioned earlier, humanin was discovered during a search for neurosurvival factors in unaffected areas of an AD patient's brain. Thus, it is not surprising that the best-studied aspect of humanin biology is in its beneficial effects in protecting against AD-related stress. These initial studies were first performed in cell culture and then followed by *in vivo* studies using both pharmacological mimetics of AD as well as mutant APP. The most recent studies have now started using different transgenic models of AD to examine the potential use of humanin to ameliorate AD.

The earliest humanin study showed that it could protect cells from apoptosis induced by amyloid β ($\text{A}\beta$)_{1–43} and a follow-up study went even further showing that humanin could protect against other mutated genes that cause familial AD such as APP, presenilin 1, and presenilin 2 (Hashimoto *et al.* 2001a,b). Although other factors such as IGF1 and basic FGF could protect neurons from ($\text{A}\beta$)_{1–43},

Figure 1

The humanin open reading frame (ORF). Humanin is encoded by an ORF within the gene for the 16s ribosomal subunit within the mitochondrial genome.

only humanin could protect against all the tested AD genes (Hashimoto *et al.* 2001a).

The first *in vivo* study to examine the physiological potential of humanin used scopolamine, a muscarinic acetylcholine receptor antagonist that causes memory deficits similar to AD. I.c.v. injections of modified humanin (HNG) ameliorated the short-term memory deficits seen with scopolamine injection (Mamiya 2001). This modification replaced serine 14 for a glycine residue and was shown to increase the neuroprotective effects of humanin by 1000-fold (Hashimoto *et al.* 2001a,b). Using an alternative anticholinergic drug, 3-quinclidinyl benzilate, Krejcova *et al.* (2004) demonstrated that HNG injected i.p. could also help abrogate the memory impairments caused by this drug. This experiment also demonstrated that the positive effects of humanin did not require i.c.v. injections, which would help facilitate the use of humanin as a possible therapeutic drug.

Although the previous studies used a surrogate drug to examine the effects of humanin/HNG on AD-like memory deficits, the first *in vivo* studies that actually used A β also demonstrated that HNG could help prevent the effects of A β toxicity on memory defects

(Tajima *et al.* 2005). Injecting A β 25–35 into mice (i.c.v.) causes memory deficits within 3 weeks, which was ameliorated by HNG treatment (Tajima *et al.* 2005). Supporting the memory improvements seen with HNG administration, *ex vivo* experiments using hippocampal brain slices have also shown that HNG can prevent the inhibition of long-term potentiation caused by A β 25–35 applied an hour before recording (Zhang *et al.* 2009). Examination of acetylcholine-producing cells in the brain by staining for choline acetyltransferase, an enzyme used in the synthesis of acetylcholine, showed that A β 25–35 decreased the number of immunoreactive cells and that HNG was able to ameliorate this decrease (Tajima *et al.* 2005). Further examination of the *in vivo* effects of humanin/HNG on A β toxicity showed that HNG was able to reduce the inflammatory response caused by A β 25–35 as demonstrated by a decrease in reactive astrocytes, microglial activation, TNF α levels, and IL6 levels (Miao *et al.* 2008).

More recent studies have administered HNG to different transgenic models of AD. HNG administered intranasally to 1-year-old mice for 3 months reduced accumulation of A β levels in the brain and reduced the memory deficits in the triple transgenic AD model (APP_{SWE}, tau_{p310L}, PS-1_{M146V}) (Niikura *et al.* 2011). In mice with preexisting amyloid plaques, HNG administered to double transgenic mice (APP_{SWE}, PS-1_{DE9}) reduced the A β accumulation and improved the cognitive deficits compared with controls (Zhang *et al.* 2012). Thus, it seems that humanin may have therapeutic and/or prophylactic promise for treating AD.

Other neurodegenerative diseases

Although AD is the number one cause of dementia in adults, many other neurodegenerative diseases exist. These diseases cause neuronal stress by varying mechanisms but lead to the eventual death of neuronal cells. Because of the anti-apoptotic effects of humanin, several laboratories have examined the use of humanin to prevent neurodegeneration with mixed results.

Polyglutamine toxicity is the underlying cause of a number of different diseases such as spinocerebellar ataxia and Huntington's disease, where mitochondria dysfunction has been noted (reviewed in Takahashi *et al.* (2010)). The potential use of humanin to prevent polyglutamine toxicity has been investigated with divergent results. In one case, humanin had no protective ability, while in another, humanin partially protected the neuronal cell line (F11 and PC12 cells respectively) (Hashimoto *et al.*

2001b, Kariya *et al.* 2005). The many differences between the two studies could explain the equivocal results, but more studies of humanin as a possible treatment for polyglutamine diseases are certainly required.

Prions also cause neuronal stress and cell death and the possibility of humanin helping against this type of stress has also been examined. An initial study demonstrated no protective effect of humanin when using prion peptide 106–126 to induce cell stress and apoptosis (Hashimoto *et al.* 2001a). A later study confirmed this fact but found that humanin did have a protective effect against prion protein 118–135 (Sponne *et al.* 2004). Further studies will be required to determine the potential use of humanin in prion diseases.

Oxidative stress

Oxidative stress has been implicated in a multitude of diseases, and because of the success of humanin treatment against A β stress, several labs have tested humanin's ability to protect against oxidative stress. Treatment of rat retinal cultures with humanin has shown that it can protect cells from oxidative stress induced by hydrogen peroxide (Yang *et al.* 2008). The same study also demonstrated that hydrogen peroxide treatment decreases the endogenous protein levels of the rat homolog of humanin, Rattin (Yang *et al.* 2008). Further examination of humanin's ability to protect against oxidative stress focused on humanin's potential to help against atherosclerosis. In human carotid atherosclerotic plaques, humanin was found to be twice as high in symptomatic patients compared with asymptomatic, suggesting a role for humanin in atherosclerosis (Zacharias *et al.* 2012). An *in vitro* study found that humanin was able to protect endothelial cells from oxidative stress induced by oxidized LDL (Bachar *et al.* 2010). An *in vivo* follow-up on that observation demonstrated that HNG was able to improve cardiovascular function in *Apoe*-deficient mice placed on a high-cholesterol diet (Oh *et al.* 2011). Combining these data suggests that the increased levels of humanin found in atherosclerotic plaques are a protective response to stress and that humanin may be able to protect against oxidative stress.

Ischemia and reperfusion

Ischemic and reperfusion stress contributes to a large number of different physiological injuries such as stroke, myocardial infarction, and sickle cell disease, where it causes a number of different stresses such as oxidative

stress, hypoxia, and nutrient deprivation. Accumulating evidence suggests that humanin may be beneficial in attenuating the stresses caused by ischemia and reperfusion. One recent study examined humanin's ability to protect against the cellular effects of the hypoxia mimetic cobalt chloride (CoCl₂) (Men *et al.* 2012). Examining transformed rat ganglion cells, Men *et al.* (2012) found that humanin could help reduce the number of apoptotic cells caused by this treatment. A lack of growth factors and nutrients is another result of ischemia and serum deprivation increases apoptosis (Kariya *et al.* 2002). Kariya *et al.* (2002, 2003) as well as Guo *et al.* (2003) demonstrated that humanin can protect against serum starvation stress as well. Interestingly, they found that humanin increased metabolic activity (measured by MTS assay) and ATP levels of serum-starved cells above that of cells in complete media (Kariya *et al.* 2003).

In vivo studies have also demonstrated the benefits of humanin treatment. In a stroke model, mice underwent middle cerebral artery occlusion for 75 min. Mice that were either pretreated or posttreated up to 4 h after the occlusion with HNG i.c.v. had up to a 50% reduction in infarct volume size (Xu *et al.* 2006). Looking at a more clinically applicable method of humanin administration, Xu *et al.* (2006) also showed that an i.p. injection 2 h following the occlusion could also reduce the infarct size to a similar extent as the i.c.v. injection at 2 h. Similar to the stroke study, humanin can also ameliorate the effects of left coronary occlusion, and both pretreatment with humanin or treatment immediately following the occlusion can confer benefits (Muzumdar *et al.* 2010). In this study, both infarct size and loss of cardiac function were reduced with humanin treatment (Muzumdar *et al.* 2010). Thus, growing evidence supports the use of humanin as a cytoprotective agent against ischemia and reperfusion stress.

Humanin and its mechanism of action

Although all studies show a protective effect of humanin, the mechanisms by which humanin acts is still unclear. Ying *et al.* (2004) found that both A β 42 and humanin activate the same receptor, the G protein-coupled formylpeptide-like-1 receptor (FPRL1). Both acted as chemoattractants to monocytes and activated microglia, but humanin prevented the intracellular accumulation of A β 42 in these cells (Ying *et al.* 2004). Independently confirming the previous finding, Harada *et al.* (2004) screened for ligands for orphan G-coupled receptors and found that humanin directly interacts with the FPRL1 and

FPRL2 receptors in Chinese hamster ovary (CHO) cells transfected with the corresponding receptor. On the other hand, using a different type of cell, F11 hybrid neuronal cells, Hashimoto *et al.* (2005) found that the A β protection of humanin did not require FPRL1 but did require STAT signaling. Following on that observation, Hashimoto *et al.* (2009) also found that humanin requires a tripartite cytokine-like receptor complex comprising the CNTF receptor, WSX1, and gp130 (IL6ST) to protect neurons from A β . To reconcile these differences, it may be that different cell types employ different receptors to respond to humanin as both FPRL1 and FPRL2 are mainly found on phagocytic cells and not on neurons (reviewed in Le *et al.* (2002)). Humanin has also been shown to directly interact with A β and change the morphology of the A β aggregates and this could also be another method by which humanin protects against A β (Zou *et al.* 2003, Maftai *et al.* 2012). Additionally, HNG administration alone has been shown to increase insulin/IGF signaling in primary mouse neurons as assessed by AKT-1 phosphorylation, although studies in mouse heart has not found the same increase in phosphorylation and instead find an increase in phospho-AMPK (Xu *et al.* 2008, Muzumdar *et al.* 2010). An *in vivo* experiment has shown that HNG is a potent insulin sensitizer due to its ability to stimulate STAT3 in the hypothalamus (Muzumdar *et al.* 2009).

While the above pathways demonstrate an extracellular mechanism for humanin action, an intracellular, anti-apoptotic role has also been elucidated. Over several papers, the Reed laboratory has demonstrated that humanin interacts with many proteins in the apoptosis cascade (Guo *et al.* 2003, Luciano *et al.* 2005, Zhai *et al.* 2005). Humanin's anti-apoptotic function has been shown to be due to its interaction with the pro-apoptotic proteins BAX, tBID, and BimEL (Fig. 2). Humanin interacts with each of these proteins and inhibits further downstream signaling. The Reed laboratory further demonstrated that exogenous application of humanin did not prevent apoptosis in these studies and concluded that humanin was likely acting through an intracellular mechanism (Luciano *et al.* 2005, Zhai *et al.* 2005). These data seem to contradict data that had previously shown that the anti-apoptotic effects of humanin required extracellular secretion, although the papers used different cell types (Yamagishi *et al.* 2003).

Three other studies have found additional humanin interacting proteins. Using a yeast-2-hybrid system, TRIM11 was discovered (Niikura *et al.* 2003). TRIM11 likely ubiquitinates humanin for degradation and adds another level of regulation to humanin expression

Figure 2

Function of specific amino acid residues within humanin. Mutational analysis of humanin has allowed for the importance of each amino acid to be elucidated. Important amino acids are marked with their function as noted in the key.

(Niikura *et al.* 2003). In a different yeast-2-hybrid screen, a secreted antagonist of humanin was also found (Rossini *et al.* 2011). V-set and transmembrane domain containing 2 like (VSTM2L) was discovered to be a secreted protein that inhibits the positive effects of humanin action on neuronal apoptosis and this may be an additional way that the body regulates humanin action (Rossini *et al.* 2011). Lastly, humanin was found to interact with α -actinin-4, suggesting that humanin may have an unknown role in kidney function (Kigawa *et al.* 2004). Although there have been many studies examining the mechanism of humanin function, seemingly contradictory results necessitate further investigation.

Few studies have examined the mechanism or conditions by which humanin expression is increased. Humanin levels are known to be increased in endothelial cells of *ApoE*-deficient mice on a high-cholesterol diet, but the primary cause for this is unknown (Oh *et al.* 2011). Conversely, humanin levels are known to decrease with age in both rats and humans, and although the primary cause of this increase is unknown as well, it suggests that humanin may play a role in the aging process (Muzumdar *et al.* 2009, Bachar *et al.* 2010). More studies are needed to identify possible regulators of humanin transcription and translation.

Structure and function

The exact length of endogenous humanin is still unknown due to the differences in the mitochondrial and cytoplasmic translational machinery. If humanin is translated within the mitochondria, the peptide will be 21 amino acids; on the other hand, if it is translated in the cytoplasm, then the result is a 24 amino acid peptide.

Figure 3

Humanin's mechanism of action and interacting partners. Humanin has shown to have both intracellular and extracellular mechanisms of action. Many of its potential effects are mediated by the FPRL receptor as well as the trimeric WSX1/CNTF/gp130 receptor. Intracellularly, it interacts with many different proteins acting as a pro-survival signal.

Both peptides have been shown to have biological activity (Guo *et al.* 2003). Because humanin is a relatively short peptide, exhaustive mutational analysis of the importance of each amino acid has been possible (results summarized in Fig. 3). P3, S7, C8, L9, L12, T13, and S14 are important for humanin's protective effects against A β toxicity (Hashimoto *et al.* 2001b, Yamagishi *et al.* 2003). These residues may be important for the interaction with A β as the region between the 5th and 15th amino acid has been shown to interact with A β by affinity mass spectrometry (Maftai *et al.* 2012). Conversely, a mutation substituting glycine for S14 increases the neuroprotective ability of humanin by 1000 times (Hashimoto *et al.* 2001b). In addition to being important for the neuroprotective effects of humanin, C8 has also been shown to be important for humanin to bind BAX and tBID and also required to protect against prion proteins (Guo *et al.* 2003, Sponne *et al.* 2004, Zhai *et al.* 2005). Some laboratories have shown that humanin secretion is required for it to have its cytoprotective effects, and the amino acid residues

L9, L10, L11, P19, and V20 are important for secretion (Yamagishi *et al.* 2003). In terms of its IGFBP3 binding ability, both F6 and K21 are important for this interaction (Ikonen *et al.* 2003). Humanin is secreted from cells and can be found in the media of transfected cells (Yamagishi *et al.* 2003). Although there is no signal peptide to direct the peptide for secretion, humanin itself can act as a signal peptide, and when fused to larger proteins, it can cause the protein to be secreted into the media (Yamagishi *et al.* 2003).

Humanin-like proteins

In addition to the question of where humanin is translated, another major question that has not yet been fully resolved is whether humanin is translated from a mitochondrially encoded transcript, or if it is a product of a nuclear gene, or both. DNA from the mitochondria is often found dispersed within the nuclear genome, and, in fact, Ramos *et al.* (2011) found 755 insertion events in the human genome. Of these insertions, ~75% of them are <500 bp in length and 60% had <80% identity to the mitochondrial sequences (Ramos *et al.* 2011). With respect to humanin, 13 humanin-like ORFs were found within the nuclear genome, although only ten of them were found to be expressed in tissue samples (Bodzioch *et al.* 2009). Of these ten ORFs, only one of these codes for a possibly identical protein depending on the polymorphism used (Bodzioch *et al.* 2009). The rest of the ORFs code for peptides that have amino acid changes in critical residues

HN	MAPRGFSCLL	LLTSEIDLVP	KRRA
1	MAPRGFSCLL	LSTSEIDLVP	KRRT
2	MAPRGFSCLL	LSTSEIDLVP	KRLSSVF
3	MATRRFSCLL	LSTSEIDLVP	KRRI
4	MATQGFSCLL	LSVSEIDLSP	KROYKQIR
5	MATPGFSCLL	LSTSEIDLPM	KRRV
6	MTPRGFSCLL	LPTSETDLPV	KRRT
7	MATGGFSCLL	LLIREIDLSP	KROI
8*	MAPRGFSCLL	LSTSEIDLVP	KRRA
9	MARRGFSCLL	LSTTATDLPV	KRRT
10	MTTRGFSCLL	LLIREIDLSP	KRRI

Figure 4

Alignment of mitochondrial humanin with possible nuclear homologs. Ten different nuclear-encoded humanin (HN) homologs have been reported with all but one encoding for slightly different proteins than humanin.

*Humanin-like protein 8 has a reported polymorphism that would cause it to be identical to mitochondrially encoded humanin.

that may change their function, although this still needs to be carefully tested (see Fig. 4). Thus, the question of the origin of humanin still needs to be clarified.

Future studies

As mentioned earlier, one of the major questions in the field is whether endogenous humanin is derived from the mitochondria, and thus constitutes a novel category of peptides, or if humanin is nuclear in origin. Although the mitochondrial sequence of humanin has been identified independently several times as a cytoprotective peptide, its sequence similarity to other nuclear humanin-like peptides creates ambiguity as to whether it is endogenously expressed or whether its amino acid sequence is merely similar to an endogenously expressed peptide. Additionally, whether humanin is translated in the cytoplasm or mitochondria is also up for debate as both translations have bioactivity (Guo *et al.* 2003). Examination of mitochondrial RNA has found a high number of poly-adenylation sites within mitochondrial genes, and poly-adenylated mitochondrial sequences have been found in the cytoplasm, providing evidence that cytoplasmic translation may occur (Nakamichi *et al.* 1998, Mercer *et al.* 2011). Even if humanin is nuclear in origin, it can still be considered a MDP as the original nucleotide sequence has been transferred from the mitochondrial genome.

From a more practical point of view, it is clear that humanin is a cytoprotective peptide for a number of different stresses. Whether it is A β toxicity, serum starvation, hypoxia, oxidative stress, or any of the other stresses previously mentioned, both *in vitro* and *in vivo* studies have shown that administration of humanin has a beneficial effect.

The discovery of humanin as a novel, stress-responsive, and cytoprotective peptide derived from a mitochondrial ORF has opened up a new field of research. For decades, the number of genes within the mitochondria has been invariant, but this discovery forces the field to reexamine this assumption. Depending on how one analyzes the genome, there can be anywhere from 100 to over 500 ORFs. The majority are likely non-transcribed ORFs, but even if only a small percentage are real MDPs, there will be many novel discoveries. Because of the mitochondria's central role in stress resistance, these peptides are also likely to play a role in cellular stress resistance and may lead to new pharmacological agents and targets.

Declaration of interest

Dr P C is a consultant and stockholder of CohBar, Inc.

Funding

This research was supported by a Glenn Foundation Award and National Institute of Health Grants to P C (1R01AG 034430, 1R01GM 090311, 1R01ES 020812, and 1R21DK 089447) and an Ellison Medical Foundation/AFAR fellowship grant to K Y.

References

- Anderson S, Bankier AT, Barrell BG, de Bruijn MH, Coulson AR, Drouin J, Eperon IC, Nierlich DP, Roe BA, Sanger F *et al.* 1981 Sequence and organization of the human mitochondrial genome. *Nature* **290** 457–465. (doi:10.1038/290457a0)
- Bachar AR, Scheffer L, Schroeder AS, Nakamura HK, Cobb LJ, Oh YK, Lerman LO, Pagano RE, Cohen P & Lerman A 2010 Humanin is expressed in human vascular walls and has a cytoprotective effect against oxidized LDL-induced oxidative stress. *Cardiovascular Research* **88** 360–366. (doi:10.1093/cvr/cvq191)
- Balaban R, Nemoto S & Finkel T 2005 Mitochondria, oxidants, and aging. *Cell* **120** 483–495. (doi:10.1016/j.cell.2005.02.001)
- Bodzioch M, Lapicka-Bodzioch K, Zapala B, Kamysz W, Kiec-Wilk B & Dembinska-Kiec A 2009 Evidence for potential functionality of nuclearly-encoded humanin isoforms. *Genomics* **94** 247–256. (doi:10.1016/j.ygeno.2009.05.006)
- Drago I, Pizzo P & Pozzan T 2011 After half a century mitochondrial calcium in- and efflux machineries reveal themselves. *EMBO Journal* **30** 4119–4125. (doi:10.1038/emboj.2011.337)
- Durieux J, Wolff S & Dillin A 2011 The cell-non-autonomous nature of electron transport chain-mediated longevity. *Cell* **144** 79–91. (doi:10.1016/j.cell.2010.12.016)
- Guo B, Zhai D, Cabezas E, Welsh K, Nouraini S, Satterthwait AC & Reed JC 2003 Humanin peptide suppresses apoptosis by interfering with Bax activation. *Nature* **423** 456–461. (doi:10.1038/nature01627)
- Harada M, Habata Y, Hosoya M, Nishi K, Fujii R, Kobayashi M & Hinuma S 2004 N-formylated humanin activates both formyl peptide receptor-like 1 and 2. *Biochemical and Biophysical Research Communications* **324** 255–261. (doi:10.1016/j.bbrc.2004.09.046)
- Hashimoto Y, Niikura T, Ito Y, Sudo H, Hata M, Arakawa E, Abe Y, Kita Y & Nishimoto I 2001a Detailed characterization of neuroprotection by a rescue factor humanin against various Alzheimer's disease-relevant insults. *Journal of Neuroscience* **21** 9235–9245.
- Hashimoto Y, Niikura T, Tajima H, Yasukawa T, Sudo H, Ito Y, Kita Y, Kawasumi M, Kouyama K, Doyu M *et al.* 2001b A rescue factor abolishing neuronal cell death by a wide spectrum of familial Alzheimer's disease genes and A β . *PNAS* **98** 6336–6341. (doi:10.1073/pnas.101133498)
- Hashimoto YY, Suzuki HH, Aiso S, Niikura T, Nishimoto I & Matsuoka MM 2005 Involvement of tyrosine kinases and STAT3 in humanin-mediated neuroprotection. *Life Sciences* **77** 3092–3104. (doi:10.1016/j.lfs.2005.03.031)
- Hashimoto Y, Kurita M & Aiso S 2009 Humanin inhibits neuronal cell death by interacting with a cytokine receptor complex or complexes involving CNTF receptor α /WSX-1/gp130. *Molecular Biology of the Cell* **20** 2864–2873. (doi:10.1091/mbc.E09-02-0168)
- Haynes CM, Yang Y, Blais SP, Neubert TA & Ron D 2010 The matrix peptide exporter HAF-1 signals a mitochondrial UPR by activating the transcription factor ZC376.7 in *C. elegans*. *Molecular Cell* **37** 529–540. (doi:10.1016/j.molcel.2010.01.015)
- Ikonen M, Liu B, Hashimoto Y, Ma L, Lee K-W, Niikura T, Nishimoto I & Cohen P 2003 Interaction between the Alzheimer's survival peptide humanin and insulin-like growth factor-binding protein 3 regulates cell survival and apoptosis. *PNAS* **100** 13042–13047. (doi:10.1073/pnas.2135111100)

- Kantari C & Walczak H 2011 Caspase-8 and bid: caught in the act between death receptors and mitochondria. *Biochimica et Biophysica Acta* **1813** 558–563. (doi:10.1016/j.bbamcr.2011.01.026)
- Kariya S, Takahashi N, Ooba N, Kawahara M, Nakayama H & Ueno S 2002 Humanin inhibits cell death of serum-deprived PC12h cells. *Neuroreport* **13** 903–907. (doi:10.1097/00001756-200205070-00034)
- Kariya S, Takahashi N & Hirano M 2003 Humanin improves impaired metabolic activity and prolongs survival of serum-deprived human lymphocytes. *Molecular and Cellular Biochemistry* **254** 83–89. (doi:10.1023/A:1027372519726)
- Kariya S, Hirano M, Nagai Y, Furiya Y, Fujikake N, Toda T & Ueno S 2005 Humanin attenuates apoptosis induced by DRPLA proteins with expanded polyglutamine stretches. *Journal of Molecular Neuroscience* **25** 165–169. (doi:10.1385/JMN:25:2:165)
- Kigawa A, Wakui H, Maki N, Okuyama S, Masai R, Ohtani H, Komatsuda A, Suzuki D, Toyoda M, Kobayashi R *et al.* 2004 Interaction of the spectrin-like repeats of α -actinin-4 with humanin peptide. *Clinical and Experimental Nephrology* **8** 331–338. (doi:10.1007/s10157-004-0322-y)
- Krejcová G, Patocka J & Slaninová J 2004 Effect of humanin analogues on experimentally induced impairment of spatial memory in rats. *Journal of Peptide Science* **10** 636–639. (doi:10.1002/psc.569)
- Le Y, Murphy PM & Wang JM 2002 Formyl-peptide receptors revisited. *Trends in Immunology* **23** 541–548. (doi:10.1016/S1471-4906(02)02316-5)
- Luciano F, Zhai D, Zhu X, Bailly-Maitre B, Ricci J-E, Satterthwait AC & Reed JC 2005 Cytoprotective peptide humanin binds and inhibits proapoptotic Bcl-2/Bax family protein BimEL. *Journal of Biological Chemistry* **280** 15825–15835. (doi:10.1074/jbc.M413062200)
- Maftai M, Tian X, Manea M, Exner TE, Schwanzer D, Arnimvon CAF & Przybylski M 2012 Interaction structure of the complex between neuroprotective factor humanin and Alzheimer's β -amyloid peptide revealed by affinity mass spectrometry and molecular modeling. *Journal of Peptide Science* **18** 373–382. (doi:10.1002/psc.2404)
- Mamiya T 2001 [Gly14]-humanin improved the learning and memory impairment induced by scopolamine *in vivo*. *British Journal of Pharmacology* **134** 1597–1599. (doi:10.1038/sj.bjp.0704429)
- Men J, Zhang X, Yang Y & Gao D 2012 An AD-related neuroprotector rescues transformed rat retinal ganglion cells from CoCl₂(2)-induced apoptosis. *Journal of Molecular Neuroscience* **47** 144–149. (doi:10.1007/s12031-011-9701-5)
- Mercer TR, Neph S, Dinger ME, Crawford J, Smith MA, Shearwood A-MJ, Haugen E, Bracken CP, Rackham O, Stamatoyannopoulos JA *et al.* 2011 The human mitochondrial transcriptome. *Cell* **146** 645–658. (doi:10.1016/j.cell.2011.06.051)
- Miao J, Zhang W & Li Z 2008 S14G-Humanin ameliorates A β 25–35-induced behavioral deficits by reducing neuroinflammatory responses and apoptosis in mice. *Neuropeptides* **42** 557–567. (doi:10.1016/j.npep.2008.08.004)
- Muzumdar R, Huffman D, Atzmon G, Buettner C, Cobb L, Fishman S, Budagov T, Cui L, Einstein F, Poduval A *et al.* 2009 Humanin: a novel central regulator of peripheral insulin action. *PLoS ONE* **4** e6334. (doi:10.1371/journal.pone.0006334)
- Muzumdar RH, Huffman DM, Calvert JW, Jha S, Weinberg Y, Cui L, Nemkal A, Atzmon G, Klein L, Gundewar S *et al.* 2010 Acute humanin therapy attenuates myocardial ischemia and reperfusion injury in mice. *Arteriosclerosis, Thrombosis, and Vascular Biology* **30** 1940–1948. (doi:10.1161/ATVBAHA.110.205997)
- Nakamichi N, Rhoads DD, Hayashi JI, Kagawa Y & Matsumura T 1998 Detection, localization, and sequence analyses of mitochondrial regulatory region RNAs in several mammalian species. *Journal of Biochemistry* **123** 392–398. (doi:10.1093/oxfordjournals.jbchem.a021950)
- Niikura T, Hashimoto Y, Tajima H, Ishizaka M, Yamagishi Y, Kawasumi M, Nawa M, Terashita K, Aiso S & Nishimoto I 2003 A tripartite motif protein TRIM11 binds and destabilizes humanin, a neuroprotective peptide against Alzheimer's disease-relevant insults. *European Journal of Neuroscience* **17** 1150–1158. (doi:10.1046/j.1460-9568.2003.02553.x)
- Niikura T, Sidahmed E, Hirata-Fukae C, Aisen PS & Matsuoka Y 2011 A humanin derivative reduces amyloid β accumulation and ameliorates memory deficit in triple transgenic mice. *PLoS ONE* **6** e16259. (doi:10.1371/journal.pone.0016259)
- Oh YK, Bachar AR, Zacharias DG, Kim SG, Wan J, Cobb LJ, Lerman LO, Cohen P & Lerman A 2011 Humanin preserves endothelial function and prevents atherosclerotic plaque progression in hypercholesterolemic ApoE deficient mice. *Atherosclerosis* **219** 65–73. (doi:10.1016/j.atherosclerosis.2011.06.038)
- Ramos A, Barben A, Mateiu L, del Mar González M, Mairal Q, Lima M, Montiel R, Aluja MP & Santos C 2011 Nuclear insertions of mitochondrial origin: database updating and usefulness in cancer studies. *Mitochondrion* **11** 946–953. (doi:10.1016/j.mito.2011.08.009)
- Ristow M & Zarse K 2010 How increased oxidative stress promotes longevity and metabolic health: the concept of mitochondrial hormesis (mitohormesis). *Experimental Gerontology* **45** 410–418. (doi:10.1016/j.exger.2010.03.014)
- Ristow M, Zarse K, Oberbach A, Klitting N, Birringer M, Kiehnopf M, Stumvoll M, Kahn C & Blüher M 2009 Antioxidants prevent health-promoting effects of physical exercise in humans. *PNAS* **106** 8665–8670. (doi:10.1073/pnas.0903485106)
- Rossini L, Hashimoto Y, Suzuki H, Kurita M, Gianfriddo M, Scali C, Roncarati R, Franceschini D, Pollio G, Traballini L *et al.* 2011 VSTM2L is a novel secreted antagonist of the neuroprotective peptide humanin. *FASEB Journal* **25** 1983–2000. (doi:10.1096/fj.10-163535)
- Salway JG 1994 Metabolism at a Glance. edn 5, pp 10, Malden, MA: Blackwell Publishing Ltd.
- Sponne I, Fife A, Koziel V, Kriem B, Oster T & Pillot T 2004 Humanin rescues cortical neurons from prion-peptide-induced apoptosis. *Molecular and Cellular Neurosciences* **25** 95–102. (doi:10.1016/j.mcn.2003.09.017)
- Tajima H, Kawasumi M, Chiba T, Yamada M, Yamashita K, Nawa M, Kita Y, Kouyama K, Aiso S, Matsuoka M *et al.* 2005 A humanin derivative, S14G-HN, prevents amyloid- β -induced memory impairment in mice. *Journal of Neuroscience Research* **79** 714–723. (doi:10.1002/jnr.20391)
- Takahashi T, Katada S & Onodera O 2010 Polyglutamine diseases: where does toxicity come from? what is toxicity? where are we going? *Journal of Molecular Cell Biology* **2** 180–191. (doi:10.1093/jmcb/mjq005)
- Wang X 2001 The expanding role of mitochondria in apoptosis. *Genes and Development* **15** 2922–2933.
- Xu X, Chua CC, Gao J, Hamdy RC & Chua BHL 2006 Humanin is a novel neuroprotective agent against stroke. *Stroke; a Journal of Cerebral Circulation* **37** 2613–2619. (doi:10.1161/01.STR.0000242772.94277.1f)
- Xu X, Chua C, Gao J, Chua K & Wang H 2008 Neuroprotective effect of humanin on cerebral ischemia/reperfusion injury is mediated by a PI3K/Akt pathway. *Brain Research* **1227** 12–18. (doi:10.1016/j.brainres.2008.06.018)
- Yamagishi Y, Hashimoto Y, Niikura T & Nishimoto I 2003 Identification of essential amino acids in humanin, a neuroprotective factor against Alzheimer's disease-relevant insults. *Peptides* **24** 585–595. (doi:10.1016/S0196-9781(03)00106-2)
- Yang Z, Zhang Q, Ge J & Tan Z 2008 Protective effects of tetramethylpyrazine on rat retinal cell cultures. *Neurochemistry International* **52** 1176–1187. (doi:10.1016/j.neuint.2007.12.008)
- Ying G, Iribarren P, Zhou Y, Gong W, Zhang N, Yu Z-X, Le Y, Cui Y & Wang JM 2004 Humanin, a newly identified neuroprotective factor, uses the G protein-coupled formylpeptide receptor-like-1 as a functional receptor. *Journal of Immunology* **172** 7078–7085.
- Zacharias DG, Kim SG, Massat AE, Bachar AR, Oh YK, Herrmann J, Rodriguez-Porcel M, Cohen P, Lerman LO & Lerman A 2012 Humanin, a cytoprotective Peptide, is expressed in carotid atherosclerotic plaques in humans. *PLoS ONE* **7** e31065. (doi:10.1371/journal.pone.0031065)

- Zhai D, Luciano F, Zhu X, Guo B, Satterthwait AC & Reed JC 2005 Humanin binds and nullifies Bid activity by blocking its activation of Bax and Bak. *Journal of Biological Chemistry* **280** 15815–15824. (doi:10.1074/jbc.M411902200)
- Zhang W, Miao J, Hao J, Li Z, Xu J, Liu R, Cao F, Wang R, Chen J & Li Z 2009 Protective effect of S14G-humanin against β -amyloid induced LTP inhibition in mouse hippocampal slices. *Peptides* **30** 1197–1202. (doi:10.1016/j.peptides.2009.02.017)

- Zhang W, Zhang W, Li Z, Hao J, Zhang Z, Liu L, Mao N, Miao J & Zhang L 2012 S14G-humanin improves cognitive deficits and reduces amyloid pathology in the middle-aged APPswe/PS1dE9 mice. *Pharmacology, Biochemistry, and Behavior* **100** 361–369. (doi:10.1016/j.pbb.2011.09.012)
- Zou P, Ding Y, Sha Y, Hu B & Nie S 2003 Humanin peptides block calcium influx of rat hippocampal neurons by altering fibrogenesis of A β (1–40). *Peptides* **24** 679–685. (doi:10.1016/S0196-9781(03)00131-1)

Received in final form 27 November 2012

Accepted 10 December 2012

Accepted Preprint published online 10 December 2012