Nuclear receptors and chromatin: an inducible couple

Raffaella Maria Gadaleta and Luca Magnani

Division of Cancer, Imperial Centre for Translational and Experimental Medicine, Imperial College London, London, W12 0NN, UK

Correspondence
should be addressed
to L. Magnani
Email
l.magnani@imperial.ac.uk

Abstract

The nuclear receptor (NR) family comprises 48 transcription factors (TFs) with essential and diverse roles in development, metabolism and disease. Differently from other TFs, NRs engage with well-defined DNA-regulatory elements, mostly after ligand-induced structural changes. However, NR binding is not stochastic, and only a fraction of the cognate regulatory elements within the genome actively engage with NRs. In this review, we summarize recent advances in the understanding of the interactions between NRs and DNA. We discuss how chromatin accessibility and epigenetic modifications contribute to the recruitment and transactivation of NRs. Lastly, we present novel evidence of the interplay between non-coding RNA and NRs in the mediation of the assembly of the transcriptional machinery.

Key Words

- ▶ nuclear receptors
- ▶ chromatin
- ▶ epigenetics
- ▶ oestrogen receptors
- ▶ gene expression

*Journal of Molecular
Endocrinology*
(2014) 52, R137–R149

Introduction/overview

Nuclear receptors (NRs) are evolutionarily related DNA-binding transcription factors (TFs) (Gronemeyer *et al.* 2004, Chambon 2005, Evans 2005). They regulate many aspects of mammalian physiology, including metabolism, development and reproduction (Mangelsdorf *et al.* 1995, Willy *et al.* 1995, Chawla *et al.* 2001), priming the activation of multiple genes to achieve accurate, synchronized and coherent functional responses. Most NRs regulate gene transcription in response to a ligand activation process, being activated by several hormones, vitamins, lipophilic metabolites and dietary lipids. NRs are also able to recruit a variety of classical TFs, co-regulators and proteins of the basal transcriptional machinery to target gene promoters and coordinate the whole process of gene transcription. The NR superfamily has a high potential for drug targeting, because the lipophilic ligands of NRs can easily pass through biological membranes. Consequently, NRs are the object of intensive translational research, which has resulted in their

establishment as drug targets for several pathological conditions. Currently, 13% of all FDA-approved drugs target NRs (Overington *et al.* 2006). These include thiazolidinediones targeting the peroxisome proliferator-activated receptor (PPAR) in type II diabetes patients, steroids targeting the glucocorticoid receptor (GR) in inflammatory diseases, tamoxifen targeting the oestrogen receptor- α (ER α) in breast cancer and antiandrogens targeting the androgen receptor (AR) in prostate cancer.

In 1985, the GR was the first NR to be cloned (Hollenberg *et al.* 1985), followed by the ER and thyroid hormone receptor (THR; Walter *et al.* 1985, Greene *et al.* 1986, Sap *et al.* 1986). To date, the NR superfamily comprises 48 members in humans. NRs mediate gene transcription via distinct mechanisms, including both classical genomic transactivation and transrepression activities. Furthermore, NRs can also modulate different pathways in a non-genomic fashion, and their activities can be influenced by post-translational modifications

(e.g. acetylation, phosphorylation, sumoylation and ubiquitination), making them targets of other cellular pathways. In this review, we mostly focus on the genomic activities of NRs.

Despite their broad range of activities, NRs share a common structural organization. They consist of an N-terminal regulatory domain, containing the ligand-independent activation function 1 (AF1; [Wärnmark et al. 2003](#)) that usually has a very weak transcriptional activity (with the notable exception of the AR), but is capable of synergizing with another activation domain, activation function 2 (AF2), present in the ligand-binding domain (LBD; see below). The most conserved domain is the DNA-binding domain (DBD), which contains the P-box, responsible for direct DNA interaction and DNA-binding specificity. It consists of two zinc fingers, which recognize specific NR-responsive elements (NRREs) in the DNA-regulatory regions of their target genes. NRs can bind to NRREs located in the proximity of target gene promoters and to more distal elements such as enhancer regions, as monomers, homodimers (e.g. GR, ER and AR) or heterodimers with the retinoid X receptor (RXR) (e.g. farnesoid X receptor (FXR), liver X receptor (LXR) and others). NRREs consist of two hexameric core half-site motifs, separated by a variable number of nucleotides. The bipartite elements can form direct, inverted or everted repeats ([Mangelsdorf & Evans 1995](#)). The DBD also contains a dimerization interface ([Zechel et al. 1994](#), [Germain et al. 2006](#), [Bain et al. 2007](#)), which can also be the site of post-translational modifications, resulting in different effects for different NRs. In some NRs, DBD phosphorylation results in decreased DNA binding (e.g. THR, [Tzagarakis-Foster & Privalsky \(1998\)](#) and ER, [Chen et al. \(1999\)](#)); in others, it increases DNA binding and co-activator recruitment (e.g. FXR, [Gineste et al. 2008](#)). The DBD and the C-terminal LBD are connected by a flexible domain called hinge region, which houses a nuclear localization sequence ([Giguère et al. 1986](#)). The LBD presents with a structure forming a hydrophobic pocket for ligand recognition and lodging. It also contains the AF2 domain, the action of which is dependent on the presence of bound ligand ([Wärnmark et al. 2003](#)), as well as motifs responsible for dimerization and co-regulator-binding regions.

For the majority of NRs, ligand binding is the critical event shifting their function from an inactive to an active state by inducing a conformational change in the LBD ([Bourguet et al. 2000](#)). The active conformation allows the second step of NR activation: recruitment of co-regulatory complexes. In recent years, it has become clear that many of these co-factors function as chromatin-remodelling

factors and can catalyse histone modifications ([Kouzarides 2007](#), [Suganuma & Workman 2008](#)). Small modifications of ligand structure seem to affect the co-activator-binding interface, providing a molecular basis for NR–ligand binding specificity and potency ([Wu et al. 2003](#)). The LBD is also a target for post-translational modifications, which govern a variety of cellular functions, including NR activity, DNA-binding affinity, ligand sensitivity, receptor stability and subcellular distribution. Therefore, NR post-translational modifications can ultimately result in either increased or decreased target gene expression ([Balasubramanian et al. 2012](#)) or in transrepression of other pathways (such as sumoylation of PPAR γ ([Pascual et al. 2005](#)) and FXR, leading to NF- κ B transrepression ([Vavassori et al. 2009](#))).

NR classification

NRs can be classified in several ways. Based on ligand-binding properties, they can be divided into three classes: hormone NRs, metabolic NRs and orphan NRs ([Table 1](#)). Hormone NRs are often located in the cytosol. Once bound to their ligands, they translocate to the nucleus, usually as homodimers, and bind to their cognate NRREs. The members of this subfamily include the AR, GR and progesterone receptor (PR) and ligands include hormonal lipids. Upon ligand binding, some hormone NRs bind to the DNA on accessible chromatin at regulatory sites in the

Table 1 Classification of some members of the nuclear receptor superfamily. NRs can be classified in several ways. Based on ligand-binding property, they can be divided into hormone, metabolic and orphan NRs. Hormone NRs generally bind to the DNA as homodimers. On the contrary, most of the metabolic NRs bind to DNA as heterodimers with their obligate partner RXR. Orphan NRs can work as either heterodimers or monomers. For a more complete classification, see [Nuclear Receptors Nomenclature Committee \(1999\)](#)

Hormone NRs ligands: hormones	Metabolic NRs ligands: dietary lipids	Orphan NRs ligands: unknown
ER α , β	RXR α , β , γ	COUP-TF1,II,III
GR	PPAR α , β , γ	NUR77, NURR1, NOR1
MR	LXR α , β	GCNF
AR	FXR α , β	LRH1
PR	PXR/SXR	Rev-ERB α , β
RAR α , β , γ	CAR	SHP
THR α , β		THR2,3,4
VDR		HNF4 α , β , γ
ECR		DAX1

target genes, such as ER and GR; others bind to chromatin and attract co-activator factors that have chromatin-remodelling properties, as in the case of ER and PR (Vicent *et al.* 2011). DNA binding and chromatin remodelling then enhance the recruitment and/or function of the general transcription machinery (Fondell *et al.* 1996, Lefterova *et al.* 2008).

Metabolic receptors are typically located in the nucleus, bound to the DNA mostly as heterodimers with the RXR. In the absence of ligands, they are often complexed on the chromatin with co-repressor proteins. In this instance, ligand binding to NRs causes the dissociation of co-repressor and recruitment of co-activator proteins. Additional proteins of the basal transcriptional machinery, including RNA polymerase II, are then recruited to the NR–DNA complex to start the transcription process. The members of this subfamily include the FXR, LXR, PPARs and others, while ligands for this subgroup encompass dietary lipids. The third subgroup of NRs is the so-called family of orphan receptors, for which regulatory ligands are still unknown or may not exist (true orphans) or for which candidates have only recently been identified (adopted orphans). Members include the chicken ovalbumin upstream transcription factors (COUP-TFs), reverse ERBA- α / β , liver receptor homologue-1 (LRH1) and others.

In this review, we summarize current knowledge regarding the bidirectional crosstalk between NRs and the chromatin landscape in the context of gene regulation.

Genomic features of NR binding

Similarly to most TFs, NRs modulate transcription by binding to well-defined DNA sequences (NRREs) interspersed in the genome (Kininis & Kraus 2008). NRs can also bind to DNA via tethering mechanisms; for example, ER α can ‘piggyback’ on AP1, MAF and RUNX1 TFs (Stender *et al.* 2010, Heldring *et al.* 2011). However, for the sake of clarity, we focus only on direct binding events. NRREs are extremely well conserved and share a significant homology. Concordantly, NR DBDs share extensive homologies (Kininis & Kraus 2008, Willis & Griffin 2011). Despite this, a recent NR binding survey has identified dramatically distinct topological differences in regulatory element selection (Kittler *et al.* 2013). Kittler *et al.* (2013) used a tag-based ChIP-seq approach to circumvent the noise created by different antibody efficiencies and characterized the binding profile of 24 NRs. Several NRs appear to have a strong preference for distal binding (ER α , RORC, AR and PR). Others are found

in proximity to promoter elements (testicular receptors TR2L, TR2S, TR4, LXR α and THR α (Kittler *et al.* 2013). Interestingly, only 32% of over 2×10^5 individual genomic loci have been found to recruit a single NR, suggesting that NRs exhibit extensive cooperative binding (Voss *et al.* 2011). For example, ten of the 24 NRs assayed are recruited to the *CCND1* promoter. These genomic regions with multiple NR-binding sites (HOT regions) exhibit a significant increase in features normally associated with active regulatory elements, including epigenetic modifications and increased chromatin accessibility (Kittler *et al.* 2013). HOT regions are reminiscent of super-enhancers described, for example, in embryonic stem cells (Whyte *et al.* 2013). In this case, several regions termed super-enhancers span significantly large stretches of DNA (median size 8 kb) and recruit large arrays of TFs (e.g. OCT4, SOX2, NANOG, KLF4 and ESRRB). Not surprisingly, breast cancer-relevant genes are enriched with HOT regions (e.g. *CCND1* and *MYC* genes; Kittler *et al.* 2013). These data suggest that combinatorial binding patterns may play an unexpected role in hormone-dependent cancer and other physiological processes regulated by NRs.

Chromatin accessibility predisposes regulatory elements to NR binding

NRs invariably bind to regulatory elements to control cell type-specific transcription in response to external stimuli. The search for putative regulatory elements, such as promoters and enhancers, has been simplified by the recent development of epigenomics (ENCODE Project Consortium *et al.* 2007, 2012). Cell type-specific regulatory elements can be classified using several epigenetic modifications, including chromatin accessibility (Ernst *et al.* 2011). With its nucleosome-rich structure, chromatin represents the first barrier for TF binding. In agreement with this, regulatory elements are commonly found in nucleosome-depleted pockets, a conformation thought to be essential for DNA motif recognition and binding (Thurman *et al.* 2012). Uniquely among other TFs, several NRs bind to DNA only after conformational changes induced by their ligands (Kininis & Kraus 2008). This raises the fundamental question about pre-induction and post-induction chromatin determinants of NR binding (Magnani & Lupien 2013) (Fig. 1).

Chromatin accessibility pre-induction

Most of our understanding of NR–chromatin interaction comes from dissecting ER α , GR and AR signalling. Initial

Figure 1

Schematic representation of nuclear receptors and chromatin accessibility. Nuclear receptors bind to responsive elements present in the DNA with different mechanisms. Chromatin accessibility and histone modifications contribute to NR binding via regulatory element bookmarking before ligands mediate activation (A). In some instances, NRs bind even in the

absence of ligands while in complex with other proteins (B). Upon ligand activation, several NRs can guide further modifications of the chromatin landscape by recruiting histone modifiers and chromatin-remodelling complexes (C).

studies considering a few loci of GR-bound chromatin found some sites to be constitutively accessible using DNase I hypersensitivity site (DHS) assays and others to be remodelled upon ligand stimulation (John *et al.* 2008). However, a follow-up genome-wide screen using DHS assays coupled with next-generation sequencing (DHS-seq) revealed that the majority (95%) of GR binding occurs at pre-induction accessible sequences. These data demonstrate that chromatin accessibility can pre-select regulatory elements for GR binding. Adult cells present unique accessibility profiles (Thurman *et al.* 2012), suggesting that cell type-specific effects of NR activation could be partly mediated by the chromatin topology (John *et al.* 2008).

Nucleosome occupancy plays a different role in the context of AR binding at enhancers. After stimulation with dihydrotestosterone (DHT), He *et al.* (2012) observed the destabilization of a central nucleosome while the flanking nucleosomes remained occupied. In agreement with this, the central nucleosome is built around a modified histone protein called H2A.Z, normally associated with fragile nucleosomes (Jin *et al.* 2009). Another study confirmed and expanded these observations and described a significant correlation between changes in DHS and AR-mediated transcription, suggesting that AR binding relieves nucleosome occupancy and potentially increases access for collaborating factors (Tewari *et al.* 2012). However, a later report suggested an intermediate model in which enhancers recruiting AR are in a stable equilibrium between

nucleosome occupancy and nucleosome-free state in uninduced cells. Androgen treatment then destabilizes this equilibrium in favour of the more accessible nucleosome-free state (Andreu-Vieyra *et al.* 2011).

The situation is less clear-cut for ER α . Using formaldehyde-assisted isolation of regulatory element (FAIRE) assays, a technique that allows enrichment of nucleosome-depleted regions (Giresi *et al.* 2007), Hurtado *et al.* (2010) reported that a substantial amount of ER α can bind to more compacted chromatin regions. More importantly, they found that a significant proportion of accessible sites are present before oestrogen induction and they can recruit ER α (Hurtado *et al.* 2010). Identical conclusions can be drawn from a study that profiled chromatin accessibility before and after oestradiol (E2) stimulation, again using FAIRE assays (Joseph *et al.* 2010). These data are in substantial agreement with a more recent work that profiled ER α chromatin accessibility using DHS assays (He *et al.* 2012). In addition, it was noted that ER α binding that leads to a substantial increase in chromatin accessibility clusters closer to oestrogen-responsive genes, suggesting that basal accessibility state can be further modified (discussed below, He *et al.* (2012)).

Chromatin accessibility post-induction

In the case of ligand-mediated DNA binding, nucleosome remodelling appears to continue even if the chromatin is

already in an accessible structure (He *et al.* 2012). Post-induction remodelling may have functional consequences for NR crosstalk. The assisted loading model predicts that binding of a NR may increase the chances of binding of a second NR via chromatin remodelling (Stratmann & Schibler 2011). ER α directly or indirectly recruits several chromatin-remodelling complexes on the chromatin after oestrogen stimulation (Belandia *et al.* 2002, Jeong *et al.* 2009). Similarly, GR is associated with SWI/SNF complexes, proteins involved in ATP-mediated nucleosome remodelling (reviewed elsewhere, King *et al.* (2012)). Accordingly, GR activation using dexamethasone can increase binding of E2-induced ER α at proximal chromatin locations (Voss *et al.* 2011, Miranda *et al.* 2013). Very similar observations were made with regard to the PR (Grøntved & Hager 2012). Furthermore, another recent report has identified local interactions between ER α and LRH1 (Lai *et al.* 2013). Depletion of LRH1 was found to cause decreased ER α binding, while LRH1 overexpression was found to facilitate ER α recruitment in the absence of ER α protein levels changes, again supporting the notion of assisted loading. These data are consistent with the transient nature of the interaction of NRs with the chromatin. Indeed, ER α and the molecular machinery that accompanies ER α binding can contact the chromatin at relatively predictable intervals in synchronized MCF7 cells (Métivier *et al.* 2003, 2008, Reid *et al.* 2003). Although speculative, it is conceivable that activated NRs could also influence binding of other TFs. Collectively, these data highlight the importance of chromatin accessibility in dictating NR binding and how NRs modulate post-induction remodelling processes.

Induction-independent

Ligand-mediated activation does not always lead to chromatin binding. For example, several NRs are constitutively bound to the chromatin even in the absence of ligands. While there is evidence for agonist-dependent changes in binding of some hormone NRs to chromatin (as described above), the influence of agonist binding on DNA occupancy for metabolic NRs is complex and not fully understood. ChIP-seq analysis of mouse liver chromatin for FXR-binding sites revealed that the majority are located in distal intergenic regions or introns and fewer sites are localized in more proximal promoter regions (Chong *et al.* 2010). The high distribution of binding sites within intergenic and intronic regions is consistent with similar reports for other steroid and metabolic NRs, including PPAR γ (Lefterova *et al.* 2008, Nielsen *et al.*

2008), ER α (Carroll *et al.* 2005) and AR (Bolton *et al.* 2007). Chong *et al.* (2010) found no statistically significant difference in genome-wide binding of FXR in the livers of a control group fed a normal chow diet vs a group fed chow supplemented with GW4064, the potent synthetic FXR agonist, suggesting that FXR binding is unaffected by ligand stimulation. These data are in substantial agreement with another genome-wide FXR binding study carried out in WT and obese mice, revealing a ligand-independent co-occupancy of FXR and RXR (Lee *et al.* 2012). The same pattern was observed in another genome-wide binding site study, in mouse liver and intestine, where it was shown that FXR was already bound to its own REs in the regulatory regions of FXR target genes (Thomas *et al.* 2010). Although further studies need to be carried out, and different ligands studied, these results support the concept that metabolic NRs are bound to the DNA regardless of the presence of the ligand at accessible regulatory elements.

However, it is important to underline that these data might also be influenced by the basal FXR activation by bile acids, the FXR endogenous ligands. Nevertheless, the binding intensity was found to be boosted upon treatment with GW4064, in agreement with other *in vitro* studies in a liver cell line, HepG2, where it was shown that FXR binds to its REs weakly in the absence of ligands, and this binding is increased upon ligand treatment (Rizzo *et al.* 2005, Fang *et al.* 2008). Once again, this effect might be due to a ligand-switch effect, and it would be interesting to dissect the binding patterns of FXR upon treatment with natural ligands. Noteworthy, as pointed out by Chong *et al.* (2010), although bioinformatics analysis revealed over 1.7 million sites in the mouse genome matching the canonical FXR-REs, a very small percentage (<0.1%) were shown to be occupied by FXR in their study. These results further support the idea that NRs are able to bind to their REs, although RE specificity is not sufficient. To achieve fully efficient NR-DNA binding, specific conditions are needed, including specific nucleosome positioning and epigenetic modifications able to alter chromatin structure and DNA accessibility.

Histone post-translational modifications and their role in NR recruitment

Epigenetic modifications have a well-characterized function during normal development where they bookmark regulatory elements to maintain heritable gene expression (Bernstein *et al.* 2006, Christophersen & Helin 2010, Ernst & Kellis 2010, Heinz *et al.* 2010, Ernst *et al.* 2011,

Zhu *et al.* 2013). Only recently have we started to understand the relationship between histone modifications and NR activity. Several studies have shown direct interactions between the activation of NRs and the remodelling of chromatin through histone modifications. Analogously to chromatin remodelling, histone modifications play different roles and can be divided into pre-induction and post-induction.

Pre-induction histone modifications

Regulatory elements are characterized by an epigenetic grammar where combinatorial patterns of epigenetic modifications are associated with active, poised or inactive promoters and enhancers (Ernst & Kellis 2010, Heinz *et al.* 2010, Ernst *et al.* 2011). For example, active enhancers are commonly marked by histone H3 lysine K4 mono- and/or dimethylation (H3K4me1/2) (Lupien *et al.* 2008, Ernst & Kellis 2010). On the other hand, nucleosomes near the transcription start sites of active promoters are enriched with histone H3 lysine K4 trimethylation (H3K4me3; Bernstein *et al.* 2006, Thurman *et al.* 2012). NRs, similar to many other TFs, interpret the chromatin to find their cognate binding sequences (Magnani & Lupien 2013). Therefore, preloaded epigenetic modifications can contribute to the reduction of the number of potential binding sites by 'flagging' active regulatory elements. For example, Lupien *et al.* (2008) found that binding of sex steroid NRs ER α and AR significantly correlated with H3K4me2 patterns in breast and prostate cancer cell lines respectively. As expected, breast and prostate cells are characterized by markedly different H3K4me2 histone mark layouts in agreement with cell type specificity of enhancer elements (Ernst *et al.* 2011, Zhu *et al.* 2013). The H3K4me2 as well as the H4K4me1 signal does not change in response to oestrogen stimulation in breast cancer cells (Lupien *et al.* 2008, Joseph *et al.* 2010, Magnani *et al.* 2011a), suggesting that histone methylation is independent of and precedes ER α binding. Similarly, in prostate cancer cells, H3K4me1/2 is found at active enhancers capable of recruiting AR before DHT stimulation in support of its role as a bookmarking modification (Wang *et al.* 2009). Limited functional studies assessing the effect of H3K4me removal on ER α signalling (Lupien *et al.* 2008, Magnani *et al.* 2011a) using overexpression of a histone lysine (K)-specific demethylase 1 (LSD1) indeed found a significant loss in ER α binding.

Interaction between epigenetic modification and NR binding is in some cases mediated by a group of proteins called pioneer factors (Magnani *et al.* 2011b, Zaret &

Carroll 2011, Magnani & Lupien 2013). Pioneer factors are a functionally defined family of proteins, including structurally unrelated TFs, capable of binding compacted chromatin (for extensive reviews, see Magnani *et al.* 2011b, Zaret & Carroll 2011). These factors can relieve chromatin compaction during lineage commitment, to maintain accessibility at cell type-specific enhancers (Cirillo & Zaret 1999). Pioneer factors can also act as molecular beacons for other TFs; for example, the fork-head family members FoxD3 and FoxA1 mark the liver-specific Alb1 enhancer starting from early embryonic development, preceding C/EBP, Ey and NFF1 binding and Alb1 expression (Zaret *et al.* 2008). Similarly, the homeobox pre-B-cell leukaemia transcription factor (PBX1) was found to precede MyoD recruitment during muscle development (Berkes *et al.* 2004). Accumulating evidence suggests that pioneer factors are preloaded on the chromatin at ER α -recruiting regulatory elements before ligand activation and contribute to increased chromatin accessibility (Jozwik & Carroll 2012). Depletion of pioneer factors including FOXA1 and PBX1 has a significant negative impact on ER α -chromatin interaction (Lupien *et al.* 2008, Magnani *et al.* 2011a). It is not clear whether all NRs are associated with these proteins; however, it appears that different NRs share pioneer factors as in the case of ER α , AR and retinoic acid receptor gamma (RAR γ) and the pioneer factors FoxA1 and the trans-acting T-cell-specific transcription factor (GATA3) (Lupien *et al.* 2008, Hua *et al.* 2009). Pioneer factors lack protein domains typically interacting with modified histones (e.g. bromo- or cromodomains), and although some data suggest that pioneer factors can recognize specific histone modifications (Lupien *et al.* 2008, Magnani *et al.* 2011a), more studies are needed to establish the exact epigenetic-pioneer factor-NR hierarchy (Magnani & Lupien 2013).

Post-induction histone modifications

Upon ligand binding, ER α recruits a large cohort of accessory proteins as co-activators and co-repressors (O'Malley & Kumar 2009, Mohammed *et al.* 2013). Interestingly, several chromatin modifiers belong to this class, including the histone acetyl transferase (HAT) p300 (Kraus & Kadonaga 1998), arginine methyltransferase CARM1 (Lee *et al.* 2005), histone deacetylase HDAC2 (Mohammed *et al.* 2013) and histone demethylase KDM1A (aka LSD1) (Hu *et al.* 2008). Similarly, the H3K9me3 histone demethylase KDM3A is recruited by AR and PPAR γ at NRREs (Yamane *et al.* 2006, Tateishi *et al.* 2009). There is a functional significance in HAT recruitment to the

chromatin as acetylation of histones destabilizes the nucleosome structure by neutralizing the basic charge of the lysine (Yamane *et al.* 2006, Tateishi *et al.* 2009). In addition, acetylated lysine acts as molecular bait for bromodomain-containing proteins, including SWI/SNF chromatin-remodelling complexes, another group of NR co-factors (Trotter & Archer 2008). Not surprisingly, chromatin remodellers and HAT proteins seem to trail very closely ER α binding dynamics in synchronized cells stimulated with E2 (Métivier *et al.* 2003).

Histone-modifying proteins also interact with other NRs. Known interactors of GR include HDAC2 (Ito *et al.* 2000), HDAC1 (Qiu *et al.* 2011) and CBP/p300 (Pfitzner *et al.* 1998). Assembly of such multi-tasking complexes is compatible with the dual roles of NRs as activators and repressors. It is also likely that co-activator/co-repressors are cell type specific, considering the diverse number of cellular processes controlled by NRs.

As described for hormone NRs, once activated by its own natural ligands (bile acids), FXR recruits a number of co-activator complexes able to remodel the chromatin structure and facilitate the assembly of the transcriptional machinery to activate target gene expression. Ananthanarayanan *et al.* (2004) demonstrated with ChIP assay experiments that upon ligand binding, a simultaneous increase in FXR and CARM1 occupation occurs at the FXR target *BSEP* (*ABCB11*) (bile salt export pump) gene promoter. This binding was found to correspond with increased deposition of Arg-17 methylation and Lys-9 acetylation of histone H3 within the FXREs of *BSEP*, leading to increased *BSEP* gene transactivation. Furthermore, FXR can be methylated by Set7/9 within its hinge region at lysine-206. ChIP assays demonstrated the recruitment of Set7/9 and the presence of the associated H3K4 monomethylation activation mark at the loci of two of the known FXR hepatic targets, the small heterodimer partner *NROB2* (*SHP*) and *BSEP*. This significantly enhanced the transcription of these two FXR target genes (Balasubramanian *et al.* 2012). Similar data for AR (Ko *et al.* 2011) and ER (Subramanian *et al.* 2008) indicate that this mechanism could also possibly stabilize FXR and promote its interaction with its heterodimerization partner RXR and the FXRE (Balasubramanian *et al.* 2012). Furthermore, recruitment of chromatin-remodelling proteins and NR-dependent gene transcription is a very dynamic process and different co-regulators and chromatin-remodelling proteins can occupy different REs of the same NR. In the hepatic FXR/SHP pathway, FXR senses elevated intrahepatic bile acid levels and indirectly suppresses the expression of the key-limiting enzyme of

the bile acid synthesis pathway, CYP7A1, via SHP. Chromatin immunoprecipitation and chromatin remodelling studies in both HepG2 cells and mouse liver have shown that upon treatment with FXR agonists, the transcription activator Brg-1 is recruited to the *SHP* promoter and it interacts with FXR, resulting in accessible chromatin, thereby enhancing FXR-dependent transactivation of *SHP*. On the other hand, Brm is recruited to both *SHP* and *CYP7A1* promoters, resulting in inactive, inaccessible chromatin (Miao *et al.* 2009). This indicates that chromatin remodellers Brm and Brg-1 have distinct functions in the regulation of two key genes within a single physiological pathway, by differentially modulating accessibility at SHP- and FXR-regulatory elements. In summary, these data demonstrate how NR binding facilitates the deposition of histone modifications, transiently altering the epigenetic profile of regulatory elements while locally increasing chromatin accessibility for the transcriptional machinery.

Epigenetic modification contributes to stimulus-specific activation of regulatory elements

NRs respond to a large array of molecules of diverse nature. Additionally, individual NRs can be promiscuous in terms of ligands. For example, metabolic NRs participate in complex, overlapping transcriptional regulation networks involving metabolic homeostasis. Some of these receptors, such as the pregnane X receptor (PXR) and the constitutive androstane receptor (CAR), are promiscuous with respect to the structurally wide range of ligands that act as agonists. In fact, they can bind to metabolites that are also ligands for other NRs, such as bile acids or steroid hormones and others (Kliewer & Willson 2002). This could be due to the presence of multiple LBDs, a very large binding pocket, or a binding site with flexibility to alter size and shape depending on the size of the ligand (reviewed in Krasowski *et al.* (2011)). In this context, the ER α has been the object of a substantial amount of work considering its role in breast cancer (Ali *et al.* 2011). ER α binding localization (its cistrome) has a significant impact on gene expression (Krum *et al.* 2008). Importantly, the ER α cistrome partially varies dependent on the type of ligand. Tamoxifen is one of the key molecules used in the treatment of breast cancer (Jordan 2003). Tamoxifen competes with E2 for the LBD of ER α and imparts a specific tridimensional structure not compatible with the recruitment of co-activators, at least in some tissues (Shang & Brown 2002). Recent evidence suggests that tamoxifen

alters the ER α -DNA binding profile (Welboren *et al.* 2009). In agreement, cells that acquire resistance to tamoxifen recruit ER α at a distinct set of loci compared with that of non-resistant cells (Ross-Innes *et al.* 2012). ER α can also be activated via growth factor-mediated phosphorylation induced by, for example, tyrosine kinase receptors such as the epidermal growth factor receptor (Kato *et al.* 1995). Genes regulated by phosphorylated ER α are distinct from oestrogen-responsive genes (Lupien *et al.* 2010). Importantly, oestrogen and EGF can induce ER α recruitment at three classes of enhancers: enhancers bound with stimulation with either oestrogen or EGF, enhancers bound exclusively following oestrogen treatment and enhancers bound exclusively following EGF treatment. These data are in agreement with ligand-specific NR binding (Lupien *et al.* 2010). However, the first class of regulatory elements (e.g. recruiting ER α in response to both stimuli) still exhibits a ligand-specific response as transcription is induced by only one ligand. This apparent contradiction is resolved taking into account ER α -mediated post-induction epigenetic modifications. Indeed, a fraction of these shared sites acquire stimulus-specific histone acetylation (H3K18ac; Lupien *et al.* 2010) and correlate with EGF- or oestrogen-regulated genes. A recent study probing different classes of ER α ligands has demonstrated that very minute changes in the chemical structure have significant effects on ER α -regulated cellular processes such as proliferation (Srinivasan *et al.* 2013). These results were observed even in response to different allosteric interactions of the same ligand (Srinivasan *et al.* 2013). Although the ER α cistrome was not assessed, it is conceivable that each ligand induces topologically different binding events.

Functional role of non-coding RNA in NR signalling

We have previously mentioned that a large class of NRs exhibit a significant tendency towards binding enhancers (Kittler *et al.* 2013). Recent work pioneered by studies on X-chromosome inactivation has highlighted a central role for long non-coding RNAs (lncRNAs) in epigenetic silencing and enhancer activation (Calabrese *et al.* 2012). Enhancer RNAs (eRNAs) are a class of lncRNAs with a potential role in the mediation of epigenetic modifications as well as interactions with TFs (Gupta *et al.* 2010, Kogo *et al.* 2011, Calabrese *et al.* 2012). Enhancer RNAs are typically bidirectional short RNAs transcribed by RNA polymerase II and centred around epigenetically defined enhancer elements (H3K4me1/2 and H3K27ac; Gupta *et al.* 2010, Kim *et al.* 2010, Kogo *et al.* 2011). They have

been implicated in several molecular processes including polycomb-mediated repression (Tsai *et al.* 2010), metastatic progression (Gupta *et al.* 2010) and enhancer activity (Orom & Shiekhattar 2011). Nonetheless, only recently, scientists have been able to elucidate their functional role partly through the use of NR-dependent cell lines (Orom & Shiekhattar 2011, Li *et al.* 2013). ER binding is thought to promote enhancer-promoter looping to activate the transcriptional machinery at target genes (Perillo *et al.* 2008, Fullwood *et al.* 2009, Li *et al.* 2012, 2013). The Rosenfeld group has described a strong relationship between E2-ER α -induced eRNA production and gene regulation (Li *et al.* 2013). Using siRNA and locked nucleic acid antisense oligonucleotides, the group was able to show that eRNA disruption suppresses E2-ER α -mediated transcription by destabilizing long-distance interactions between promoters and enhancers (Li *et al.* 2013). The production of eRNAs substantially correlates with numerous markers of enhancer activation, including post-induction epigenetic modifications and increased chromatin accessibility (Hah *et al.* 2013). Interestingly, eRNA production does not seem to influence pre-induction histone modifications such as H3K4me1 and H3K4me3 (Hah *et al.* 2013). However, Hah *et al.* (2013) suggest that eRNA production may be occurring after the assembly of the transcriptional machinery on the chromatin and is decoupled by promoter-enhancer interaction. While we are just beginning to understand eRNAs, their interaction with NRs seems to be conserved as demonstrated by orphan NRs Rev-ERB α and Rev-ERB β . In this case, Rev-Erb receptors function as strong repressors by suppressing eRNA transcription at lineage-specific enhancers (Lam *et al.* 2013). Altogether, these data suggest that NR activity at enhancers is far more complex than initially postulated.

Future directions

NRs have been studied extensively for over 30 years; nevertheless, we are still gathering new information about the way they function. Several aspects are yet to be defined, including how NRs cooperate physically and mechanistically with each other and the DNA structure. For example, in hormone-dependent cancers, it will be crucial to establish the role of understudied NRs and understand whether they could replace the classical sex hormone receptors once anti-cancer drugs have targeted ER α and AR. It will also be important to determine the role of unliganded NRs in development as well as disease. An intriguing study has found that unliganded PR forms a repressor complex with a non-coding RNA and several

epigenetic silencers including histone demethylases LSD1, KDM5B and histone deacetylases HDAC1/2 (Vicent *et al.* 2013). In other words, unactivated PR maintains repressed chromatin structures at progesterone target gene enhancers, while the epigenetic landscape is reversed upon the interaction of PR with its ligand (Vicent *et al.* 2013). These data further highlight the complex interaction between non-coding RNA, epigenetic modifications, NRs and transcription. More questions concerning NR binding topology would need an answer, such as how heterogeneous is the binding in normal and diseased tissues and how targeted therapies can alter NR binding profiles. Considering that enhancer elements are inherently variable between individuals (Kasowski *et al.* 2013), it is reasonable to expect that NRs will also target different regions in different patients. Indeed, a recent study that profiled ER α binding in breast cancer patients has found little consensus between individuals for most of the ER α -binding regions (Ross-Innes *et al.* 2012). Furthermore, in a study of hepatic genomic binding of ligand-activated FXR in healthy and obese mice, Lee *et al.* (2012) demonstrated that almost 50% of the sites were unique to either healthy or obese mice and that the altered FXR binding led to direct repression of many of these unique targets. Finally, it has been shown that the H3K4me3 epigenetic mark, essential for the transactivation of a number of hepatic FXR target genes, is reduced at their FXREs in a mouse model of cholestasis (Ananthanarayanan *et al.* 2004, Lee *et al.* 2012). Whether this is a reflection of an intrinsic heterogeneity in the epigenetic landscape of diseased tissues still requires clarification.

Ultimately, in the last few years, there has been a surge in studies addressing the fundamental role of chromatin and epigenetics in directing transcription. The inducible nature of NRs and the study of their interaction with the DNA allow us to mechanistically explore the dynamic nature of transcription and how diseases alter TF–chromatin crosstalk. Further understanding of these processes will contribute to the discovery of new relevant clinical implications (e.g. location-specific chromatin interactions as a prognostic factor) and the development of novel and more targeted treatments. Considering the importance of NRs in development, metabolism and disease and the high potential for pharmacological manipulation, we foresee new breakthroughs in the coming years.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review.

Funding

This work was supported by Imperial Junior Research Fellowships to L M and R M G.

Acknowledgements

The authors thank Dr Charlotte Bevan and Dr Simak Ali for critical reading of the manuscript.

References

- Ali S, Buluwela L & Coombes RC 2011 Antiestrogens and their therapeutic applications in breast cancer and other diseases. *Annual Review of Medicine* **62** 217–232. (doi:10.1146/annurev-med-052209-100305)
- Ananthanarayanan M, Li S, Balasubramanian N, Suchy FJ & Walsh MJ 2004 Ligand-dependent activation of the farnesoid X-receptor directs arginine methylation of histone H3 by CARM1. *Journal of Biological Chemistry* **279** 54348–54357. (doi:10.1074/jbc.M410021200)
- Andreu-Vieyra C, Lai J, Berman BP, Frenkel B, Jia L, Jones PA & Coetzee GA 2011 Dynamic nucleosome-depleted regions at androgen receptor enhancers in the absence of ligand in prostate cancer cells. *Molecular and Cellular Biology* **31** 4648–4662. (doi:10.1128/MCB.05934-11)
- Bain DL, Heneghan AF, Connaghan-Jones KD & Miura MT 2007 Nuclear receptor structure: implications for function. *Annual Review of Physiology* **69** 201–220. (doi:10.1146/annurev.physiol.69.031905.160308)
- Balasubramanian N, Ananthanarayanan M & Suchy FJ 2012 Direct methylation of FXR by Set7/9, a lysine methyltransferase, regulates the expression of FXR target genes. *American Journal of Physiology. Gastrointestinal and Liver Physiology* **302** G937–G947. (doi:10.1152/ajpgi.00441.2011)
- Belandia B, Orford RL, Hurst HC & Parker MG 2002 Targeting of SWI/SNF chromatin remodelling complexes to estrogen-responsive genes. *EMBO Journal* **21** 4094–4103. (doi:10.1093/emboj/cdf412)
- Berkes CA, Bergstrom DA, Penn BH, Seaver KJ, Knoepfler PS & Tapscott SJ 2004 Pbx marks genes for activation by MyoD indicating a role for a homeodomain protein in establishing myogenic potential. *Molecular Cell* **14** 465–477. (doi:10.1016/S1097-2765(04)00260-6)
- Bernstein BE, Mikkelsen TS, Xie X, Kamal M, Huebert DJ, Cuff J, Fry B, Meissner A, Wernig M, Plath K *et al.* 2006 A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* **125** 315–326. (doi:10.1016/j.cell.2006.02.041)
- Bolton EC, So AY, Chaivorapol C, Haqq CM, Li H & Yamamoto KR 2007 Cell- and gene-specific regulation of primary target genes by the androgen receptor. *Genes and Development* **21** 2005–2017. (doi:10.1101/gad.1564207)
- Bourguet W, Germain P & Gronemeyer H 2000 Nuclear receptor ligand-binding domains: three-dimensional structures, molecular interactions and pharmacological implications. *Trends in Pharmacological Sciences* **21** 381–388. (doi:10.1016/S0165-6147(00)01548-0)
- Calabrese JM, Sun W, Song L, Mugford JW, Williams L, Yee D, Starmer J, Mieczkowski P, Crawford GE & Magnuson T 2012 Site-specific silencing of regulatory elements as a mechanism of X inactivation. *Cell* **151** 951–963. (doi:10.1016/j.cell.2012.10.037)
- Carroll JS, Liu XS, Brodsky AS, Li W, Meyer CA, Szary AJ, Eeckhoutte J, Shao W, Hestermann EV, Geistlinger TR *et al.* 2005 Chromosome-wide mapping of estrogen receptor binding reveals long-range regulation requiring the forkhead protein FoxA1. *Cell* **122** 33–43. (doi:10.1016/j.cell.2005.05.008)
- Chambon P 2005 The nuclear receptor superfamily: a personal retrospect on the first two decades. *Molecular Endocrinology* **19** 1418–1428. (doi:10.1210/me.2005-0125)

- Chawla A, Boisvert WA, Lee CH, Laffitte BA, Barak Y, Joseph SB, Liao D, Nagy L, Edwards PA, Curtiss LK *et al.* 2001 A PPAR γ -LXR-ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis. *Molecular Cell* **7** 161–171. (doi:10.1016/S1097-2765(01)00164-2)
- Chen D, Pace PE, Coombes RC & Ali S 1999 Phosphorylation of human estrogen receptor α by protein kinase A regulates dimerization. *Molecular and Cellular Biology* **19** 1002–1015.
- Chong HK, Infante AM, Seo Y-K, Jeon T-I, Zhang Y, Edwards PA, Xie X & Osborne TF 2010 Genome-wide interrogation of hepatic FXR reveals an asymmetric IR-1 motif and synergy with LRH-1. *Nucleic Acids Research* **38** 6007–6017. (doi:10.1093/nar/gkq397)
- Christophersen NS & Helin K 2010 Epigenetic control of embryonic stem cell fate. *Journal of Experimental Medicine* **207** 2287–2295. (doi:10.1084/jem.20101438)
- Cirillo LA & Zaret KS 1999 An early developmental transcription factor complex that is more stable on nucleosome core particles than on free DNA. *Molecular Cell* **4** 961–969. (doi:10.1016/S1097-2765(00)80225-7)
- ENCODE Project Consortium, Birney E, Stamatoyannopoulos JA, Dutta A, Guigó R, Gingeras TR, Margulies EH, Weng Z, Snyder M, Dermitzakis ET *et al.* 2007 Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* **447** 799–816. (doi:10.1038/nature05874)
- ENCODE Project Consortium, Bernstein BE, Birney E, Dunham I, Green ED, Gunter C & Snyder M 2012 An integrated encyclopedia of DNA elements in the human genome. *Nature* **489** 57–74. (doi:10.1038/nature11247)
- Ernst J & Kellis M 2010 Discovery and characterization of chromatin states for systematic annotation of the human genome. *Nature Biotechnology* **28** 817–825. (doi:10.1038/nbt.1662)
- Ernst J, Kheradpour P, Mikkelsen TS, Shores N, Ward LD, Epstein CB, Zhang X, Wang L, Issner R, Coyne M *et al.* 2011 Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature* **473** 43–49. (doi:10.1038/nature09906)
- Evans RM 2005 The nuclear receptor superfamily: a rosetta stone for physiology. *Molecular Endocrinology* **19** 1429–1438. (doi:10.1210/me.2005-0046)
- Fang S, Tsang S, Jones R, Ponugoti B, Yoon H, Wu S-Y, Chiang C-M, Willson TM & Kemper JK 2008 The p300 acetylase is critical for ligand-activated farnesoid X receptor (FXR) induction of SHP. *Journal of Biological Chemistry* **283** 35086–35095. (doi:10.1074/jbc.M803531200)
- Fondell JD, Ge H & Roeder RG 1996 Ligand induction of a transcriptionally active thyroid hormone receptor coactivator complex. *PNAS* **93** 8329–8333. (doi:10.1073/pnas.93.16.8329)
- Fullwood MJ, Liu MH, Pan YF, Liu J, Xu H, Mohamed YB, Orlov YL, Velkov S, Ho A, Mei PH *et al.* 2009 An oestrogen-receptor- α -bound human chromatin interactome. *Nature* **462** 58–64. (doi:10.1038/nature08497)
- Germain P, Staels B, Dacquet C, Spedding M & Laudet V 2006 Overview of nomenclature of nuclear receptors. *Pharmacological Reviews* **58** 685–704. (doi:10.1124/pr.58.4.2)
- Giguère V, Hollenberg SM, Rosenfeld MG & Evans RM 1986 Functional domains of the human glucocorticoid receptor. *Cell* **46** 645–652. (doi:10.1016/0092-8674(86)90339-9)
- Gineste R, Sirvent A, Paumelle R, Helleboid S, Aquilina A, Darteil R, Hum DW, Fruchart J-C & Staels B 2008 Phosphorylation of farnesoid X receptor by protein kinase C promotes its transcriptional activity. *Molecular Endocrinology* **22** 2433–2447. (doi:10.1210/me.2008-0092)
- Giresi PG, Kim J, McDaniell RM, Iyer VR & Lieb JD 2007 FAIRE (formaldehyde-assisted isolation of regulatory elements) isolates active regulatory elements from human chromatin. *Genome Research* **17** 877–885. (doi:10.1101/gr.5533506)
- Greene GL, Gilna P, Waterfield M, Baker A, Hort Y & Shine J 1986 Sequence and expression of human estrogen receptor complementary DNA. *Science* **231** 1150–1154. (doi:10.1126/science.3753802)
- Gronemeyer H, Gustafsson J-A & Laudet V 2004 Principles for modulation of the nuclear receptor superfamily. *Nature Reviews. Drug Discovery* **3** 950–964. (doi:10.1038/nrd1551)
- Grøntved L & Hager GL 2012 Impact of chromatin structure on PR signaling: transition from local to global analysis. *Molecular and Cellular Endocrinology* **357** 30–36. (doi:10.1016/j.mce.2011.09.006)
- Gupta RA, Shah N, Wang KC, Kim J, Horlings HM, Wong DJ, Tsai M-C, Hung T, Argani P, Rinn JL *et al.* 2010 Long non-coding RNA HOTAIR reprograms chromatin state to promote cancer metastasis. *Nature* **464** 1071–1076. (doi:10.1038/nature08975)
- Hah N, Murakami S, Nagari A, Danko CG & Kraus WL 2013 Enhancer transcripts mark active estrogen receptor binding sites. *Genome Research* **23** 1210–1223. (doi:10.1101/gr.152306.112)
- He HH, Meyer CA, Chen MW, Jordan VC, Brown M & Liu XS 2012 Differential DNase I hypersensitivity reveals factor-dependent chromatin dynamics. *Genome Research* **22** 1015–1025. (doi:10.1101/gr.133280.111)
- Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H & Glass CK 2010 Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Molecular Cell* **38** 576–589. (doi:10.1016/j.molcel.2010.05.004)
- Heldring N, Isaacs GD, Diehl AG, Sun M, Cheung E, Ranish JA & Kraus WL 2011 Multiple sequence-specific DNA-binding proteins mediate estrogen receptor signaling through a tethering pathway. *Molecular Endocrinology* **25** 564–574. (doi:10.1210/me.2010-0425)
- Hollenberg SM, Weinberger C, Ong ES, Cerelli G, Oro A, Lebo R, Thompson EB, Rosenfeld MG & Evans RM 1985 Primary structure and expression of a functional human glucocorticoid receptor cDNA. *Nature* **318** 635–641. (doi:10.1038/318635a0)
- Hu Q, Kwon Y-S, Nunez E, Cardamone MD, Hutt KR, Ohgi KA, Garcia-Bassets I, Rose DW, Glass CK, Rosenfeld MG *et al.* 2008 Enhancing nuclear receptor-induced transcription requires nuclear motor and LSD1-dependent gene networking in interchromatin granules. *PNAS* **105** 19199–19204. (doi:10.1073/pnas.0810634105)
- Hua S, Kittler R & White KP 2009 Genomic antagonism between retinoic acid and estrogen signaling in breast cancer. *Cell* **137** 1259–1271. (doi:10.1016/j.cell.2009.04.043)
- Hurtado A, Holmes KA, Ross-Innes CS, Schmidt D & Carroll JS 2010 FOXA1 is a key determinant of estrogen receptor function and endocrine response. *Nature Genetics* **43** 27–33. (doi:10.1038/ng.730)
- Ito K, Barnes PJ & Adcock IM 2000 Glucocorticoid receptor recruitment of histone deacetylase 2 inhibits interleukin-1 β -induced histone H4 acetylation on lysines 8 and 12. *Molecular and Cellular Biology* **20** 6891–6903. (doi:10.1128/MCB.20.18.6891-6903.2000)
- Jeong KW, Lee Y-H & Stallcup MR 2009 Recruitment of the SWI/SNF chromatin remodeling complex to steroid hormone-regulated promoters by nuclear receptor coactivator flightless-I. *Journal of Biological Chemistry* **284** 29298–29309. (doi:10.1074/jbc.M109.037010)
- Jin C, Zang C, Wei G, Cui K, Peng W, Zhao K & Felsenfeld G 2009 H3.3/H2A.Z double variant-containing nucleosomes mark ‘nucleosome-free regions’ of active promoters and other regulatory regions. *Nature Genetics* **41** 941–945. (doi:10.1038/ng.409)
- John S, Sabo PJ, Johnson TA, Sung M-H, Biddie SC, Lightman SL, Voss TC, Davis SR, Meltzer PS, Stamatoyannopoulos JA *et al.* 2008 Interaction of the glucocorticoid receptor with the chromatin landscape. *Molecular Cell* **29** 611–624. (doi:10.1016/j.molcel.2008.02.010)
- Jordan VC 2003 Tamoxifen: a most unlikely pioneering medicine. *Nature Reviews. Drug Discovery* **2** 205–213. (doi:10.1038/nrd1031)
- Joseph R, Orlov YL, Huss M, Sun W, Kong SL, Ukil L, Pan YF, Li G, Lim M, Thomsen JS *et al.* 2010 Integrative model of genomic factors for determining binding site selection by estrogen receptor- α . *Molecular Systems Biology* **6** 456. (doi:10.1038/msb.2010.109)
- Jozwik KM & Carroll JS 2012 Pioneer factors in hormone-dependent cancers. *Nature Reviews. Cancer* **12** 381–385. (doi:10.1038/nrc3263)

- Kasowski M, Kyriazopoulou-Panagiotopoulou S, Grubert F, Zaugg JB, Kundaje A, Liu Y, Boyle AP, Zhang QC, Zakharia F, Spacek DV *et al.* 2013 Extensive variation in chromatin states across humans. *Science* **342** 750–752. (doi:10.1126/science.1242510)
- Kato S, Endoh H, Masuhiro Y, Kitamoto T, Uchiyama S, Sasaki H, Masushige S, Gotoh Y, Nishida E, Kawashima H *et al.* 1995 Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science* **270** 1491–1494. (doi:10.1126/science.270.5241.1491)
- Kim T-K, Hemberg M, Gray JM, Costa AM, Bear DM, Wu J, Harmin DA, Laptewicz M, Barbara-Haley K, Kuersten S *et al.* 2010 Widespread transcription at neuronal activity-regulated enhancers. *Nature* **465** 182–187. (doi:10.1038/nature09033)
- King HA, Trotter KW & Archer TK 2012 Chromatin remodeling during glucocorticoid receptor regulated transactivation. *Biochimica et Biophysica Acta* **1819** 716–726. (doi:10.1016/j.bbagr.2012.02.019)
- Kininis M & Kraus WL 2008 A global view of transcriptional regulation by nuclear receptors: gene expression, factor localization, and DNA sequence analysis. *Nuclear Receptor Signaling* **6** e005.
- Kittler R, Zhou J, Hua S, Ma L, Liu Y, Pendleton E, Cheng C, Gerstein M & White KP 2013 A comprehensive nuclear receptor network for breast cancer cells. *Cell Reports* **3** 538–551. (doi:10.1016/j.celrep.2013.01.004)
- Kliwer SA & Willson TM 2002 Regulation of xenobiotic and bile acid metabolism by the nuclear pregnane X receptor. *Journal of Lipid Research* **43** 359–364.
- Ko S, Ahn J, Song CS, Kim S, Knapczyk-Stwora K & Chatterjee B 2011 Lysine methylation and functional modulation of androgen receptor by Set9 methyltransferase. *Molecular Endocrinology* **25** 433–444. (doi:10.1210/me.2010-0482)
- Kogo R, Shimamura T, Mimori K, Kawahara K, Imoto S, Sudo T, Tanaka F, Shibata K, Suzuki A, Komune S *et al.* 2011 Long noncoding RNA HOTAIR regulates polycomb-dependent chromatin modification and is associated with poor prognosis in colorectal cancers. *Cancer Research* **71** 6320–6326. (doi:10.1158/0008-5472.CAN-11-1021)
- Kouzarides T 2007 Chromatin modifications and their function. *Cell* **128** 693–705. (doi:10.1016/j.cell.2007.02.005)
- Krasowski MD, Ni A, Hagey LR & Ekins S 2011 Evolution of promiscuous nuclear hormone receptors: LXR, FXR, VDR, PXR, and CAR. *Molecular and Cellular Endocrinology* **334** 39–48. (doi:10.1016/j.mce.2010.06.016)
- Kraus WL & Kadonaga JT 1998 p300 and estrogen receptor cooperatively activate transcription via differential enhancement of initiation and reinitiation. *Genes and Development* **12** 331–342. (doi:10.1101/gad.12.3.331)
- Krum SA, Miranda-Carboni GA, Lupien M, Eeckhoutte J, Carroll JS & Brown M 2008 Unique ER α cistromes control cell type-specific gene regulation. *Molecular Endocrinology* **22** 2393–2406. (doi:10.1210/me.2008-0100)
- Lai C-F, Flach KD, Alexi X, Fox SP, Ottaviani S, Thiruchelvam PTR, Kyle FJ, Thomas RS, Launchbury R, Hua H *et al.* 2013 Co-regulated gene expression by oestrogen receptor α and liver receptor homolog-1 is a feature of the oestrogen response in breast cancer cells. *Nucleic Acids Research* **41** 10228–10240. (doi:10.1093/nar/gkt827)
- Lam MTY, Cho H, Lesch HP, Gosselin D, Heinz S, Tanaka-Oishi Y, Benner C, Kaikkonen MU, Kim AS, Kosaka M *et al.* 2013 Rev-Erbs repress macrophage gene expression by inhibiting enhancer-directed transcription. *Nature* **498** 511–515. (doi:10.1038/nature12209)
- Lee Y-H, Coonrod SA, Kraus WL, Jelinek MA & Stallcup MR 2005 Regulation of coactivator complex assembly and function by protein arginine methylation and demethylation. *PNAS* **102** 3611–3616. (doi:10.1073/pnas.0407159102)
- Lee J, Seok S, Yu P, Kim K, Smith Z, Rivas-Astroza M, Zhong S & Kemper JK 2012 Genomic analysis of hepatic farnesoid X receptor binding sites reveals altered binding in obesity and direct gene repression by farnesoid X receptor in mice. *Hepatology* **56** 108–117. (doi:10.1002/hep.25609)
- Lefterova MI, Zhang Y, Steger DJ, Schupp M, Schug J, Cristancho A, Feng D, Zhuo D, Stoeckert CJ, Liu XS *et al.* 2008 PPAR γ and C/EBP factors orchestrate adipocyte biology via adjacent binding on a genome-wide scale. *Genes and Development* **22** 2941–2952. (doi:10.1101/gad.1709008)
- Li G, Ruan X, Auerbach RK, Sandhu KS, Zheng M, Wang P, Poh HM, Goh Y, Lim J, Zhang J *et al.* 2012 Extensive promoter-centered chromatin interactions provide a topological basis for transcription regulation. *Cell* **148** 84–98. (doi:10.1016/j.cell.2011.12.014)
- Li W, Notani D, Ma Q, Tanasa B, Nunez E, Chen AY, Merkurjev D, Zhang J, Ohgi K, Song X *et al.* 2013 Functional roles of enhancer RNAs for oestrogen-dependent transcriptional activation. *Nature* **498** 516–520. (doi:10.1038/nature12210)
- Lupien M, Eeckhoutte J, Meyer CA, Wang Q, Zhang Y, Li W, Carroll JS, Liu XS & Brown M 2008 FoxA1 translates epigenetic signatures into enhancer-driven lineage-specific transcription. *Cell* **132** 958–970. (doi:10.1016/j.cell.2008.01.018)
- Lupien M, Meyer CA, Bailey ST, Eeckhoutte J, Cook J, Westerling T, Zhang X, Carroll JS, Rhodes DR, Liu XS *et al.* 2010 Growth factor stimulation induces a distinct ER(α) cistrome underlying breast cancer endocrine resistance. *Genes and Development* **24** 2219–2227. (doi:10.1101/gad.1944810)
- Magnani L & Lupien M 2013 Chromatin and epigenetic determinants of estrogen receptor α (ESR1) signaling. *Molecular and Cellular Endocrinology* **382** 633–641. (doi:10.1016/j.mce.2013.04.026)
- Magnani L, Ballantyne EB, Zhang X & Lupien M 2011a PBX1 genomic pioneer function drives ER α signaling underlying progression in breast cancer. *PLoS Genetics* **7** e1002368. (doi:10.1371/journal.pgen.1002368)
- Magnani L, Eeckhoutte J & Lupien M 2011b Pioneer factors: directing transcriptional regulators within the chromatin environment. *Trends in Genetics* **27** 465–474. (doi:10.1016/j.tig.2011.07.002)
- Mangelsdorf DJ & Evans RM 1995 The RXR heterodimers and orphan receptors. *Cell* **83** 841–850. (doi:10.1016/0092-8674(95)90200-7)
- Mangelsdorf DJ, Thummel C, Beato M, Herrlich P, Schütz G, Umesono K, Blumberg B, Kastner P, Mark M, Chambon P *et al.* 1995 The nuclear receptor superfamily: the second decade. *Cell* **83** 835–839. (doi:10.1016/0092-8674(95)90199-X)
- Métivier R, Penot G, Hübner MR, Reid G, Brand H, Kos M & Gannon F 2003 Estrogen receptor- α directs ordered, cyclical, and combinatorial recruitment of cofactors on a natural target promoter. *Cell* **115** 751–763. (doi:10.1016/S0092-8674(03)00934-6)
- Métivier R, Gallais R, Tiffiche C, Le Péron C, Jurkowska RZ, Carmouche RP, Ibberson D, Barath P, Demay F, Reid G *et al.* 2008 Cyclical DNA methylation of a transcriptionally active promoter. *Nature* **452** 45–50. (doi:10.1038/nature06544)
- Miao J, Fang S, Lee J, Comstock C, Knudsen KE & Kemper JK 2009 Functional specificities of Brm and Brg-1 Swi/Snf ATPases in the feedback regulation of hepatic bile acid biosynthesis. *Molecular and Cellular Biology* **29** 6170–6181. (doi:10.1128/MCB.00825-09)
- Miranda TB, Voss TC, Sung M-H, Baek S, John S, Hawkins M, Grøntved L, Schiltz RL & Hager GL 2013 Reprogramming the chromatin landscape: interplay of the estrogen and glucocorticoid receptors at the genomic level. *Cancer Research* **73** 5130–5139. (doi:10.1158/0008-5472.CAN-13-0742)
- Mohammed H, D'Santos C, Sérandour AA, Ali HR, Brown GD, Atkins A, Rueda OM, Holmes KA, Theodorou V, Robinson JLL *et al.* 2013 Endogenous purification reveals GREB1 as a key estrogen receptor regulatory factor. *Cell Reports* **3** 342–349. (doi:10.1016/j.celrep.2013.01.010)
- Nielsen R, Pedersen TA, Hagenbeek D, Moulos P, Siersbaek R, Megens E, Denissov S, Børgesen M, Francois K-J, Mandrup S *et al.* 2008 Genome-wide profiling of PPAR γ :RXR and RNA polymerase II occupancy reveals temporal activation of distinct metabolic pathways and changes in RXR dimer composition during adipogenesis. *Genes and Development* **22** 2953–2967. (doi:10.1101/gad.501108)

- Nuclear Receptors Nomenclature Committee 1999 A unified nomenclature system for the nuclear receptor superfamily. *Cell* **97** 161–163. (doi:10.1016/S0092-8674(00)80726-6)
- O'Malley BW & Kumar R 2009 Nuclear receptor coregulators in cancer biology. *Cancer Research* **69** 8217–8222. (doi:10.1158/0008-5472.CAN-09-2223)
- Orom UA & Shiekhattar R 2011 Long non-coding RNAs and enhancers. *Current Opinion in Genetics & Development* **21** 194–198. (doi:10.1016/j.gde.2011.01.020)
- Overington JP, Al-Lazikani B & Hopkins AL 2006 How many drug targets are there? *Nature Reviews. Drug Discovery* **5** 993–996. (doi:10.1038/nrd2199)
- Pascual G, Fong AL, Ogawa S, Gamliel A, Li AC, Perissi V, Rose DW, Willson TM, Rosenfeld MG & Glass CK 2005 A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR- γ . *Nature* **437** 759–763. (doi:10.1038/nature03988)
- Perillo B, Ombra MN, Bertoni A, Cuozzo C, Sacchetti S, Sasso A, Chiariotti L, Malorni A, Abbondanza C & Avvedimento EV 2008 DNA oxidation as triggered by H3K9me2 demethylation drives estrogen-induced gene expression. *Science* **319** 202–206. (doi:10.1126/science.1147674)
- Pfützner E, Jähne R, Wissler M, Stoecklin E & Groner B 1998 p300/CREB-binding protein enhances the prolactin-mediated transcriptional induction through direct interaction with the transactivation domain of Stat5, but does not participate in the Stat5-mediated suppression of the glucocorticoid response. *Molecular Endocrinology* **12** 1582–1593. (doi:10.1210/me.12.10.1582)
- Qiu Y, Stavreva DA, Luo Y, Indrawan A, Chang M & Hager GL 2011 Dynamic interaction of HDAC1 with a glucocorticoid receptor-regulated gene is modulated by the activity state of the promoter. *Journal of Biological Chemistry* **286** 7641–7647. (doi:10.1074/jbc.M110.185488)
- Reid G, Hübner MR, Métivier R, Brand H, Denger S, Manu D, Beaudouin J, Ellenberg J & Gannon F 2003 Cyclic, proteasome-mediated turnover of unliganded and liganded ER α on responsive promoters is an integral feature of estrogen signaling. *Molecular Cell* **11** 695–707. (doi:10.1016/S1097-2765(03)00090-X)
- Rizzo G, Renga B, Antonelli E, Passeri D, Pellicciari R & Fiorucci S 2005 The methyl transferase PRMT1 functions as co-activator of farnesoid X receptor (FXR)/9-cis retinoid X receptor and regulates transcription of FXR responsive genes. *Molecular Pharmacology* **68** 551–558. (doi:10.1124/mol.105.012104)
- Ross-Innes CS, Stark R, Teschendorff AE, Holmes KA, Ali HR, Dunning MJ, Brown GD, Gojis O, Ellis IO, Green AR *et al.* 2012 Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. *Nature* **481** 389–393.
- Sap J, Muñoz A, Damm K, Goldberg Y, Ghysdael J, Leutz A, Beug H & Vennström B 1986 The c-erb-A protein is a high-affinity receptor for thyroid hormone. *Nature* **324** 635–640. (doi:10.1038/324635a0)
- Shang Y & Brown M 2002 Molecular determinants for the tissue specificity of SERMs. *Science* **295** 2465–2468. (doi:10.1126/science.1068537)
- Srinivasan S, Nwachukwu JC, Parent AA, Cavett V, Nowak J, Hughes TS, Kojetin DJ, Katzenellenbogen JA & Nettles KW 2013 Ligand-binding dynamics wire cellular signaling via estrogen receptor- α . *Nature Chemical Biology* **9** 326–332. (doi:10.1038/nchembio.1214)
- Stender JD, Kim K, Charn TH, Komm B, Chang KCN, Kraus WL, Benner C, Glass CK & Katzenellenbogen BS 2010 Genome-wide analysis of estrogen receptor α DNA binding and tethering mechanisms identifies Runx1 as a novel tethering factor in receptor-mediated transcriptional activation. *Molecular and Cellular Biology* **30** 3943–3955. (doi:10.1128/MCB.00118-10)
- Stratmann M & Schibler U 2011 Transcription factor loading: please take my place! *Cell* **146** 497–499. (doi:10.1016/j.cell.2011.07.037)
- Subramanian K, Jia D, Kapoor-Vazirani P, Powell DR, Collins RE, Sharma D, Peng J, Cheng X & Vertino PM 2008 Regulation of estrogen receptor α by the SET7 lysine methyltransferase. *Molecular Cell* **30** 336–347. (doi:10.1016/j.molcel.2008.03.022)
- Suganuma T & Workman JL 2008 Crosstalk among histone modifications. *Cell* **135** 604–607. (doi:10.1016/j.cell.2008.10.036)
- Tateishi K, Okada Y, Kallin EM & Zhang Y 2009 Role of Jhdm2a in regulating metabolic gene expression and obesity resistance. *Nature* **458** 757–761. (doi:10.1038/nature07777)
- Tewari AK, Yardimci GG, Shibata Y, Sheffield NC, Song L, Taylor BS, Georgiev SG, Coetzee GA, Ohler U, Furey TS *et al.* 2012 Chromatin accessibility reveals insights into androgen receptor activation and transcriptional specificity. *Genome Biology* **13** R88. (doi:10.1186/gb-2012-13-10-r88)
- Thomas AM, Hart SN, Kong B, Fang J, Zhong X-B & Guo GL 2010 Genome-wide tissue-specific farnesoid X receptor binding in mouse liver and intestine. *Hepatology* **51** 1410–1419. (doi:10.1002/hep.23450)
- Thurman RE, Rynes E, Humbert R, Vierstra J, Maurano MT, Haugen E, Sheffield NC, Stergachis AB, Wang H, Vernot B *et al.* 2012 The accessible chromatin landscape of the human genome. *Nature* **489** 75–82. (doi:10.1038/nature11232)
- Trotter KW & Archer TK 2008 The BRG1 transcriptional coregulator. *Nuclear Receptor Signaling* **6** e004.
- Tsai M-C, Manor O, Wan Y, Mosammaparast N, Wang JK, Lan F, Shi Y, Segal E & Chang HY 2010 Long noncoding RNA as modular scaffold of histone modification complexes. *Science* **329** 689–693. (doi:10.1126/science.1192002)
- Tzagarakis-Foster C & Privalsky ML 1998 Phosphorylation of thyroid hormone receptors by protein kinase A regulates DNA recognition by specific inhibition of receptor monomer binding. *Journal of Biological Chemistry* **273** 10926–10932. (doi:10.1074/jbc.273.18.10926)
- Vavassori P, Mencarelli A, Renga B, Distrutti E & Fiorucci S 2009 The bile acid receptor FXR is a modulator of intestinal innate immunity. *Journal of Immunology* **183** 6251–6261. (doi:10.4049/jimmunol.0803978)
- Vicent GP, Nacht AS, Font-Mateu J, Castellano G, Gaveglia L, Ballaré C & Beato M 2011 Four enzymes cooperate to displace histone H1 during the first minute of hormonal gene activation. *Genes and Development* **25** 845–862. (doi:10.1101/gad.621811)
- Vicent GP, Nacht AS, Zaurin R, Font-Mateu J, Soronellas D, Le Dily F, Reyes D & Beato M 2013 Unliganded progesterone receptor-mediated targeting of an RNA-containing repressive complex silences a subset of hormone-inducible genes. *Genes and Development* **27** 1179–1197. (doi:10.1101/gad.215293.113)
- Voss TC, Schiltz RL, Sung M-H, Yen PM, Stamatoiyannopoulos JA, Biddie SC, Johnson TA, Miranda TB, John S & Hager GL 2011 Dynamic exchange at regulatory elements during chromatin remodeling underlies assisted loading mechanism. *Cell* **146** 544–554. (doi:10.1016/j.cell.2011.07.006)
- Walter P, Green S, Greene G, Krust A, Bornert JM, Jeltsch JM, Staub A, Jensen E, Scrae G & Waterfield M 1985 Cloning of the human estrogen receptor cDNA. *PNAS* **82** 7889–7893. (doi:10.1073/pnas.82.23.7889)
- Wang Q, Li W, Zhang Y, Yuan X, Xu K, Yu J, Chen Z, Beroukhim R, Wang H, Lupien M *et al.* 2009 Androgen receptor regulates a distinct transcription program in androgen-independent prostate cancer. *Cell* **138** 245–256. (doi:10.1016/j.cell.2009.04.056)
- Wärnmark A, Treuter E, Wright APH & Gustafsson J-A 2003 Activation functions 1 and 2 of nuclear receptors: molecular strategies for transcriptional activation. *Molecular Endocrinology* **17** 1901–1909. (doi:10.1210/me.2002-0384)
- Welboren W-J, van Driel MA, Janssen-Megens EM, van Heeringen SJ, Sweep FC, Span PN & Stunnenberg HG 2009 ChIP-Seq of ER α and RNA polymerase II defines genes differentially responding to ligands. *EMBO Journal* **28** 1418–1428. (doi:10.1038/emboj.2009.88)
- Whyte WA, Orlando DA, Hnisz D, Abraham BJ, Lin CY, Kagey MH, Rahl PB, Lee TI & Young RA 2013 Master transcription factors and mediator establish super-enhancers at key cell identity genes. *Cell* **153** 307–319. (doi:10.1016/j.cell.2013.03.035)

- Willis S & Griffin PR 2011 Mutual information identifies sequence positions conserved within the nuclear receptor superfamily: approach reveals functionally important regions for DNA binding specificity. *Nuclear Receptor Signaling* **9** e001.
- Willy PJ, Umesono K, Ong ES, Evans RM, Heyman RA & Mangelsdorf DJ 1995 LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Genes and Development* **9** 1033–1045. (doi:10.1101/gad.9.9.1033)
- Wu Y, Chin WW, Wang Y & Burris TP 2003 Ligand and coactivator identity determines the requirement of the charge clamp for coactivation of the peroxisome proliferator-activated receptor γ . *Journal of Biological Chemistry* **278** 8637–8644. (doi:10.1074/jbc.M210910200)
- Yamane K, Toumazou C, Tsukada Y-I, Erdjument-Bromage H, Tempst P, Wong J & Zhang Y 2006 JHDM2A, a JmJC-containing H3K9 demethylase, facilitates transcription activation by androgen receptor. *Cell* **125** 483–495. (doi:10.1016/j.cell.2006.03.027)
- Zaret KS & Carroll JS 2011 Pioneer transcription factors: establishing competence for gene expression. *Genes and Development* **25** 2227–2241. (doi:10.1101/gad.176826.111)
- Zaret KS, Watts J, Xu J, Wandzioch E, Smale ST & Sekiya T 2008 Pioneer factors, genetic competence, and inductive signaling: programming liver and pancreas progenitors from the endoderm. *Cold Spring Harbor Symposia on Quantitative Biology* **73** 119–126. (doi:10.1101/sqb.2008.73.040)
- Zeuchel C, Shen XQ, Chambon P & Gronemeyer H 1994 Dimerization interfaces formed between the DNA binding domains determine the cooperative binding of RXR/RAR and RXR/TR heterodimers to DR5 and DR4 elements. *EMBO Journal* **13** 1414–1424.
- Zhu J, Adli M, Zou JY, Verstappen G, Coyne M, Zhang X, Durham T, Miri M, Deshpande V, De Jager PL *et al.* 2013 Genome-wide chromatin state transitions associated with developmental and environmental cues. *Cell* **152** 642–654. (doi:10.1016/j.cell.2012.12.033)

Received in final form 28 November 2013

Accepted 19 December 2013

Accepted Preprint published online 20 December 2013