A new pathway in the control of the initiation of puberty: the *MKRN3* gene

Ana Paula Abreu, Delanie B Macedo¹, Vinicius N Brito¹, Ursula B Kaiser and Ana Claudia Latronico¹

Division of Endocrinology, Diabetes and Hypertension, Harvard Medical School, Brigham and Women's Hospital, Boston, Massachusetts, USA

¹Unidade de Endocrinologia do Desenvolvimento, Disciplina de Endocrinologia e Metabologia, Laboratório de Hormônios e Genética Molecular, LIM 42, Hospital das Clínicas, Faculdade de Medicina da Universidade de São Paulo, Avenida Dr Enéas de Carvalho Aguiar, 255, 7º andar, sala 7037, CEP: 05403-900, São Paulo, Brazil

Correspondence should be addressed to A C Latronico
Email
anacl@usp.br

Abstract

Pubertal timing is influenced by complex interactions among genetic, nutritional, environmental, and socioeconomic factors. The role of *MKRN3*, an imprinted gene located in the Prader–Willi syndrome critical region (chromosome 15q11–13), in pubertal initiation was first described in 2013 after the identification of deleterious *MKRN3* mutations in five families with central precocious puberty (CPP) using whole-exome sequencing analysis. Since then, additional loss-of-function mutations of *MKRN3* have been associated with the inherited premature sexual development phenotype in girls and boys from different ethnic groups. In all of these families, segregation analysis clearly demonstrated autosomal dominant inheritance with complete penetrance, but with exclusive paternal transmission, consistent with the monoallelic expression of *MKRN3* (a maternally imprinted gene). Interestingly, the hypothalamic *Mkfn3* mRNA expression pattern in mice correlated with a putative inhibitory input on puberty initiation. Indeed, the initiation of puberty depends on a decrease in factors that inhibit the release of GnRH combined with an increase in stimulatory factors. These recent human and animal findings suggest that *MKRN3* plays an inhibitory role in the reproductive axis to represent a new pathway in pubertal regulation.

Key Words

- ▶ hypothalamus and neuroendocrinology
- ▶ gonadotropins
- ▶ mutations
- ▶ secretion

Journal of Molecular Endocrinology
(2015) **54**, R131–R139

Introduction

Puberty, a complex biological process involving sexual maturation and accelerated linear growth, is initiated when the pulsatile secretion of gonadotropin-releasing hormone (GnRH) increases after a quiescent period during childhood. The regulation of puberty initiation remains as one of the great mysteries of human biology and it is thought that a conjunction of factors plays a role to initiate puberty.

Environmental and metabolic factors are important regulators of pubertal development, but these influences

are superimposed upon substantial genetic control. Similar timing of puberty shared by mothers and her children, and within racial groups, suggests a genetic component to the onset of puberty (Zacharias & Wurtman 1969, Herman-Giddens *et al.* 1997). Substantial efforts have been made to identify genes that are responsible for the initiation of puberty (Ojeda & Lomniczi 2014). The identification of these genes is critical for advancing the understanding of the neuroendocrine regulation of puberty initiation.

In an attempt to identify genes operating within the neuroendocrine brain that ultimately regulate the reproductive axis, researchers have been studying patients with pubertal disorders. The hypothalamic–pituitary–gonadal axis regulates puberty initiation and reproduction. GnRH is produced and secreted by the hypothalamus in a pulsatile manner during the embryonic and neonatal phases of life. GnRH secretion is actively inhibited during infancy and puberty begins with the reactivation of its secretion. GnRH deficiency results in hypogonadotropic hypogonadism, in which patients fail to develop puberty and are usually infertile. Conversely, early reactivation of GnRH secretion results in central precocious puberty (CPP). While several genes have been detected in association with GnRH deficiency and have contributed to the current knowledge of GnRH regulation (Bianco & Kaiser 2009, Semple & Topaloglu 2010), genes linked to CPP have until recently only been identified subsequent to their association with hypogonadotropic hypogonadism, such as *KISS1* and *KISS1R*. However, only rare genetic defects in *KISS1* and its receptor have been identified in patients with CPP (Teles *et al.* 2008, Silveira *et al.* 2010).

The advances in sequencing methods permitted the detection of new genes implicated in the neuroendocrine regulation of puberty. Using exome sequencing analysis and studying familial cases of CPP, genetic defects in a gene with no previous link to the hypothalamic–pituitary–gonadal axis, Makorin ring finger 3 (*MKRN3*), were identified as the cause of premature sexual development in one-third of the families (Abreu *et al.* 2013). *MKRN3* is located on the long arm of chromosome 15 in the

Prader–Willi syndrome (PWS) critical region and is maternally imprinted (discussed later). Subsequently, we showed that mutations in *MKRN3* are also the cause of CPP in apparently sporadic cases (Macedo *et al.* 2014). These findings were significant contributions to the field and will advance the understanding of the pubertal process. We will review the genetic defects identified in patients with CPP and their clinical implications, and discuss herein the possible role of *MKRN3* within the reproductive axis.

Loss-of-function mutations of *MKRN3* cause familial CPP

The role of *MKRN3* in pubertal initiation was first described in 2013 after a comprehensive genetic study of several families with CPP (Abreu *et al.* 2013). In this study, the authors investigated 40 members of 15 families with CPP from different ethnic groups (12 Brazilian, two American, and one Belgian families), and applying whole-exome sequencing four deleterious *MKRN3* mutations – three frameshift and a missense mutation (Fig. 1) – were detected in five of these families (33%). Both sexes were equally affected by *MKRN3* mutations (eight girls and seven boys) (Table 1).

More recently, Macedo *et al.* (2014) studied 215 unrelated children (207 girls and eight boys) with CPP from three different University Medical Centers and identified five novel heterozygous mutations in eight unrelated Brazilian girls, including four frameshift variants and one missense variant (Fig. 1 and Table 1). No family history of premature sexual development was

Figure 1

MKRN3 protein structure and mutations identified in patients with central precocious puberty. Zn, zinc; H, histidine; C, cysteine. The three C3H zinc finger motifs are shown in red, the C3HC4 RING finger motif is in blue, and the MKRN-specific Cys–His domain is shown in green. The numbers

correspond to the amino acid positions in the protein. Blue mutation labels and arrows indicate the location of frameshift mutations; red mutation labels and arrows indicate the location of the missense mutations.

Table 1 MKRN3 mutations identified in children with CPP. (Gonadarche or testicular enlargement. Breast/Genital development according to Tanner stage and bone age assessed at the time of diagnosis)

Family number	Origin	Patient number	Sex	Initial clinical manifestation (age in years)	Characteristics at time of diagnosis			MKRN3 mutation		References
					Age (years)	Breast/Genital development Tanner stage	Bone age (years)	DNA	Protein	
1	USA	I	F	Thelarche (5.7)	6.5	2	7.7	637delC	Arg213Glyfs*73	Abreu et al. (2013)
		II	M	Gonadarche (8.0)	8.7	3	11	637delC	Arg213Glyfs*73	
2	Brazil	III	F	Thelarche (6.5)	6.7	2	7.8	637delC	Arg213Glyfs*73	Abreu et al. (2013)
		I	F	Thelarche (6.2)	7.0	3	7.0	1171_1172insA	Tyr391*	
		II	F	Thelarche (5.7)	6.0	3	6.0	1171_1172insA	Tyr391*	
3	Brazil	I	M	Gonadarche and pubarche (5.9)	8.1	3	10	475_476insC	Ala162Glyfs*14	Abreu et al. (2013)
		II	M	Gonadarche and pubarche unknown	9.7	3	9.7	475_476insC	Ala162Glyfs*14	
4	USA	I	M	Gonadarche and pubarche (8.5)	8.8	3	11	475_476insC	Ala162Glyfs*14	Abreu et al. (2013)
		II	F	Thelarche (5.0)	6.5	2	8.3	475_476insC	Ala162Glyfs*14	
5	Belgium	I	F	Thelarche (6.2)	6.4	2	9.4 ^a	1095G>T	Arg365Ser	Abreu et al. (2013)
		II	M	Gonadarche and pubarche unknown	9.7	3	12	1095G>T	Arg365Ser	
6	Brazil	III	F	Thelarche and pubarche (5.4)	5.7	2	8.5	1095G>T	Arg365Ser	Macedo et al. (2014)
7	Brazil	I	F	Thelarche and pubarche (6.4)	7.8	3	11	482delC	Pro161Argfs*10	Macedo et al. (2014)
8	Brazil	I	F	Thelarche (5.4)	6.1	3	8.8	482_483insC	Pro161Argfs*16	Macedo et al. (2014)
9	Brazil	I	F	Thelarche (6.0)	7.9	4	11	482_483insC	Pro161Argfs*16	Macedo et al. (2014)
10	Brazil	I	F	Thelarche (3.0) and pubarche (6.0)	6.7	3	8.8	482delC	Pro161Argfs*10	Macedo et al. (2014)
11	Brazil	I	F	Thelarche (4.0)	6.8	4	12	675_676insA	Gln226Thrfs*6	Macedo et al. (2014)
12	Brazil	I	F	Thelarche and pubarche (6.0)	6.6	3	7.8	766_767delA	Glu256Glyfs*36	Macedo et al. (2014)
13	Brazil	I	F	Thelarche (6.0) and pubarche (6.3)	6.9	3	9.3	482_483insC	Pro161Argfs*16	Macedo et al. (2014)
14	Greece	I	F	Thelarche (6.0)	7.1	3	8.8	1249T>A	Phe417Ile	Macedo et al. (2014)
		II	F	Thelarche (6.0)	7.1	3	10	1018T>G	Cys340Gly	Settas et al. (2014)
15	Germany	I	M	Gonadarche unknown	9.2	2	10	1018T>G	Cys340Gly	Schreiner et al. 2014
		II	M	Gonadarche and pubarche (9.0)	10	3	–	331G>T	Glu111*	
16	Germany	II	F	Thelarche (6.1)	7.5	3	–	331G>T	Glu111*	Schreiner et al. (2014)
		I	F	Thelarche (7.5)	–	–	–	475_476insC	Ala162Glyfs*14	
		II	F	Thelarche (6.8)	7.1	2	–	475_476insC	Ala162Glyfs*14	
17	Israel	I	F	Thelarche and pubarche (5.5)	5.8	3	6.8	1260T>G	His420Gln	de Vries et al. (2014)
		II	M	Gonadarche and pubarche unknown	9.8	3	11	1260T>G	His420Gln	
		III	F	Thelarche and pubarche (5.5)	6.3	3	8.8	1260T>G	His420Gln	
		IV	F	Thelarche (4.5)	4.5	2	6.0	1260T>G	His420Gln	

F, female; M, male.

^aBone age at 7.4 years.

reported in the majority of these studied patients. However, segregation analysis was performed for five of these eight girls, and in all cases it has been demonstrated that the mutant allele was paternally inherited in all families with *MKRN3* mutations.

In order to investigate whether the CPP phenotype could arise from loss of *MKRN3* expression by the paternal allele due to a *de novo* deletion, maternal uniparental disomy, or an imprinting defect, Macedo *et al.* (2014) investigated 52 patients with familial and sporadic CPP, without known *MKRN3* sequence defects, by methylation-specific multiplex ligation-dependent probe amplification (MS-MLPA) of chromosome 15q11. No copy number changes or methylation abnormalities were detected in the 15q11 *locus*, suggesting that epigenetic defects involving this *locus* are likely rare mechanisms in this disorder (Macedo *et al.* 2014).

Recently, other investigators have also reported *MKRN3* defects associated with familial CPP. Settas *et al.* (2014) described a novel heterozygous missense mutation (p.Cys340Gly) in *MKRN3* in two Greek siblings, a girl with CPP and a boy with early puberty. Soon thereafter, Schreiner *et al.* (2014) identified two heterozygous *MKRN3* mutations (p.Glu111* and p.Ala162Glyfs*14) in affected members of two German families. Finally, de Vries *et al.* (2014) identified a novel missense mutation (p.His420Gln) in four siblings, including a boy, from an Ashkenazi–Sephardic Jewish family (Fig. 1 and Table 1). These studies further expanded the *MKRN3* mutational spectrum.

Clinical features of CPP associated with *MKRN3* mutations

All patients with CPP carrying loss-of-function mutations in *MKRN3* exhibited typical clinical and hormonal features of premature activation of the reproductive axis, including early pubertal signs, such as breast, testes, and pubic hair development, accelerated linear growth, advanced bone age, and elevated basal and/or GnRH-stimulated LH levels (Table 1). Except for two related patients (one girl and her brother) who presented with esotropia (Abreu *et al.* 2013), which is a minor criterion for PWS, no other signs of PWS were detected in any patient with CPP caused by *MKRN3* defects. Another female patient had mild nonspecific syndromic features, including a high-arched palate, dental abnormalities, clinodactyly, and hyperlordosis (Macedo *et al.* 2014). Regarding the therapeutic response of CPP patients with *MKRN3* defects, six of eight patients have demonstrated adequate

responses to conventional treatment with depot GnRH agonists to date (Macedo *et al.* 2014).

In patients with CPP due to *MKRN3* defects, the median age of puberty onset was 6.0 years in girls (ranging from 3.0 to 7.5 years) and 8.25 years in boys (ranging from 5.9 to 9.0 years) (Abreu *et al.* 2013, Macedo *et al.* 2014, Schreiner *et al.* 2014, Settas *et al.* 2014, de Vries *et al.* 2014), suggesting that the *MKRN3* mutations may affect girls more severely than boys (Table 1). Given the median age of pubertal onset of affected patients with *MKRN3* mutations, it is speculated that the prepubertal inhibitory tonus on GnRH secretion took place normally, but was lost prematurely in patients with *MKRN3* mutations. This clinical observation suggests that *MKRN3* may not be crucial for GnRH suppression after the mini-puberty of early infancy, but that the downregulation of *MKRN3* plays a relevant role for the re-emergence of GnRH pulses in the pubertal onset (Macedo *et al.* 2014).

MKRN3 mutations and polymorphisms

Currently, 12 distinct loss-of-function mutations of *MKRN3* have been described in 30 patients (22 girls and eight boys) with CPP from 17 families from different ethnicities (Fig. 1 and Table 1). Remarkably, eight of these mutations encode either premature stop codons or frameshift mutations (Table 1). The four missense mutations (p.Cys340Gly, p.Arg365Ser, p.Phe417Ile, and p.His420Gln) were located within a zinc finger motif or a RING finger motif (Fig. 1), regions predicted to be involved in RNA binding and ubiquitin ligase activity of the protein respectively, and essential for protein function. All missense mutations are predicted to be pathogenic by *in silico* analysis. Additionally, *ab initio* modeling of the mutations p.Arg365Ser and p.Cys340Gly predicts that these mutations lead to significant structural perturbations in the 3D structure of the RING finger motif (Settas *et al.* 2014). The substitution of histidine 420 with glutamine in the *MKRN3* protein is predicted to reduce the affinity between the Zn ion-binding site and the relevant Zn, disrupting the binding pocket leading to unfolding of the finger (de Vries *et al.* 2014). Interestingly, most of the *MKRN3* mutations (64%) were located in the amino-terminal region of the protein, which is encoded by a poly-C-rich sequence, suggesting that this area may be a potential hotspot.

The important role of *MKRN3* in human puberty initiation has been reinforced recently by large genome-wide and custom-genotyping arrays in up to 182 416 women of European descent (57 studies) (Perry *et al.* 2014).

There was evidence of 123 signals at 106 genomic *loci* associated with age at menarche. Three of these *loci* were located in imprinted regions, including the *MKRN3* locus, demonstrating parent-of-origin-specific associations concordant with known parental expression patterns (Perry *et al.* 2014). This study suggests that not only are rare variants in *MKRN3* associated with CPP, but also more common variants/polymorphisms may be associated with changes in the timing of puberty (as reflected by the age of menarche) within the normal range and within the general population. This was a remarkable extension of the findings of rare variants in a disease such as CPP to more common variants in a polygenic trait such as age of menarche. The lack of parent-of-origin-specific analysis in previous GWAS studies may explain why *MKRN3* has not been previously associated with the age of menarche (He *et al.* 2009, Elks *et al.* 2010).

MKRN3 gene and protein structure and expression

MKRN3 was first cloned in 1999 by Jong *et al.* (1999a) during a study of the Prader–Willi/Angelman syndrome (PW/AS) critical region. They identified a cDNA in the PW/AS region encoding a zinc finger protein, initially named zinc finger protein 127 (*ZNF127*) and later renamed *MKRN3*. The functional and physiological relevance of *MKRN3* is not known and despite its location in the PWS critical region, its role in this syndrome is also unclear. An antisense transcript was concomitantly identified and named *ZNF127AS* (*MKRN3-AS1*). The antisense gene is not translated and is expressed weakly during fetal development and at very low levels in adult brain regions (Jong *et al.* 1999a). Similar to other antisense genes, *ZNF127AS* may regulate the expression of the 'sense' gene (*MKRN3*).

MKRN3 is located on human chromosome 15q11–13 (chromosome 7C in mouse) in a region that contains a cluster of imprinted genes associated with two different neurobehavioral disorders (Lalande 1996, Nicholls *et al.* 1998). Mutations or loss of expression of the maternally expressed gene *UBE3A* lead to Angelman syndrome (AS), while PWS is thought to be a contiguous gene syndrome arising from the loss of expression of multiple paternally expressed genes, including *MKRN3*. The report of two patients with PWS with a deletion of the 15q11–13 locus that did not include *MKRN3* suggested that this gene is not required for the development of the syndrome, but we still cannot rule out a role for *MKRN3* in some of the clinical features of PWS (Kanber *et al.* 2009).

MKRN3 differential allele expression occurs through silencing of the *MKRN3* maternal allele, which is associated with 5' CpG island methylation (Hershko *et al.* 1999, Jong *et al.* 1999b). Methylation of *MKRN3* and the other imprinted genes located within the PW/AS region of chromosome 15 is coordinately regulated by a bipartite imprinting control center (IC), composed of a sequence around the *SNRPN* promoter, the PWS-IC, and an 880 bp sequence located 35 kb upstream, the AS-IC (Rabinovitz *et al.* 2012). The molecular mechanisms of IC function are not well understood, but it has been postulated, based on studies in humans and mouse models, that the PWS-IC is a positive regulatory element and activates paternal-specific gene expression (Bielinska *et al.* 2000, Bressler *et al.* 2001), whereas the AS-IC functions in the maternal imprint by allele-specific repression of the PWS-IC to prevent the paternal imprinting program (Jong *et al.* 1999b, Ohta *et al.* 1999, Shemer *et al.* 2000, Rodriguez-Jato *et al.* 2005).

MKRN3 is an intronless gene, and probably arose by germline retrotransposition from the Makorin ring finger protein 1 gene (*MKRN1*), a highly transcribed, intron-containing gene, 80–90 million years ago (Rapkins *et al.* 2006). *MKRN1* is the ancestral founder of the Makorin gene family (Gray *et al.* 2000). Several other retrocopies of *MKRN1* have been identified in mammalian genomes, most of them probably corresponding to pseudogenes. Nine *MKRN* family *loci* distributed throughout the human genome have been identified thus far, with only three functional genes – *MKRN1*, *MKRN2*, and *MKRN3* – identified in vertebrates (Gray *et al.* 2000, Bohne *et al.* 2010).

The MAKORIN family

The *MKRN* gene family encodes putative ribonucleoproteins with a distinctive array of zinc-finger motifs including several C3H zinc fingers, a makorin-specific Cys–His arrangement, and a RING zinc finger (Jong *et al.* 1999a,b). In particular, *MKRN3* has a centrally located RING finger motif (C3HC4), two amino-terminal C3H zinc finger motifs followed by the unique pattern of conserved Cys–His residues called a Makorin zinc finger motif, and a carboxy-terminal C3H zinc finger motif (Jong *et al.* 1999a; Fig. 1). *MKRN3* is highly conserved among species, and the mouse and human *MKRN3* amino acid sequences share 69% identity and 82% similarity (Jong *et al.* 1999a). Mice and humans usually do not have conserved UTRs, yet the *MKRN3* 3'-UTR has 90% identity between these two species, implying a functional significance to this region of *MKRN3*. *MKRN3* is ubiquitously expressed in

adult human tissue, with the highest expression levels in the testis. In the fetus, it is highly expressed in the CNS and is expressed in post-meiotic sperm germ cells, particularly in round spermatids (Jong *et al.* 1999a).

The characteristic arrangement of cysteine (Cys or C) and histidine (His or H) residues in the zinc finger proteins in MKRN3 can allow some predictions of its function. C3H zinc fingers have been identified in RNA-binding proteins (Barabino *et al.* 1997, Murray *et al.* 1997). RING zinc fingers have been shown to mediate protein:protein interactions (Schwabe & Klug 1994). More recent evidence suggests that the RING zinc finger is a signature domain for E3 ligases, a category of enzymes mediating the transfer of ubiquitin from an E2 ubiquitin-conjugating enzyme to target protein substrates (Deshaies & Joazeiro 2009). Ubiquitination can have diverse effects on the protein substrate, varying from proteasome-dependent proteolysis to modulation of protein function and/or localization (Deshaies & Joazeiro 2009, Behrends & Harper 2011). The multiple types and number of zinc finger motifs in makorin proteins suggest possible multiple cellular functions for this protein.

Makorin proteins share a highly homologous amino acid sequence, particularly in the zinc finger domains (Fig. 2), suggesting that they may share similar functions or regulatory mechanisms. Although there are not many studies on MKRN3 function to date, MKRN1 function has been better explored. A possible important cellular function for MKRN1 is supported by the high identity (92%) between the human and murine orthologs and

the ubiquitous protein expression in human and mouse tissues (Gray *et al.* 2000). MKRN1 acts as an E3 ubiquitin ligase, inducing degradation of human telomerase reverse transcriptase (hTERT), viral capsid proteins, p53 and p21 cell cycle regulators, and peroxisome proliferator-activated receptor gamma (PPAR γ) (Kim *et al.* 2005, 2014, Lee *et al.* 2009).

An alternative mechanism of action for MKRN1 was described when it was identified as a repressor of c-Jun transcriptional activity using a yeast assay. Through regulation of RNA polymerase II-dependent transcription, MKRN1 can have either negative or positive effects on gene expression (Omwancha *et al.* 2006). Interestingly, it has been demonstrated that the disruption of ubiquitin ligase activity of MKRN1 did not affect its inhibitory transcriptional activity, suggesting that this function is independent of protein ubiquitination (Omwancha *et al.* 2006).

MKRN2 is also ubiquitously expressed (Gray *et al.* 2000). Similar to other members of this family, MKRN2 is highly conserved throughout evolution. It has been shown that mkrn2 negatively regulates neurogenesis via PI3K/Akt signaling in *Xenopus* embryos; however, the detailed molecular mechanisms of this effect and the potential functions of mammalian MKRN2 remain yet to be studied (Yang *et al.* 2008).

MKRN3 presumptively possesses ubiquitin-protein isopeptide ligase (E3) activity, intimated by the presence of the highly conserved C3HC4 RING finger domain. The tandem repeat of C3H zinc fingers may provide

Figure 2

MKRN1 and MKRN3 protein sequence alignment. The two protein sequences share high homology and similarity, especially in the RING finger domains. Bold letters represent conserved amino acids, and squares similar amino acids. Sections highlighted in red represent the three zinc fingers

C3H motifs, in yellow the MKRN-specific Cys–His motif, and in green the RING finger domain. MKRN1 NCBI Reference Sequence NP_038474, UniProtKB/Swiss-Prot: Q9UHC7.3. MKRN3 NP_005655.1 and UniProtKB/Swiss-Prot Q13064.1.

high-specificity RNA binding (Hudson *et al.* 2004), and the unique Cys–His makorin motif has been suggested to be a DNA-binding domain. Taken together, the studies of MKRN1 and MKRN2 imply that MKRN3 can similarly act as an E3 ligase, based on the high homology of the RING finger domain (Fig. 2). A previous study on mice demonstrated that *Mkrm3* is highly expressed in the hypothalamic arcuate nucleus during the infantile and early juvenile periods, with a reduction in expression at postnatal days 12–15, before puberty initiation (Abreu *et al.* 2013). The arcuate nucleus is the site of expression of critical regulators of GnRH secretion, such as kisspeptin, neurokinin B, and dynorphin (Navarro *et al.* 2011). Puberty is initiated with a loss of inhibitory inputs and a gain in excitatory inputs (Ojeda & Lomniczi 2014). The *Mkrm3* expression pattern in the hypothalamic arcuate nucleus suggests that this peptide plays a role in the inhibition of GnRH secretion during the prepubertal quiescent period (Fig. 3). The decrease in *Mkrm3* expression is hypothesized to be associated with an increase in GnRH stimulatory factors and/or GnRH expression, and it can be postulated that MKRN3 may be acting at the hypothalamic level as an E3 ligase to inhibit stimulatory inputs, so that loss-of-function mutations of MKRN3 result in early activation of the hypothalamic–pituitary–gonadal axis, expressed phenotypically as CPP. It is also possible that MKRN3 can act as a transcriptional regulator, as has been demonstrated for MKRN1. Data from the Human Protein Atlas indicate that MKRN3 is located primarily in the plasma membrane and cytoplasm, and also in the nucleus (www.proteinatlas.org). Based on its location in

the plasma membrane, we can speculate that MKRN3 may also be involved in endocytosis and downregulation of receptors, as has been demonstrated for some other E3 ligases (Hershko & Ciechanover 1998). The genetic findings from patients with CPP are in agreement with the hypothesis that MKRN3 may act as a ‘brake’ or inhibitor of GnRH secretion during childhood (Hughes 2013). Further studies are needed to elucidate the precise mechanism(s) of action of MKRN3.

Conclusion

Age of puberty initiation is associated with causes of substantial morbidity and mortality. Early age of menarche is associated with breast cancer and cardiovascular disease (Kvale 1992, Lakshman *et al.* 2009). Therefore, it is important to understand the mechanisms controlling puberty initiation. In an attempt to identify genes that will broaden the knowledge on GnRH regulation to bring new genetic screening, and diagnostic and treatment tools, researchers have been trying to identify genes associated with CPP for several years. By studying individuals with hypogonadotropic hypogonadism, a genetic disorder caused by absence or deficiency of GnRH secretion, we learned about important stimulators of GnRH secretion (Bianco & Kaiser 2009, Semple & Topaloglu 2010). Owing to the complexity of the mechanisms involved in puberty initiation and the initial failure to identify genes associated with CPP, it has been hypothesized that CPP was not caused by mutations in a single gene, but rather was a consequence of complex interactions among environment, metabolic factors, and polygenic defects. However, with the development of new sequencing methodologies, the combination of whole-exome sequencing analysis with detailed phenotypic characterization and the careful selection of the correct cohort of patients led to the identification of *MKRN3* as an important regulator of pubertal development. *MKRN3* is the first gene with a probable inhibitory effect on GnRH secretion, with mutations identified within humans. To date, *MKRN3* is the most common genetic defect associated with CPP. The identification of loss-of-function mutations in this gene can contribute to the diagnosis of CPP, especially in boys in whom the signs of puberty initiation are not easily detectable, thereby helping to make the diagnosis earlier and facilitating treatment decisions. In addition, the presence of *MKRN3* mutations can contribute to early diagnosis of CPP in familial cases and guide genetic counseling. It is not clear if there is a sex difference in the effect of *MKRN3* mutations, but it

Figure 3

Schematic representation of MKRN3 mechanism of action. Human and mouse studies suggest that MKRN3 acts as an inhibitor of GnRH secretion during childhood (diagram on A), and that a decrease in *MKRN3* expression is associated with an increase in GnRH stimulatory factors and/or GnRH resulting in puberty initiation (diagram on B). NKB, neurokinin B; – inhibition, + stimulation. ↑ increase ↓ decrease.

is evident that mutations in this gene accelerate puberty initiation in both sexes. Although the precise mechanism of regulation of GnRH secretion by MKRN3 is not yet understood, its importance in the hypothalamic–pituitary–gonadal axis is indisputable. The studies of *Mkrm3* expression in the hypothalamic arcuate nucleus of mice support the findings that loss-of-function mutations in humans lead to early puberty initiation and strengthen the hypothesis that MKRN3 acts as an inhibitor of GnRH secretion during childhood (Fig. 3). Ultimately, the recent GWAS findings linking MKRN3 with the age of menarche have consolidated the involvement of this gene in pubertal timing.

The identification of a putative inhibitory factor in the hypothalamic–pituitary–gonadal axis has opened an exciting new arena in the neuroendocrine field. Further studies will elucidate the precise mechanism of action of this important regulator of GnRH secretion.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was supported by FAPESP grant numbers 2013-06391-1 to DBM and 2013-03236-5 to ACL; Eunice Kennedy Shriver NICHD/NIH through cooperative agreement U54HD28138 as part of the Specialized Cooperative Centers Program in Reproduction and Infertility Research and Eunice Kennedy Shriver NICHD/NIH R01HD019938, to UK; and 1F05HD072773-03, to APA.

References

- Abreu AP, Dauber A, Macedo DB, Noel SD, Brito VN, Gill JC, Cukier P, Thompson IR, Navarro VM, Gagliardi PC *et al.* 2013 Central precocious puberty caused by mutations in the imprinted gene MKRN3. *New England Journal of Medicine* **368** 2467–2475. (doi:10.1056/NEJMoa1302160)
- Barabino SM, Hubner W, Jenny A, Minvielle-Sebastia L & Keller W 1997 The 30-kD subunit of mammalian cleavage and polyadenylation specificity factor and its yeast homolog are RNA-binding zinc finger proteins. *Genes and Development* **11** 1703–1716. (doi:10.1101/gad.11.13.1703)
- Behrends C & Harper JW 2011 Constructing and decoding unconventional ubiquitin chains. *Nature Structural & Molecular Biology* **18** 520–528. (doi:10.1038/nsmb.2066)
- Bianco SD & Kaiser UB 2009 The genetic and molecular basis of idiopathic hypogonadotropic hypogonadism. *Nature Reviews. Endocrinology* **5** 569–576. (doi:10.1038/nrendo.2009.177)
- Bielinska B, Blaydes SM, Buiting K, Yang T, Krajewska-Walasek M, Horsthemke B & Brannan CI 2000 *De novo* deletions of SNRPN exon 1 in early human and mouse embryos result in a paternal to maternal imprint switch. *Nature Genetics* **25** 74–78. (doi:10.1038/75629)
- Bohne A, Darras A, D'Cotta H, Baroiller JF, Galiana-Arnoux D & Volff JN 2010 The vertebrate makorin ubiquitin ligase gene family has been shaped by large-scale duplication and retroposition from an ancestral gonad-specific, maternal-effect gene. *BMC Genomics* **11** 721. (doi:10.1186/1471-2164-11-721)
- Bressler J, Tsai TF, Wu MY, Tsai SF, Ramirez MA, Armstrong D & Beaudet AL 2001 The SNRPN promoter is not required for genomic imprinting of the Prader–Willi/Angelman domain in mice. *Nature Genetics* **28** 232–240. (doi:10.1038/90067)
- Deshais RJ & Joazeiro CA 2009 RING domain E3 ubiquitin ligases. *Annual Review of Biochemistry* **78** 399–434. (doi:10.1146/annurev.biochem.78.101807.093809)
- Elks CE, Perry JR, Sulem P, Chasman DI, Franceschini N, He C, Lunetta KL, Visser JA, Byrne EM, Cousminer DL *et al.* 2010 Thirty new loci for age at menarche identified by a meta-analysis of genome-wide association studies. *Nature Genetics* **42** 1077–1085. (doi:10.1038/ng.714)
- Gray TA, Hernandez L, Carey AH, Schaldach MA, Smithwick MJ, Rus K, Marshall Graves JA, Stewart CL & Nicholls RD 2000 The ancient source of a distinct gene family encoding proteins featuring RING and C(3)H zinc-finger motifs with abundant expression in developing brain and nervous system. *Genomics* **66** 76–86. (doi:10.1006/geno.2000.6199)
- He C, Kraft P, Chen C, Buring JE, Pare G, Hankinson SE, Chanock SJ, Ridker PM, Hunter DJ & Chasman DI 2009 Genome-wide association studies identify loci associated with age at menarche and age at natural menopause. *Nature Genetics* **41** 724–728. (doi:10.1038/ng.385)
- Herman-Giddens ME, Slora EJ, Wasserman RC, Bourdony CJ, Bhapkar MV, Koch GG & Hasemeier CM 1997 Secondary sexual characteristics and menses in young girls seen in office practice: a study from the Pediatric Research in Office Settings network. *Pediatrics* **99** 505–512. (doi:10.1542/peds.99.4.505)
- Hershko A & Ciechanover A 1998 The ubiquitin system. *Annual Review of Biochemistry* **67** 425–479. (doi:10.1146/annurev.biochem.67.1.425)
- Hershko A, Razin A & Shemer R 1999 Imprinted methylation and its effect on expression of the mouse Zfp127 gene. *Gene* **234** 323–327. (doi:10.1016/S0378-1119(99)00192-4)
- Hudson BP, Martinez-Yamout MA, Dyson HJ & Wright PE 2004 Recognition of the mRNA AU-rich element by the zinc finger domain of TIS11d. *Nature Structural & Molecular Biology* **11** 257–264. (doi:10.1038/nsmb738)
- Hughes IA 2013 Releasing the brake on puberty. *New England Journal of Medicine* **368** 2513–2515. (doi:10.1056/NEJMe1306743)
- Jong MT, Gray TA, Ji Y, Glenn CC, Saitoh S, Driscoll DJ & Nicholls RD 1999a A novel imprinted gene, encoding a RING zinc-finger protein, and overlapping antisense transcript in the Prader–Willi syndrome critical region. *Human Molecular Genetics* **8** 783–793. (doi:10.1093/hmg/8.5.783)
- Jong MT, Carey AH, Caldwell KA, Lau MH, Handel MA, Driscoll DJ, Stewart CL, Rinchik EM & Nicholls RD 1999b Imprinting of a RING zinc-finger encoding gene in the mouse chromosome region homologous to the Prader–Willi syndrome genetic region. *Human Molecular Genetics* **8** 795–803. (doi:10.1093/hmg/8.5.795)
- Kanber D, Giltay J, Wieczorek D, Zogel C, Hochstenbach R, Caliebe A, Kuechler A, Horsthemke B & Buiting K 2009 A paternal deletion of MKRN3, MAGEL2 and NDN does not result in Prader–Willi syndrome. *European Journal of Human Genetics* **17** 582–590. (doi:10.1038/ejhg.2008.232)
- Kim JH, Park SM, Kang MR, Oh SY, Lee TH, Muller MT & Chung IK 2005 Ubiquitin ligase MKRN1 modulates telomere length homeostasis through a proteolysis of hTERT. *Genes and Development* **19** 776–781. (doi:10.1101/gad.1289405)
- Kim JH, Park KW, Lee EW, Jang WS, Seo J, Shin S, Hwang KA & Song J 2014 Suppression of PPAR γ through MKRN1-mediated ubiquitination and degradation prevents adipocyte differentiation. *Cell Death and Differentiation* **21** 594–603. (doi:10.1038/cdd.2013.181)
- Kvale G 1992 Reproductive factors in breast cancer epidemiology. *Acta Oncologica* **31** 187–194. (doi:10.3109/02841869209088901)

- Lakshman R, Forouhi NG, Sharp SJ, Luben R, Bingham SA, Khaw KT, Wareham NJ & Ong KK 2009 Early age at menarche associated with cardiovascular disease and mortality. *Journal of Clinical Endocrinology and Metabolism* **94** 4953–4960. (doi:10.1210/jc.2009-1789)
- Lalande M 1996 Parental imprinting and human disease. *Annual Review of Genetics* **30** 173–195. (doi:10.1146/annurev.genet.30.1.173)
- Lee EW, Lee MS, Camus S, Ghim J, Yang MR, Oh W, Ha NC, Lane DP & Song J 2009 Differential regulation of p53 and p21 by MKRN1 E3 ligase controls cell cycle arrest and apoptosis. *EMBO Journal* **28** 2100–2113. (doi:10.1038/emboj.2009.164)
- Macedo DB, Abreu AP, Reis AC, Montenegro LR, Dauber A, Beneduzzi D, Cukier P, Silveira LF, Teles MG, Carroll RS *et al.* 2014 Central precocious puberty that appears to be sporadic caused by paternally inherited mutations in the imprinted gene makorin ring finger 3. *Journal of Clinical Endocrinology and Metabolism* **99** E1097–E1103. (doi:10.1210/jc.2013-3126)
- Murray MV, Turnage MA, Williamson KJ, Steinhauer WR & Searles LL 1997 The *Drosophila* suppressor of sable protein binds to RNA and associates with a subset of polytene chromosome bands. *Molecular and Cellular Biology* **17** 2291–2300.
- Navarro VM, Gottsch ML, Wu M, Garcia-Galiano D, Hobbs SJ, Bosch MA, Pinilla L, Clifton DK, Dearth A, Ronnekleiv OK *et al.* 2011 Regulation of NKB pathways and their roles in the control of Kiss1 neurons in the arcuate nucleus of the male mouse. *Endocrinology* **152** 4265–4275. (doi:10.1210/en.2011-1143)
- Nicholls RD, Saitoh S & Horsthemke B 1998 Imprinting in Prader–Willi and Angelman syndromes. *Trends in Genetics* **14** 194–200. (doi:10.1016/S0168-9525(98)01432-2)
- Ohta T, Buiting K, Kokkonen H, McCandless S, Heeger S, Leisti H, Driscoll DJ, Cassidy SB, Horsthemke B & Nicholls RD 1999 Molecular mechanism of angelman syndrome in two large families involves an imprinting mutation. *American Journal of Human Genetics* **64** 385–396. (doi:10.1086/302232)
- Ojeda SR & Lomniczi A 2014 Puberty in 2013: unravelling the mystery of puberty. *Nature Reviews. Endocrinology* **10** 67–69. (doi:10.1038/nrendo.2013.233)
- Omwancha J, Zhou XF, Chen SY, Baslan T, Fisher CJ, Zheng Z, Cai C & Shemshedini L 2006 Makorin RING finger protein 1 (MKRN1) has negative and positive effects on RNA polymerase II-dependent transcription. *Endocrine* **29** 363–373. (doi:10.1385/ENDO:29:2:363)
- Perry JR, Day F, Elks CE, Sulem P, Thompson DJ, Ferreira T, He C, Chasman DI, Esko T, Thorleifsson G *et al.* 2014 Parent-of-origin-specific allelic associations among 106 genomic loci for age at menarche. *Nature* **514** 92–97. (doi:10.1038/nature13545)
- Rabinovitz S, Kaufman Y, Ludwig G, Razin A & Shemer R 2012 Mechanisms of activation of the paternally expressed genes by the Prader–Willi imprinting center in the Prader–Willi/Angelman syndromes domains. *PNAS* **109** 7403–7408. (doi:10.1073/pnas.1116661109)
- Rapkins RW, Hore T, Smithwick M, Ager E, Pask AJ, Renfree MB, Kohn M, Hameister H, Nicholls RD, Deakin JE *et al.* 2006 Recent assembly of an imprinted domain from non-imprinted components. *PLoS Genetics* **2** e182. (doi:10.1371/journal.pgen.0020182)
- Rodriguez-Jato S, Nicholls RD, Driscoll DJ & Yang TP 2005 Characterization of cis- and trans-acting elements in the imprinted human SNURF-SNRPN locus. *Nucleic Acids Research* **33** 4740–4753. (doi:10.1093/nar/gki786)
- Schreiner F, Gohlke B, Hamm M, Korsch E & Woelfle J 2014 MKRN3 mutations in familial central precocious puberty. *Hormone Research in Paediatrics* **82** 122–126. (doi:10.1159/000362815)
- Schwabe JW & Klug A 1994 Zinc mining for protein domains. *Nature Structural Biology* **1** 345–349. (doi:10.1038/nsb0694-345)
- Sample RK & Topaloglu AK 2010 The recent genetics of hypogonadotrophic hypogonadism – novel insights and new questions. *Clinical Endocrinology* **72** 427–435. (doi:10.1111/j.1365-2265.2009.03687.x)
- Settas N, Dacou-Voutetakis C, Karantzis M, Kanaka-Gantenbein C, Chrousos GP & Voutetakis A 2014 Central precocious puberty in a girl and early puberty in her brother caused by a novel mutation in the MKRN3 gene. *Journal of Clinical Endocrinology and Metabolism* **99** E647–E651. (doi:10.1210/jc.2013-4084)
- Shemer R, Hershko AY, Perk J, Mostoslavsky R, Tsuberi B, Cedar H, Buiting K & Razin A 2000 The imprinting box of the Prader–Willi/Angelman syndrome domain. *Nature Genetics* **26** 440–443. (doi:10.1038/82571)
- Silveira LG, Noel SD, Silveira-Neto AP, Abreu AP, Brito VN, Santos MG, Bianco SD, Kuohung W, Xu S, Gryngarten M *et al.* 2010 Mutations of the KISS1 gene in disorders of puberty. *Journal of Clinical Endocrinology and Metabolism* **95** 2276–2280. (doi:10.1210/jc.2009-2421)
- Teles MG, Bianco SD, Brito VN, Trarbach EB, Kuohung W, Xu S, Seminara SB, Mendonca BB, Kaiser UB & Latronico AC 2008 A GPR54-activating mutation in a patient with central precocious puberty. *New England Journal of Medicine* **358** 709–715. (doi:10.1056/NEJMoa073443)
- de Vries L, Gat-Yablonski G, Dror N, Singer A & Phillip M 2014 A novel MKRN3 missense mutation causing familial precocious puberty. *Human Reproduction* **29** 2838–2843. (doi:10.1093/humrep/deu256)
- Yang PH, Cheung WK, Peng Y, He ML, Wu GQ, Xie D, Jiang BH, Huang QH, Chen Z, Lin MC *et al.* 2008 Makorin-2 is a neurogenesis inhibitor downstream of phosphatidylinositol 3-kinase/Akt (PI3K/Akt) signal. *Journal of Biological Chemistry* **283** 8486–8495. (doi:10.1074/jbc.M704768200)
- Zacharias L & Wurtman RJ 1969 Age at menarche. Genetic and environmental influences. *New England Journal of Medicine* **280** 868–875. (doi:10.1056/NEJM196904172801606)

Received in final form 10 February 2015

Accepted 6 March 2015