Reversal of fortune: estrogen receptor- β in endometriosis

Rosalia C M Simmen¹ and Angela S Kelley²

¹Department of Physiology and Biophysics, University of Arkansas for Medical Sciences, Little Rock, Arkansas, USA

²Department of Obstetrics and Gynecology, University of Michigan Health System, Ann Arbor, Michigan, USA

Correspondence should be addressed to R C M Simmen
Email
simmenrosalia@uams.edu

Abstract

Enhanced inflammation and reduced apoptosis sustain the growth of endometriotic lesions. Alterations in the expression of estrogen receptor-alpha (ER α) and estrogen receptor-beta (ER β) accompany the conversion of resident endometrial cells within the normal uterine environment to ectopic lesions located in extrauterine sites. Recent studies highlighted in this focused review linked ER β to dysregulation of apoptotic and inflammatory networks involving novel interacting partners in endometriosis. The elucidation of these nongenomic actions of ER β using human cells and mouse models is an important step in understanding key regulatory pathways that are disrupted leading to disease establishment and progression.

Key Words

- ▶ endometriosis
- ▶ estrogen receptor-beta
- ▶ inflammation
- ▶ non-genomic
- ▶ apoptosome

Journal of Molecular Endocrinology
(2016) **57**, F23–F27

Estrogens are key mediators of endometrial homeostasis; hence, any dysregulation in their synthesis, metabolism, and/or activities irrefutably leads to a broad range of endometrial pathologies. Two major estrogen receptor proteins, estrogen receptor-alpha (ER α) and estrogen receptor-beta (ER β) which are encoded by distinct genes (White *et al.* 1987, Kuiper *et al.* 1996), mediate the actions of estrogens in target cells. The significant homologies in DNA (94%) and ligand-binding domains (59%) between these ERs enable both proteins to bind estrogens with equal affinity and to transcriptionally regulate common subsets of ER-responsive genes. ER α and ER β display distinct spatial, temporal, and physiological expression. Genetic deletions support most roles for ER α in the uterus and for ER β to prevail in the ovary (Hamilton *et al.* 2014). Although ligand-bound ER α is a requisite for mitogenesis of uterine cells, ER β is considered to inhibit ER α -dependent cell proliferation in part, through its ability to form ER α /ER β heterodimers with different ligand specificity, interacting partners,

and transcriptional targets than ER α homodimers (Pace *et al.* 1997).

Endometriosis is an estrogen-dependent gynecological disorder, defined as the growth of endometrial stroma and glands in extrauterine sites such as in the peritoneum and the ovary (Burney & Giudice 2012). The disease affects 1 in 10 reproductive age women and can lead to debilitating pelvic pain, dysmenorrhea, and reduced fertility. Although benign, endometriosis is a chronic condition that requires long-term treatment throughout a woman's reproductive years. The goals of treatment for endometriosis include hormonal suppression of active endometriosis or surgical excision/ablation of visible lesions (Vercellini *et al.* 2014). However, medical therapies are not without side effects and surgical removal of the ovaries and uterus, although considered a definitive management for the condition, is not without associated morbidity.

In recent years, ER β has emerged as an important player in the pathogenesis of endometriosis. Human endometriotic lesions, whether of ovarian or peritoneal

locations, display higher ER β expression compared with normal human endometrial cells; this was not shown for ER α (Bulun *et al.* 2012). The reversal in the ER β -to-ER α ratio in lesions ($>>1$), relative to normal endometrial cells ($<<1$), has been similarly demonstrated in many animal models of endometriosis (Fazleabas *et al.* 2003, Greaves *et al.* 2014, Heard *et al.* 2014). It raises the intriguing question of how the conventionally antiproliferative ER β in the endometrium, in the context of endogenous estrogens, dismantles its customary role and orchestrates a pro-proliferative/antiapoptotic/proinflammatory response to drive disease pathogenesis. In a recent issue of *Cell* (Han *et al.* 2015), an elegant study by O'Malley and coworkers unequivocally demonstrated that ER β is required for the progression of endometriosis in mice and defined the apoptosome and inflammasome as endogenous targets of nongenomic ER β action. Katzenellenbogen and colleagues in a related study published in *Science Translational Medicine* (Zhao *et al.* 2015) described two ER antagonists, one specific for ER α (chloroindazole (CLI)) and the other specific for ER β (oxabicycloheptane sulfonate (OBHS)), that individually function to inhibit the estrogen-inflammatory axis to suppress endometriosis in mice. These two studies collectively define a novel role for ER β in the upper echelon of the inflammatory regulatory hierarchy. Importantly, as inflammation is well considered as one of the major contributors to the development and progression of endometriosis, the study findings offer promise for novel therapeutic strategies that may be relevant to endometriosis and other reproductive and nonreproductive diseases associated with chronic inflammation.

To uncover the underlying mechanism(s) by which ER β promotes endometriosis, O'Malley and his group used ovariectomized+E₂-pelleted immunocompetent mouse models in which endometriosis was surgically induced through autotransplantation of ER β -overexpressing (ER β -OE) (gain-of-ER β -function) and ER β -null (loss-of-ER β -function) endometrial tissues in the peritoneal cavity. Ectopic lesions generated from ER β -OE endometrium had larger volumes, whereas those from ER β -null endometrium had smaller volumes, than did wild-type (WT; control) ectopic lesions. Consistent with these observations, the use of the ER β -selective antagonist PHTPP (a pyrazolo[1,5- α] pyrimidine-based ligand) suppressed ectopic lesion growth relative to vehicle alone. The inhibitory effect of PHTPP on lesion growth was accompanied by the loss of recruitment of CD163-positive monocyte/macrophage cells that normally infiltrate lesions. Conversely, the

ER β -specific agonist ERB-041 enhanced the growth of mouse ectopic lesions compared with vehicle alone. Because these noted changes in lesion growth occurred in the absence of perturbations in ER α , enhanced ER β expression and activity appear sufficient to promote endometriosis progression. In an earlier study (Harris *et al.* 2005), athymic nude mice surgically implanted with human endometriotic lesions showed lesion regression when administered ERB-041, a response clearly contrasting with that obtained with the immunocompetent mouse model in the *Cell* study. Intriguingly, ectopic lesions formed in control and ERB-041-treated nude mice did not express ER β , a major departure from lesions of women and those generated in other animal models of endometriosis, in which ER β is the predominant ER isoform. Although the molecular basis underlying the differential responses of the two mouse models to ERB-041 remains unknown, these results provide support for the complex interactions between the immune system and ER-mediated signaling in the development and progression of endometriosis. Additionally, these results highlight innate limitations of the animal models utilized for many endometriosis studies, including those described in the two highlighted papers in this review (e.g., the need to implant estrogen pellets, which most likely does not reflect/only approximates the situation in humans) and underscore the continuing need for the development of more relevant models to fully understand the human disease.

How might ER β alter lesion biology distinct from its role in nondiseased endometrial cells? The authors analyzed by mass spectroscopy flag-tagged ER β -containing protein complexes that were immunoprecipitated from eutopic endometria of endometrial ER β -OE mice. Further confirmation by Western blotting revealed that a majority of proteins interacting with ER β are involved in inflammation and apoptosis signaling. Apoptosis plays an important role in inflammation and in the resolution of inflammatory reactions, and the two are irrefutably linked because cell death signaling initiated by tumor necrosis factor- α (TNF- α) activates inflammasomes to initiate IL-1 β -driven inflammation (Vince & Silke 2016). An attractive candidate identified in the study as an ER β -interacting protein is apoptosis signal-regulating kinase-1 (ASK-1). ASK-1 is a component of TNF- α -induced apoptosis complex 1, whose formation is required for TNF- α -induced apoptosis. Serine/threonine kinase receptor-associated protein (STRAP) and 14-3-3 protein were also identified to interact with ER β in the same screen. Interestingly, STRAP and 14-3-3 proteins have been previously demonstrated to bind ASK-1 in a

tripartite complex (Jung *et al.* 2010). The formation of this complex interferes with the functional association between ASK-1 and the TNF receptor-associated factor 2 in response to TNF- α signaling, resulting in the inhibition of TNF- α -induced apoptosis. To validate this model, the authors demonstrated (i) lower status of phosphorylated ASK-1 (phospho-Thr845 ASK-1), without accompanying changes in total ASK-1 protein levels, indicating loss of ASK-1 activation and hence, function in ER β -OE ectopic lesions compared with WT ectopic lesions; (ii) conversely, increased levels of phosphorylated ASK-1 in ER β -null ectopic lesions compared with WT ectopic lesions; and (iii) lower mitochondrial cytochrome *c* levels in ER β -OE ectopic lesions, consistent with the disruption of TNF- α -induced ASK-1 activation that normally leads to caspase 1 (Cas1) activation. The link between ER β and Cas1 was further illuminated by demonstrating that ER β -OE ectopic lesions had lower Cas1 levels and lacked detectable interactions between Cas1 and apoptotic peptidase-interacting factor; by contrast, the latter interaction was easily detected in ER β -null ectopic lesions. Taken together, these novel findings identify TNF- α -induced apoptosis as a key regulatory pathway disrupted by cytoplasmic-based ER β to promote lesion survival. The cytoplasmic-localized ER β antiapoptotic action likely occurs in conjunction with nuclear-localized ER β transcriptional activation of gene targets such as the serum and glucocorticoid-regulated kinase, which, by phosphorylating the proapoptotic FOXO3, inhibits apoptosis (Monsivais *et al.* 2016). However, the relative contribution of nuclear vs cytoplasmic actions of ER β to lesion survival is unknown.

The intriguing idea that ER β multitasks outside of the nucleus was further demonstrated in other experiments from the same study. The authors identified caspase-1 and the NLR family pyrin domain-containing 3 as additional ER β -interacting proteins. These findings are significant in the context of the inflammatory process, given the requisite participation of caspase-1 and NLR in the processing of pro-IL-1 β to the mature bioactive IL-1 β , a key regulator of adhesion and proliferation in endometrial cells (Kao *et al.* 2011). In this regard, ER β -OE lesions had higher IL-1 β and cleaved caspase-1 levels than did control ectopic lesions; conversely, ER β -null lesions had lower levels of both components. Given that primary human endometriotic stromal cells also showed elevated levels of IL-1 β and increased antiapoptosis signaling when treated with TNF- α , the results suggest that the key ER β -mediated events elucidated in these mouse models of endometriosis are relevant to the human disease.

Interestingly, the authors found, using immortalized human endometrial epithelial cells expressing Myc-tagged human ER α genes, that TNF- α treatment of these cells did not elicit comparable antiapoptotic responses and an increase in IL-1 β expression as noted for the same human cells expressing ER β . More intriguingly, cotransfection of ER α and ER β in these cells resulted in ER α inhibition of ER β -mediated IL-1 β production, suggesting that ER α may assume a negative regulatory role in ER β -mediated promotion of inflammation. Whether this effect of ER α reflects its competitive displacement of ER β from its interacting proteins or its sequestering of ER β through formation of ER α /ER β heterodimers is unknown. This result, while baffling, sheds new light on the contrasting activities of ER β and ER α in endometriosis and must be reconciled with previous findings that ER α is an equally active player in the pathology of this disease in mice (Burns *et al.* 2012). In the study by Katzenellenbogen and colleagues (Zhao *et al.* 2015), specific ER α (CLI) and ER β (OBHS) antagonists exhibited ER-dependent anti-inflammatory activities in a mouse model of endometriosis and in human endometriotic stromal cells. In these models, CLI and OBHS equally inhibited estrogen-dependent processes, including cell proliferation, cyst formation, vascularization, cytokine production, macrophage infiltration and lesion growth. A previous study has reported the presence of both ER α and ER β in peritoneal fluid macrophages and shown that both ERs display higher expression in peritoneal fluid macrophages of women with endometriosis compared with women without endometriosis (Montagna *et al.* 2008). Interestingly, this same study showed that only ER α expression was positively correlated with increased proinflammatory cytokine levels in peritoneal fluids of women with endometriosis. ER β levels, while higher in peritoneal fluid macrophages of women with endometriosis, were correlated with the expression of proinflammatory cytokines, irrespective of disease status, suggesting a role for macrophage ER β in basal proinflammatory cytokine production and for macrophage ER α as a likely 'driver' of increased inflammation seen in endometriosis. As mouse ectopic lesions, similar to human lesions, display higher levels of ER β than ER α , one intriguing question raised by the Katzenellenbogen study pertains to the unexpected comparable levels of inhibition of estrogen-dependent responses with disruption of ER α signaling to that mediated by the more highly expressed ER β . Perhaps the significant contribution of recruited macrophages expressing higher ER α , and hence equally higher production of pro-inflammatory cytokines, may underlie

the favorable response of lesions to ER α antagonists. It is also tempting to speculate that the cellular locations of the respective actions of ER α and ER β may be at play, given that in the O'Malley's study, ER β disruption of inflammasome and apoptosome functions takes place outside of the nucleus, invoking nongenomic actions and involving novel interactions with proteins not previously identified to interact with ER β . If the latter is true, then a follow-up question is whether ER α interacts (or not) with some of the same proteins that were identified for ER β . Yet another question is whether ER β preferentially acts outside of the nucleus in endometriotic lesions and if this may underlie the reversal of its (generally good) fortune of being the better half to ER α in controlling mitogenesis. The answer is not likely going to be straightforward given other nuclear actions attributed to ER β in ectopic lesions, one of which is its transcriptional regulation of ER α in endometriotic stromal cells (Trukhacheva *et al.* 2009). Time (and more in-depth scrutiny) will tell.

The molecular details of complex mechanisms have gained much ground from careful dissection of context-dependent cross talk among seemingly unrelated molecules. For an enigmatic condition such as endometriosis, the identification of novel partners

elucidated here for ER β begs the question of whether progesterone receptors (whose expression and function are equally compromised in endometriosis) and other steroid receptor coactivators or corepressors may similarly assume new extranuclear (cytoplasmic) roles to sustain ectopic growth. A tip-off to this possibility comes from the recent discovery of a new 70 kDa (truncated) steroid receptor coactivator 1 (SRC-1) isoform, which, similar to ER β , was found to have little expression in normal endometrium, but displayed a significant expression in endometriotic lesions (Han *et al.* 2012). Moreover, the truncated SRC-1 protein, similar to ER β , was demonstrated to be essential in the initial stages of endometriosis establishment. In the current *Cell* paper, SRC-1 isoform was shown to form a complex with ER β and caspase-8, inhibiting the latter from interacting with its usual partner Fas-associated via death domain protein to generate apoptosis complex II that augments TNF- α -induced apoptosis. Whether the truncated SRC-1 only partners with ER β or exhibits a more expansive repertoire of interacting proteins to promote endometriosis remains to be explored.

The present studies provide fundamental insights into the adaptive functions of ER β in inflammation and apoptosis (Fig. 1). Fine-tuning our awareness of the

Figure 1

Proposed model for ER β regulation of inflammation and apoptosis in endometriotic lesions. Normal endometrium and ectopic lesions display distinct levels of ER α and ER β . The direct interactions of ER β with proteins associated with inflammation and apoptosis in ectopic lesions (Han *et al.* 2015) provide novel, nongenomic mechanisms for ER β -mediated pathogenesis of endometriosis. CLI and OHBS are, respectively, specific ER α and ER β antagonists shown to inhibit ER-mediated promotion of inflammation underlying endometriosis progression (Zhao *et al.* 2015). ASK-1, STRAP, 14-3-3, NLR, Cas1, and IL-1 β are defined in the text.

different networks orchestrated by steroid hormone receptors and their changing partners in normal and endometriotic cells may offer the much-needed therapeutic opportunities to address the development and progression of endometriosis.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was supported in part by the National Institutes of Health (RO1CA136493, UL1TR000039) and the Arkansas Biosciences Institute Intramural Grant Program.

References

- Bulun SE, Monsavaiz D, Pavone ME, Dyson M, Xue Q, Attar E, Tokunaga H & Su EJ 2012 Role of estrogen receptor- β in endometriosis. *Seminars in Reproductive Medicine* **30** 39–45. (doi:10.1055/s-0031-1299596)
- Burney RO & Giudice LC 2012 Pathogenesis and pathophysiology of endometriosis. *Fertility and Sterility* **98** 511–519. (doi:10.1016/j.fertnstert.2012.06.029)
- Burns KA, Rodriguez KF, Hewitt SC, Janardhan KS, Young SL & Korach KS 2012 Role of estrogen receptor signaling required for endometriosis-like lesion establishment in a mouse model. *Endocrinology* **153** 3960–3971. (doi:10.1210/en.2012-1294)
- Fazleabas AT, Brudney A, Chai D, Langoi D & Bulun SE 2003 Steroid receptor and aromatase expression in baboon endometriotic lesions. *Fertility and Sterility* **80** (Supplement 2) 820–827. (doi:10.1016/S0015-0282(03)00982-8)
- Greaves E, Cousins FL, Murray A, Esnal-Zufiaurre A, Fassbender A, Horne AW & Saunders PT 2014 A novel mouse model of endometriosis mimics human phenotype and reveals insights into the inflammatory contribution of shed endometrium. *American Journal of Pathology* **184** 1930–1939. (doi:10.1016/j.ajpath.2014.03.011)
- Hamilton KJ, Arao Y & Korach KS 2014 Estrogen hormone physiology: reproductive findings from estrogen receptor mutant mice. *Reproductive Biology* **14** 3–8. (doi:10.1016/j.repbio.2013.12.002)
- Han SJ, Hawkins SM, Begum K, Jung SY, Kovanci E, Qin J, Lydon JP, DeMayo FJ & O'Malley BW 2012 A new isoform of steroid receptor coactivator-1 is crucial for pathogenic progression of endometriosis. *Nature Medicine* **18** 1102–1111. (doi:10.1038/nm.2826)
- Han SJ, Jung SY, Wu S-P, Hawkins SM, Park MJ, Kyo S, Qin J, Lydon JP, Tsai SY, Tsai M-J, et al. 2015 Estrogen receptor β modulates apoptosis complexes and the inflammasome to drive the

- pathogenesis of endometriosis. *Cell* **163** 960–974. (doi:10.1016/j.cell.2015.10.034)
- Harris HA, Bruner-Tran KL, Zhang X, Osteen KG & Lyttle CRA 2005 Selective estrogen receptor-beta agonist causes lesion regression in an experimentally induced model of endometriosis. *Human Reproduction* **20** 936–941. (doi:10.1093/humrep/deh711)
- Heard ME, Simmons CD, Simmen FA & Simmen RC 2014 Krüppel-like factor 9 deficiency in uterine endometrial cells promotes ectopic lesion establishment associated with activated notch and hedgehog signaling in a mouse model of endometriosis. *Endocrinology* **155** 1532–1546. (doi:10.1210/en.2013-1947)
- Jung H, Seong HA, Manoharan R & Ha H 2010 Serine-threonine kinase receptor-associated protein inhibits apoptosis signal-regulating kinase 1 function through direct interaction. *Journal of Biological Chemistry* **285** 54–70. (doi:10.1074/jbc.M109.045229)
- Kao AP, Wang KH, Long CY, Chai CY, Tsai CF, Hsieh TH, Hsu CY, Chang CC, Lee JN & Tsai EM 2011 Interleukin-1 β induces cyclooxygenase-2 expression and promotes the invasive ability of human mesenchymal stem cells derived from ovarian endometrioma. *Fertility and Sterility* **96** 678–684. (doi:10.1016/j.fertnstert.2011.06.041)
- Kuiper GG, Enmark E, Peltö-Huikko M, Nilsson S & Gustafsson JA 1996 Cloning of a novel receptor expressed in rat prostate and ovary. *PNAS* **93** 5925–5930. (doi:10.1073/pnas.93.12.5925)
- Monsavaiz D, Dyson MT, Yin P, Navarro A, Coon JS 5th, Pavone ME & Bulun SE 2016 Estrogen receptor β regulates endometriotic cell survival through serum and glucocorticoid-regulated kinase activation. *Fertility and Sterility* **105** 1266–1273. (doi:10.1016/j.fertnstert.2016.01.012)
- Montagna P, Capellino S, Villaggio B, Remorgida V, Ragni N, Cutolo M & Ferrero S 2008 Peritoneal fluid macrophages in endometriosis: correlation between the expression of estrogen receptors and inflammation. *Fertility and Sterility* **90** 156–164. (doi:10.1016/j.fertnstert.2006.11.200)
- Pace P, Taylor J, Suntharalingam S, Coombes RC & Ali S 1997 Human estrogen receptor beta binds DNA in a manner similar to and dimerizes with estrogen receptor alpha. *Journal of Biological Chemistry* **272** 25832–25838. (doi:10.1074/jbc.272.41.25832)
- Trukhacheva E, Lin Z, Reierstad S, Cheng YH, Milad M & Bulun SE 2009 Estrogen receptor (ER) beta regulates ERalpha expression in stromal cells derived from ovarian endometriosis. *Journal of Clinical Endocrinology and Metabolism* **94** 615–622. (doi:10.1210/jc.2008-1466)
- Vercellini P, Viganò P, Somigliana E & Fedele L 2014 Endometriosis: pathogenesis and treatment. *Nature Reviews Endocrinology* **10** 261–275. (doi:10.1038/nrendo.2013.255)
- Vince JE & Silke J 2016 The intersection of cell death and inflammasome activation. *Cellular and Molecular Life Sciences* **73** 2349–2367. (doi:10.1007/s00018-016-2205-2)
- White R, Lees JA, Needham M, Ham J & Parker M 1987 Structural organization and expression of the mouse estrogen receptor. *Molecular Endocrinology* **1** 735–744. (doi:10.1210/mend-1-10-735)
- Zhao Y, Gong P, Chen Y, Nwachukwu JC, Srinivasan S, Ko C, Bagchi MK, Taylor RN, Korach KS, Nettles KW, et al. 2015 Dual suppression of estrogenic and inflammatory activities for targeting of endometriosis. *Science Translational Medicine* **7** 271. (doi:10.1126/scitranslmed.3010626)

Received in final form 27 May 2016

Accepted 7 June 2016

Accepted Preprint published online 7 June 2016