REVIEW

Insights into calcium-sensing receptor trafficking and biased signalling by studies of calcium homeostasis

Caroline M Gorvin^{1,2}¹Institute of Metabolism and Systems Research (IMSR), University of Birmingham, Birmingham, UK²Centre for Endocrinology, Diabetes and Metabolism (CEDAM), Birmingham Health Partners, Birmingham, UKCorrespondence should be addressed to C M Gorvin: C.Gorvin@bham.ac.uk

Abstract

The calcium-sensing receptor (CASR) is a class C G-protein-coupled receptor (GPCR) that detects extracellular calcium concentrations, and modulates parathyroid hormone secretion and urinary calcium excretion to maintain calcium homeostasis. The CASR utilises multiple heterotrimeric G-proteins to mediate signalling effects including activation of intracellular calcium release; mitogen-activated protein kinase (MAPK) pathways; membrane ruffling; and inhibition of cAMP production. By studying germline mutations in the CASR and proteins within its signalling pathway that cause hyper- and hypocalcaemic disorders, novel mechanisms governing GPCR signalling and trafficking have been elucidated. This review focusses on two recently described pathways that provide novel insights into CASR signalling and trafficking mechanisms. The first, identified by studying a CASR gain-of-function mutation that causes autosomal dominant hypocalcaemia (ADH), demonstrated a structural motif located between the third transmembrane domain and the second extracellular loop of the CASR that mediates biased signalling by activating a novel β -arrestin-mediated G-protein-independent pathway. The second, in which the mechanism by which adaptor protein-2 σ -subunit (AP2 σ) mutations cause familial hypocalciuric hypercalcaemia (FHH) was investigated, demonstrated that AP2 σ mutations impair CASR internalisation and reduce multiple CASR-mediated signalling pathways. Furthermore, these studies showed that the CASR can signal from the cell surface using multiple G-protein pathways, whilst sustained signalling is mediated only by the $G_{q/11}$ pathway. Thus, studies of FHH- and ADH-associated mutations have revealed novel steps by which CASR mediates signalling and compartmental bias, and these pathways could provide new targets for therapies for patients with calcaemic disorders.

Key Words

- ▶ calcium-sensing receptor
- ▶ adaptor protein-2
- ▶ familial hypocalciuric hypercalcaemia
- ▶ autosomal dominant hypocalcaemia
- ▶ biased signalling

*Journal of Molecular
Endocrinology*
(2018) **61**, R1–R12

Introduction to the CASR

Extracellular calcium (Ca^{2+}_e) is required for diverse biological functions ranging from blood coagulation, mineralisation of bone matrix, muscle contraction

and hormone secretion (Brown 1991). Thus, calcium concentrations within the blood are tightly regulated. The parathyroid gland plays an essential role in calcium

homeostasis by detecting Ca^{2+}_e in the blood, and in response to hypocalcaemia, secretes the parathyroid hormone (PTH) to normalize serum calcium concentrations. PTH achieves this by: enhancing bone resorption; activating calcium reabsorption at the kidneys; and stimulating the synthesis of 1,25-dihydroxyvitamin D_3 , which mobilises intestinal calcium absorption (Fig. 1A) (Riccardi & Brown 2010). The net effect of these three pathways is to increase Ca^{2+}_e , which provides feedback inhibition to the parathyroid gland, to suppress PTH secretion (Conigrave & Ward 2013). The parathyroid is able to detect Ca^{2+}_e concentrations in the blood using the cell-surface-expressed calcium-sensing receptor (CASR), a class C G-protein-coupled receptor (GPCR), for which Ca^{2+} is the major ligand (Riccardi & Brown 2010, Conigrave & Ward 2013).

The CASR is a 1078-amino acid protein that exists at cell surfaces as a disulphide-linked homodimer (Ward *et al.* 1998), although it is capable of forming heterodimers (Gama *et al.* 2001, Chang *et al.* 2007). The CASR has a large extracellular domain, which was recently crystallized by two independent laboratories, and consists of a bilobed venus flytrap ligand binding domain (VFTD) and a cysteine-rich domain (CRD) (Geng *et al.* 2016, Zhang *et al.* 2016) (Fig. 1B). Ca^{2+}_e binds between the two lobes (lobe 1 and lobe 2) of the VFTD, which initiates a conformational change, facilitating lobe 2–lobe 2 dimerisation and allowing the CRDs to interact (Geng *et al.* 2016, Zhang *et al.* 2016). These conformational changes are predicted to re-orientate the seven-transmembrane (TM) domain (Geng *et al.* 2016),

and consequently activate the associated G-proteins and initiate signal transduction (Standfuss *et al.* 2011). The CASR has been reported to couple to multiple G-protein subtypes, but predominantly signals by: the $\text{G}_{i/o}$ pathway, to suppress cAMP and activate mitogen-activated protein kinase (MAPK) cascades (Kifor *et al.* 2001, Thomsen *et al.* 2012); the $\text{G}_{q/11}$ -phospholipase C (PLC)-mediated pathway, to generate inositol 1,4,5-trisphosphate (IP_3) and diacylglycerol, that activate intracellular calcium (Ca^{2+}_i) mobilisations (Hofer & Brown 2003) and MAPK pathways, respectively. The CASR can also couple to a G-protein-independent mechanism involving β -arrestin proteins to also activate MAPK signalling (Thomsen *et al.* 2012) (Fig. 2).

Human disorders of the CASR provide insights into receptor signalling mechanisms

The importance of the CASR in the regulation of Ca^{2+}_e is highlighted in patients with germline mutations of the receptor that have disorders of calcium homeostasis. Heterozygous loss-of-function mutations in the CASR lead to the autosomal dominant condition familial hypocalciuric hypercalcaemia (FHH) (Pollak *et al.* 1993), which is characterised by lifelong mild-to-moderate hypercalcaemia, normal or mildly raised serum PTH levels and low urinary calcium excretion (Firek *et al.* 1991, Marx 2015). FHH is considered to be a benign disorder as most patients are asymptomatic, but it can be associated with chondrocalcinosis and pancreatitis in some cases (Pearce *et al.* 1996a, Volpe *et al.* 2009, Hannan & Thakker 2013).

Figure 1

Regulation of extracellular calcium at the parathyroid glands. (A) Schematic illustrating how the parathyroid glands respond to changes in extracellular calcium concentrations $[\text{Ca}^{2+}_e]$ in the blood. In the presence of low $[\text{Ca}^{2+}_e]$ in the blood, the parathyroid glands secrete parathyroid hormone (PTH). PTH binds to PTH receptors at the kidney, resulting in calcium reabsorption, and activation of 1,25-dihydroxyvitamin D_3 ; which in turn acts upon bone to mediate calcium release, and the intestine to activate calcium resorption. PTH also acts directly on bone to activate calcium release. The tissue-specific effects on calcium are shown in the hatched box with an arrow to indicate that these are increased by low $[\text{Ca}^{2+}]_{\text{blood}}$. (B) Schematic of the homodimeric calcium-sensing receptor (CASR) showing the two protomers (CASR1 and CASR2) within the plasma membrane (PM). The CASR is expressed highly at the parathyroid gland and its major function is to detect $[\text{Ca}^{2+}_e]$ and regulate PTH secretion accordingly. The CASR has a large extracellular domain (ECD) comprising the venus flytrap domain (VFTD) that contains two lobes (lobe 1 and lobe 2), and a cysteine-rich domain (CRD). The CRD connects the ECD to the 7-transmembrane domains (TMD), and the CASR has a long cytoplasmic intracellular domain (ICD). Ionised calcium binds between the two lobes of the VFTD. In the presence of high calcium, the CASR is activated, leading to changes in its signal transduction and reduced PTH secretion.

Figure 2

Major signalling pathways of the calcium-sensing receptor (CASR). Schematic diagram of the calcium-bound homodimeric CASR and its major signalling pathways. The CASR is expressed at the plasma membrane (PM) where its function is to detect $[Ca^{2+}_e]$ in the blood. The CASR activates two major signalling pathways: $G_{q/11}$ and $G_{i/o}$. Activation of the $G_{q/11}$ pathway leads to stimulation of its major effector protein phospholipase C (PLC), which hydrolyses phosphatidylinositol 4,5-bisphosphate (PIP₂) to the second messengers diacylglycerol (DAG) and inositol trisphosphate (IP₃). DAG activates protein kinase C (PKC) and the mitogen-activated protein kinase (MAPK) signalling pathway. IP₃ binds to IP₃ receptors at the ER, which mobilise intracellular calcium (Ca^{2+}_i) release into the cytoplasm. This increase in Ca^{2+}_i can further activate MAPK signalling. The CASR also activates $G_{i/o}$ signalling pathways, which inhibits adenylate cyclase (AC), resulting in reductions in cAMP and protein kinase A (PKA) activity. This reduction in PKA relieves the inhibition on MAPK signalling and therefore provides another activation pathway for MAPK signalling. The net effect of all these signalling pathways is a change in gene transcription.

Homozygous and compound heterozygous loss-of-function mutations of the CASR are the common cause of neonatal severe hyperparathyroidism (NSHPT), which is characterised by marked elevations in serum calcium and PTH, failure to thrive and hyperparathyroid bone disease (Pollak *et al.* 1993, Chattopadhyay & Brown 2006, Hannan & Thakker 2013). On occasion, heterozygous CASR mutations may lead to an NSHPT-like phenotype, and this is largely due to the dominant-negative nature of the mutant receptor on the wild-type (WT) CASR in these cases (Pearce *et al.* 1995, Obermannova *et al.* 2009).

Gain-of-function mutations of the CASR cause autosomal dominant hypocalcaemia (ADH) characterised by mild-to-moderate hypocalcaemia and inappropriately low or normal PTH concentrations (Pearce *et al.* 1996b, Hannan & Thakker 2013). Up to 50% of patients present with hypocalcaemic symptoms of paraesthesia, carpopedal spasms, seizures and ectopic calcification of the kidneys and basal ganglia (Pearce *et al.* 1996b,

Hannan & Thakker 2013). Some patients with gain-of-function mutations in CASR may have Bartter syndrome type 5, which is characterised by renal salt wasting, hypokalaemia, hyperreninaemia and hyperaldosteronaemia (Vargas-Poussou *et al.* 2002, Watanabe *et al.* 2002).

Biased signalling of the CASR

Functional studies in HEK293 cells have demonstrated that disease-causing mutations may influence CASR signalling responses in a biased manner (Leach *et al.* 2012). Despite the capability of the CASR to signal via multiple signalling pathways, *in vitro* studies in HEK293 cells have shown that it preferentially couples to the Ca^{2+}_i signalling pathway (Leach *et al.* 2012). In contrast, disease-causing CASR mutants have been shown to switch this preferential signalling, with some FHH1-causing mutations signalling equally via the Ca^{2+}_i and MAPK pathways, or predominantly via MAPK pathways, while many ADH1-associated mutants couple more strongly to Ca^{2+}_i (Leach *et al.* 2012). Furthermore, studies of positive and negative allosteric CASR modulating compounds, revealed they too can mediate a biased signalling response, with both classes of drugs influencing Ca^{2+}_i to a greater extent than ERK1/2 phosphorylation (Leach *et al.* 2013). Thus, these findings established that agonist-induced CASR signalling may occur in a biased manner, although the GPCR structural motifs mediating ligand-dependent bias were not described.

Novel insights into mechanisms by which CASR can mediate signalling bias

Recently, a novel ADH1-associated CASR mutation, Arg680Gly, has provided some insights into the molecular mechanisms mediating signalling bias (Gorvin *et al.* 2018a). *In vitro* analyses showed that Ca^{2+}_i signalling in cells expressing the Arg680Gly mutation was not different to that observed in WT cells, in contrast to previously reported ADH1-associated CASR mutants (Gorvin *et al.* 2018a). However, the Arg680Gly mutation did enhance MAPK signalling (Gorvin *et al.* 2018a). Furthermore, this type of signalling bias, in which an ADH1 mutation enhances MAPK signalling but does not affect Ca^{2+}_i , has not previously been described and provided an opportunity to explore the role of the Arg680 residue in CASR structure-function. As MAPK acts as a convergence pathway for multiple CASR signalling pathways (Fig. 2), each was investigated using a single reporter assay measuring

luciferase upstream of a serum-response element (SRE), which acts as a measure for MAPK-mediated transcription (Gorvin *et al.* 2017b). By applying compounds that specifically block the $G_{q/11}$ (YM-245890 and UBO-QIC), the $G_{i/o}$ (pertussis toxin) or the β -arrestin (β -arrestin-1 and β -arrestin-2 targeting siRNA) pathways, it was shown that the Arg680Gly mutation enhanced MAPK signalling by a β -arrestin1/2-mediated pathway (Takasaki *et al.* 2004, Schrage *et al.* 2015, Gorvin *et al.* 2018a).

The structural location of the mutant residue within the CASR transmembrane domain (TMD) region provides some insights into the likely mechanism mediating this bias. Homology modelling of the CASR TMD, based upon the structure of the closely related human metabotropic glutamate receptor 1 (mGluR1) (Hu *et al.* 2005, Gorvin *et al.* 2018a), predicts that the Arg680 residue lies at the extracellular side of TM3, and that the residue forms salt-bridge connections with the side chain of adjacent residues in extracellular loop 2 (ECL2) (Glu767) or TM7 (Glu837) (Fig. 3). Such connections between residues within TMDs or with ECLs of GPCRs are known to be important in receptor activity. For example, TM3 and TM6 of the β 2AR form ionic interactions which lock the receptor in an inactive state, and conformational changes within these helices govern G-protein coupling at the cytoplasmic face of the receptor (Ballesteros *et al.* 2001, Rasmussen *et al.* 2007). Furthermore, disruption of a salt-bridge in β 2-AR allows lateral displacement of TM3 away from TM4 and TM5, facilitating β -arrestin binding (Shukla *et al.* 2014), and it was therefore hypothesised that the CASR Arg680Gly mutation may break a salt-bridge between TM3 and ECL2 or TM7, allowing β -arrestin to bind more readily at the CASR cytoplasmic face (Gorvin *et al.* 2018a). The importance of the Arg680, Glu767 and Glu837 residues in the CASR has previously been recognised. Mutation of Glu767 and Glu837 has been shown to increase signalling by the CASR (Hu *et al.* 2002, 2005, 2006, Uckun-Kitapci *et al.* 2005), and previous structural homology models that investigated the binding of allosteric modulator drugs with the CASR demonstrated that the three residues form critical contacts for drug binding (Miedlich *et al.* 2004, Petrel *et al.* 2004), indicating their importance in CASR structure-function. To investigate the hypothesis that a salt-bridge is disrupted by mutation of Arg680, a series of mutations were engineered at the Arg680, Glu767 and Glu837 residues. First, Glu767 and Glu837 were mutated to Arg residues, to introduce unfavourable electrostatic interactions. Under these conditions, the Glu767Arg engineered mutant was shown to enhance MAPK activity in a similar way to the Arg680Gly ADH1 mutant,

Figure 3

Disruption of an Arg680–Glu767 salt-bridge within CASR mediates signalling bias by activation of a G-protein independent, β -arrestin pathway. (A) Schematic diagram of a CASR monomer at the plasma membrane (PM) showing the seven-transmembrane domains (TM1–7) with extracellular loops 1–3 (ECL1–3) and intracellular loops 1–3 (ICL1–3). Arg680 is located at the extracellular end of TM3 and is predicted to form a salt-bridge with either a Glu767 residue on ECL2 or a Glu837 residue on TM7. (B) Homology model of the CASR TM3, TM7 and ECL2 region, reproduced with permission from (Gorvin *et al.* 2018a). The homology model is based on the published structure of mGluR1 (Wu *et al.* 2014). The Arg680 residue is shown projecting from TM3 and is predicted to form a salt-bridge with Glu767 on ECL2. The Glu837 residue on TM7 lies at a 5.7 Å distance from the Arg680 residue, and therefore formation of a salt-bridge between Arg680 and Glu837 is less likely, but was tested as the homology model may not reflect the true state of CASR.

a response that was also β -arrestin sensitive (Gorvin *et al.* 2018a). In contrast, responses in cells expressing the CASR Glu837Arg engineered mutant were similar to those in cells expressing WT CASR (Gorvin *et al.* 2018a). To confirm that a salt-bridge is required between residues 680 and 767 in the CASR, a double mutant was generated by mutating the Glu767Arg mutant receptor with an additional missense mutation of Arg680 to Glu680. This should allow the salt-bridge between the two residues to reform. Indeed, in this double mutant (Glu680-Arg767) CASR, MAPK signalling was restored to levels similar to those seen in WT cells (Gorvin *et al.* 2018a).

Thus, these studies have revealed some of the important structural motifs of CASR that mediate signalling bias, and discovery of this novel β -arrestin-specific pathway may help facilitate the development of targeted therapeutics for CASR. Furthermore, it demonstrates the importance of investigating multiple signalling outputs downstream of CASR, to ensure that potentially disease-causing mutations are not classified as benign polymorphisms (Gorvin *et al.* 2018a).

Insights from human genetic mutations in components of the CASR signalling pathway

Both FHH and ADH are genetically heterozygous conditions with mutations in the CASR accounting for approximately 65% and 70% of cases, respectively (Hannan *et al.* 2012, Nesbit *et al.* 2013a). Two further

genetically distinct forms of FHH, and one further distinct form of ADH, have been described (Mannstadt *et al.* 2013, Nesbit *et al.* 2013a,b). Heterozygous loss- and gain-of-function germline mutations in the α -subunit of G-protein 11 ($G\alpha_{11}$), a component of the CASR signalling pathway, give rise to FHH type-2 (FHH2) and ADH type-2 (ADH2), respectively (Mannstadt *et al.* 2013, Nesbit *et al.* 2013a). Only four FHH2 and six ADH2 mutations have been described to date, and therefore these mutations account for only a small number of cases of these disorders (Mannstadt *et al.* 2013, Nesbit *et al.* 2013a, Li *et al.* 2014, Gorvin *et al.* 2016, 2017a, Piret *et al.* 2016). FHH type-3 (FHH3) is due to mutations in the sigma subunit of the adaptor protein-2 ($AP2\sigma$), which plays a fundamental role in clathrin-mediated endocytosis of transmembrane proteins, such as GPCRs (Nesbit *et al.* 2013b). The $AP2\sigma$ protein is ubiquitously expressed, and clathrin-mediated endocytosis is a critical cellular process; however, FHH is a largely benign condition, and the phenotypes observed in FHH3 patients are largely CASR-specific. Thus, by studying the $AP2\sigma$ mutations identified in FHH3, which have been described in a single residue (Nesbit *et al.* 2013b, Hannan *et al.* 2015), novel insights into the trafficking and signalling mechanisms of CASR have been elucidated, and indicate an important interplay between these two processes (Gorvin *et al.* 2018b).

Trafficking of the CASR

The cell surface expression of the CASR is important for detecting extracellular ligand and signalling by the receptor, and therefore the plasma membrane expression of CASR is carefully regulated. This involves multiple pathways including: receptor synthesis and secretion, trafficking to the plasma membrane and removal of the CASR from the cell surface by endocytosis.

Regulation of cell surface expression by functional desensitisation

In contrast to most GPCRs, which undergo agonist-dependent desensitisation by phosphorylation and/or by β -arrestin proteins, functional desensitisation appears to have only a minimal impact on CASR expression, and there are inconsistencies between findings in these studies (Bouschet *et al.* 2005, Pi *et al.* 2005, Lorenz *et al.* 2007, Thomsen *et al.* 2012). Initial studies of the CASR showed that several proteins desensitise the receptor. These include: PKC, which phosphorylates the receptor following activation of signalling pathways and recruits

β -arrestin; G-protein regulatory kinase (GRK) 2, which binds to $G\alpha_q$ and inhibits its signalling; and GRK4, which phosphorylates CASR to facilitate desensitisation (Pi *et al.* 2005, Lorenz *et al.* 2007). Furthermore, GRKs and β -arrestin have independent functions as combined treatment of cells with these proteins enhances desensitisation (Lorenz *et al.* 2007). The traditional view of GPCR desensitisation and internalisation was one in which the receptor is phosphorylated and β -arrestin recruited, followed by internalisation of the receptor by clathrin-mediated endocytosis, which is facilitated by interactions between β -arrestin and AP2 (Shukla *et al.* 2011). While studies of CASR showed that β -arrestin can be recruited, this did not enhance receptor internalisation, which only occurred when cells were treated with high concentrations of Ca^{2+}_e (10 mM) (Lorenz *et al.* 2007). β -Arrestin is also now recognised to have another function, as a scaffold protein that facilitates signalling either at the plasma membrane or for some GPCRs, at the endosome (Shukla *et al.* 2014). This function of β -arrestin has been recognised for CASR in some studies, providing seemingly contradictory information to that in studies of functional desensitisation. Thus, treatment of cells with dominant-negative forms of β -arrestin1 or β -arrestin2, or with siRNA targeting β -arrestin1 or β -arrestin2, reduces the pERK and membrane ruffling signals downstream of CASR (Bouschet *et al.* 2005, Thomsen *et al.* 2012, Gorvin *et al.* 2018b). Further studies are required to determine whether the discrepancies within these data sets are due to experimental differences, differences in cell type or if both desensitisation and enhanced signalling occur downstream of CASR, but at different spatial or temporal points.

Regulation of cell surface expression by the secretory pathway and agonist-driven insertional signalling (ADIS)

The cell surface expression of CASR and mechanisms by which the receptor is synthesised and exported to the plasma membrane are generally well understood. The CASR is synthesised at the endoplasmic reticulum (ER) where it forms homodimers (Pidashveva *et al.* 2006) and undergoes quality control and immature glycosylation steps prior to progression to the Golgi (Fan *et al.* 1997). At the Golgi, the CASR undergoes further maturation. This forward trafficking through the secretory pathway involves a number of regulatory proteins that have been well described in previous reviews (Huang *et al.* 2011, Breitwieser 2013). The core-glycosylated CASR is then retained within pre-plasma membrane compartments (Fan *et al.* 1997).

Therefore, CASR is retained intracellularly within two large intracellular reserves: one at the ER and one at pre-plasma membrane compartments (Breitwieser 2012).

The large pool of fully mature CASR plays a unique role in regulation of Ca^{2+}_e . Unlike many GPCRs, the CASR is chronically exposed to its ligand at baseline concentrations, and thus the receptor undergoes very little functional desensitisation (Gama & Breitwieser 1998). Furthermore, CASR is able to elicit signalling responses for as long as elevated Ca^{2+}_e is available (Pi *et al.* 2005, Lorenz *et al.* 2007, Grant *et al.* 2011). Experimental studies have sought to explain this apparent paradox and using a combination of total internal reflection fluorescence microscopy (TIRFm) and a construct that allows the simultaneous measurement of CASR insertion within plasma membranes and endocytosis (known as BSEP-CASR), a new model for CASR cell surface expression has emerged (Fig. 4). This model proposes that in the basal state, CASR is only weakly expressed at the plasma membrane, but on exposure to increases in Ca^{2+}_e there is an increase in anterograde trafficking through the secretory pathway (Grant *et al.* 2011) (Fig. 4). This increase in the secretory pathway involves both mature CASR located in pre-plasma membrane regions, and newly synthesised CASR from the ER (Grant *et al.* 2011,

Breitwieser 2013), and at present, the mechanisms that trigger the ADIS events are not fully elucidated. However, studies in which ADIS and CASR signalling were measured simultaneously revealed that elevated Ca^{2+}_i and the $\text{G}\alpha_{q/11}$ pathway are likely involved (Grant *et al.* 2012, Gorvin *et al.* 2018b). Furthermore, the 14-3-3 proteins, which bind CASR at an Arg-rich site within the C-terminus, limit the release of CASR from the ER, and therefore regulate CASR cell surface expression by reducing the ADIS-mobilisable pool of the receptor (Grant *et al.* 2015). The presence of the ADIS system and constitutive endocytosis may also explain some of the inconsistencies between studies of CASR trafficking, as endocytosis of the receptor is balanced by the continual insertion of new CASR from the secretory pathway (Breitwieser 2013). Those studies in which a tagged construct and cell surface labelling (e.g. FLAG) was used in isolation, without also measuring insertion of new CASR within the plasma membrane, may not reflect the physiological state. Similarly, measurement of total cell surface expression of receptor cannot be used as a surrogate for determining the rate of internalisation.

These studies have demonstrated that trafficking of the CASR plays a critical role in receptor signalling and calcium homeostasis. More recent studies, building upon these initial findings, have identified further insights

Figure 4

Mechanisms by which calcium-sensing receptor (CASR) expression is regulated at plasma membranes. Schematic showing trafficking pathways regulating the expression of CASR at the plasma membrane. The CASR is continuously synthesised at the endoplasmic reticulum (ER) and undergoes post-translational modifications at the ER and Golgi, before its export to the plasma membrane surface. In addition to this constitutive anterograde trafficking of the CASR, an additional export pathway for the CASR has been described. This pathway is activated by high $[\text{Ca}^{2+}]_e$ and has been named the agonist-driven insertional signalling, or ADIS, pathway. Removal of the CASR from the cell surface is regulated by clathrin-mediated endocytosis that requires the heterotetrameric adaptor protein-2 (AP2). AP2 binds directly to transmembrane proteins using its μ -subunit or σ -subunit. The CASR has a putative dileucine motif within its C-terminus with which it is predicted to bind to AP2 σ . The α -subunit and β -subunit have large appendages that are important for binding to clathrin coat proteins, plasma membrane phospholipids and endocytic accessory proteins. Once internalised within clathrin-coated vesicles, CASR is targeted to the endosomal-lysosomal system. Experimental evidence suggests that CASR is degraded at the lysosome, and very little, if any, recycling of the receptor occurs.

into CASR cell surface expression and endocytosis, and recognised further connections between the trafficking and signalling of this receptor.

Regulation of cell surface expression by endocytosis

The CASR has been described to have two types of internalisation: a constitutive pathway (Grant *et al.* 2011), and an agonist-driven pathway (Lorenz *et al.* 2007, Gorvin *et al.* 2018b). The CASR was shown to internalise using clathrin-mediated endocytosis in early studies (Holstein *et al.* 2004). In addition, CASR has been described to associate with several proteins that facilitate clustering at the plasma membrane and therefore increase the efficiency of internalisation, or act as scaffolds to enable signalling to occur. One such protein is caveolin-1, with which CASR has been shown to coimmunoprecipitate (Kifor *et al.* 1998). CASR has been described to be enriched in caveolae structures in parathyroid chief cells and osteosarcoma cells (Kifor *et al.* 1998, Jung *et al.* 2005, Sun & Murphy 2010). CASR-mediated signalling is impaired in cells treated with caveolin-1 targeting siRNA, and it is likely that this signalling function is facilitated by caveolae acting as signalling hubs, allowing the CASR, G-proteins and PKC to cluster together (Kifor *et al.* 1998, Jung *et al.* 2005, Sun & Murphy 2010). However, it is currently unknown whether caveolin facilitates CASR clustering within caveolae structures to enhance signalling, or whether the signalling itself drives this clustering to encourage endocytosis (Breitwieser 2013). Filamin is an actin-binding cytoskeleton protein that is important for protein scaffolding (Hjalm *et al.* 2001) and may also facilitate clustering that aids in receptor endocytosis. Filamin binds to the CASR C-terminus and increases total cellular content of CASR by preventing its proteosomal degradation (Hjalm *et al.* 2001), with some studies showing CASR expression is reduced in cells treated with siRNA targeted against filamin. However, other reports show no such changes in CASR expression, and this likely requires further investigation (Huang *et al.* 2006, Mingione *et al.* 2017). The net result of filamin A binding to CASR is increased MAPK signalling by the receptor (Hjalm *et al.* 2001, Pi *et al.* 2002).

Effect of AP2 σ mutations on CASR signalling and trafficking

Mutations in AP2 σ have been demonstrated to cause FHH3, and are associated with impaired CASR-mediated Ca²⁺_i signalling (Nesbit *et al.* 2013b). The AP2 complex is a ubiquitously expressed heterotetrameric protein

that plays a fundamental role in the clathrin-mediated endocytosis of transmembrane proteins, such as GPCRs. The two larger subunits, α and β , have appendages that bind to the clathrin coat proteins, plasma membrane phospholipids and endocytic accessory proteins (e.g. β -arrestin); while the two smaller subunits, μ and σ , bind to endocytic motifs of cargo proteins (Collins *et al.* 2002, Kelly *et al.* 2008, Jackson *et al.* 2010, Kirchhausen *et al.* 2014) (Fig. 4). The AP2 μ subunit recognises tyrosine-based motifs and the AP2 σ subunit recognises dileucine-based motifs (Haucke & De Camilli 1999, Kelly *et al.* 2008) (Fig. 4). FHH3-associated mutations in the AP2 σ protein have been reported to affect the Arg15 residue, and structural modelling studies using a published structure of the AP2 complex have shown these missense mutations (to Cys15, His15 and Leu15) likely disrupt interactions with a putative dileucine motif in the CASR C-terminus (Nesbit *et al.* 2013b), and thus affect CASR endocytosis.

Initially, this hypothesis was tested using an ELISA assay of total CASR at cell surfaces in HEK293 cells stably overexpressing CASR (HEK-CASR), and transiently transfected with AP2 σ -mutant proteins. This showed that AP2 σ -mutant expressing cells had increased CASR cell surface expression when compared to AP2 σ -WT cells following stimulation with 5 mM Ca²⁺_e (Nesbit *et al.* 2013b). Thus, it was concluded that CASR endocytosis is impaired in these cells, resulting in increased total CASR cell surface expression (Nesbit *et al.* 2013b). More detailed studies characterising the ADIS and endocytosis components of CASR regulation in AP2 σ mutant cells using TIRFm and the BSEP-CASR construct showed both ADIS and CASR endocytosis were impaired, resulting in the net effect of an increased total CASR cell surface expression in cells expressing AP2 σ mutant protein compared to WT cells (Gorvin *et al.* 2018b). Additional TIRFm studies focussing on CASR and clathrin demonstrated that CASR and clathrin colocalise at plasma membranes, and that the duration of this colocalisation is prolonged in AP2 σ -mutant expressing cells (Gorvin *et al.* 2018b). Furthermore, the vesicles containing both CASR and clathrin in AP2 σ -mutant cells, when compared to WT cells, were less motile, which is an indication that the vesicles are less likely to result in viable endocytic events (Rappoport & Simon 2003, Gorvin *et al.* 2018b). Thus, cells expressing AP2 σ -mutant proteins have delayed recruitment of clathrin, and colocalisation with CASR is prolonged, resulting in impaired CASR endocytosis (Gorvin *et al.* 2018b).

The effect of the FHH3-associated AP2 σ -mutant proteins on multiple CASR-mediated signalling pathways has been characterised in a number of cellular assays

in HEK293 cells stable overexpressing AP2 σ WT and mutant proteins and lymphoblastoid cell lines derived from blood samples from FHH3 patients with the AP2 σ -Cys15 mutation (Nesbit *et al.* 2013b, Gorvin *et al.* 2018b). These studies confirmed that AP2 σ mutations of the Arg15 residue impair Ca^{2+}_i mobilisation, phosphorylated ERK1/2 (pERK1/2) MAPK signalling, membrane ruffling and suppression of cAMP, and that all of these pathways occur downstream of $\text{G}_{q/11}$ and $\text{G}_{i/o}$ (Nesbit *et al.* 2013b, Gorvin *et al.* 2018b). Therefore, these studies have shown that AP2 σ mutations reduce endocytosis, resulting in increased CASR cell surface expression, but paradoxically decrease CASR-mediated signalling (Gorvin *et al.* 2018b). To explain this paradox, we hypothesised that CASR may be able to continue signalling from within the cell (i.e. sustained signalling) (Gorvin *et al.* 2018b). Such sustained signalling has been previously reported for some class A (e.g. β 2-adrenergic receptor (β 2AR), dopamine receptor D1 (DRD1), thyroid-stimulating hormone receptor, vasopressin receptor 2 (V2R) and luteinizing hormone receptor (LHR)), and class B (e.g. parathyroid hormone 1 receptor (PTH1R)) GPCRs (Calebiro *et al.* 2009, Ferrandon *et al.* 2009, Kotowski *et al.* 2011, Feinstein *et al.* 2013, Irannejad *et al.* 2013, Jean-Alphonse *et al.* 2014). Thus, in cells with the AP2 σ mutation, in which there is impaired endocytosis, the availability of internalised receptors from which sustained signals could emanate would be reduced and thus the net effect would be impaired overall CASR-

mediated signalling. To test this hypothesis, a combination of imaging and biochemical analyses, along with chemical inhibitors, were used in HEK-CASR, HEK-AP2 σ and CRISPR-Cas generated β -arrestin knockout cells to assess CASR-mediated MAPK signalling (Gorvin *et al.* 2018b). To assess sustained signalling, two primary assays were used: (1) assessment of pERK1/2 over 60 min following treatment of cells with a 5 min pulse of 5 mM Ca^{2+}_e ; and (2) analysis of SRE luciferase reporter responses over 12 h, following a 5 min pulse of 5 mM Ca^{2+}_e . Using these methods, MAPK-sustained signals were demonstrated in HEK-CASR cells, and evidence for an internal, likely endosomal, source was shown in three ways (Gorvin *et al.* 2018b) (Fig. 5). First, addition of Dyngo, a chemical inhibitor of dynamin, which is required for vesicle scission during clathrin-mediated endocytosis, abolishes sustained signals, whilst rapid plasma membrane-mediated signals remain intact. Loss of this sustained response in Dyngo-treated cells was not due to increased apoptosis or decreased proliferation, and was unaffected by inhibition of CASR protein synthesis as the sustained rise in pERK1/2 was not blocked by tunicamycin, an inhibitor of glycosylation of newly synthesised CASR. Second, sustained signals were reduced in cells expressing a dominant-negative Rab5 protein, which delays maturation of early endosomes and therefore slows clathrin-mediated endocytosis. Third, reduction or loss of these sustained signals was observed in cells expressing AP2 σ mutant proteins (Fig. 5).

Figure 5

The calcium-sensing receptor (CASR) can activate plasma membrane and endosomal signalling pathways. Schematic diagram showing the signalling pathways that occur downstream of the CASR. As described in Fig. 2, CASR predominantly signals via: the $\text{G}_{q/11}$ pathway, leading to activation of the intracellular calcium (Ca^{2+}_i) and mitogen-activated protein kinase (MAPK) pathways; and the $\text{G}_{i/o}$ pathway, leading to activation of MAPK and reductions in the cAMP pathways. These pathways are activated rapidly (within 2–5 min), and originate at the plasma membrane. The CASR can also signal from an internal location using a $\text{G}_{q/11}$ pathway. This sustained signal that occurs later than the plasma membrane signal (from 30 min) is sensitive to: global clathrin-mediated endocytosis blockade (shown experimentally using Dyngo); inhibition of CASR-mediated internalisation (demonstrated in cells expressing FHH3-associated AP2 σ mutants); and maturation of internalised vesicles to the early endosome (demonstrated experimentally using a dominant-negative (DN) form of Rab5). Thus, this sustained signal likely arises within endosomes.

The G-proteins involved in these sustained signals were also explored as MAPK signalling is a convergence pathway for multiple CASR-mediated signalling pathways (Fig. 2). The ability of the $G_{\alpha_{q/11}}$ and $G_{\alpha_{i/o}}$ pathways to activate CASR sustained signals was investigated using G-protein-specific inhibitors (UBO-QIC and PTx) and the SRE luciferase reporter assay, which showed that inhibition of both $G_{\alpha_{q/11}}$ and $G_{\alpha_{i/o}}$ impaired the early, plasma-membrane-mediated CASR signal, while only $G_{\alpha_{q/11}}$ was important for the later sustained signal. Furthermore, confocal microscopy confirmed that G_{α_q} and phosphatidylinositol 4,5-bisphosphate (PIP₂), the lipid hydrolysed by PLC, colocalise with CASR at plasma membranes and a subpopulation of endosomes (Gorvin *et al.* 2018b). In addition, inhibitors of the PLC-DAG-IP₃ pathway (U73122, GF-109203X and 2-aminoethoxydiphenyl borate (2-APB), which inhibit PLC, PKC and the IP₃-receptor, respectively) were shown to reduce sustained pERK1/2 signals, indicating that $G_{\alpha_{q/11}}$ and its signalling pathway are important for CASR-mediated signalling (Gorvin *et al.* 2018b). Finally, the effects of the β -arrestin scaffold proteins, which are important for the sustained endosomal signalling of some GPCRs such as V2R and PTH1R (Feinstein *et al.* 2013, Wehbi *et al.* 2013), were assessed in HEK-CASR cells with deletion of the β -arrestin1 and β -arrestin2 genes. In these cells, both pERK and SRE reporter responses were unaffected by deletion of the β -arrestin proteins (Gorvin *et al.* 2018b). Thus, the CASR mediates some MAPK signals from endosomes using G_{α_q} and PLC, but does not require β -arrestin for this pathway.

Conclusions and important lessons for the future (or unanswered questions)

Since the cloning of the CASR gene 25 years ago (Brown *et al.* 1993), many insights have been gained into the role of the CASR in calcium homeostasis, and by studying patients with mutations within this gene, we have learnt much about CASR signalling and trafficking mechanisms. The discovery that CASR can signal from within the cell, most likely from an endosomal source, and that disruption of a specific structural motif can mediate G-protein independent signalling bias, opens up many new avenues of investigation. How do mutations in the CASR affect these endosomal pathways? Can drugs that target the CASR affect these pathways? How are trafficking and signalling of the CASR so intimately linked? Future studies investigating these questions will undoubtedly

reveal further complexities into the regulation of CASR, and could provide mechanisms relevant to other GPCRs.

Declaration of interest

The author declares that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

References

- Ballesteros JA, Jensen AD, Liapakis G, Rasmussen SG, Shi L, Gether U & Javitch JA 2001 Activation of the beta 2-adrenergic receptor involves disruption of an ionic lock between the cytoplasmic ends of transmembrane segments 3 and 6. *Journal of Biological Chemistry* **276** 29171–29177. (<https://doi.org/10.1074/jbc.M103747200>)
- Bouschet T, Martin S & Henley JM 2005 Receptor-activity-modifying proteins are required for forward trafficking of the calcium-sensing receptor to the plasma membrane. *Journal of Cell Science* **118** 4709–4720. (<https://doi.org/10.1242/jcs.02598>)
- Breitwieser GE 2012 Minireview: the intimate link between calcium sensing receptor trafficking and signaling: implications for disorders of calcium homeostasis. *Molecular Endocrinology* **26** 1482–1495. (<https://doi.org/10.1210/me.2011-1370>)
- Breitwieser GE 2013 The calcium sensing receptor life cycle: trafficking, cell surface expression, and degradation. *Best Practice and Research: Clinical Endocrinology and Metabolism* **27** 303–313. (<https://doi.org/10.1016/j.beem.2013.03.003>)
- Brown EM 1991 Extracellular Ca²⁺ sensing, regulation of parathyroid cell function, and role of Ca²⁺ and other ions as extracellular (first) messengers. *Physiological Reviews* **71** 371–411. (<https://doi.org/10.1152/physrev.1991.71.2.371>)
- Brown EM, Gamba G, Riccardi D, Lombardi M, Butters R, Kifor O, Sun A, Hediger MA, Lytton J & Hebert SC 1993 Cloning and characterization of an extracellular Ca(2+)-sensing receptor from bovine parathyroid. *Nature* **366** 575–580. (<https://doi.org/10.1038/366575a0>)
- Calebiro D, Nikolaev VO, Gagliani MC, de Filippis T, Dees C, Tacchetti C, Persani L & Lohse MJ 2009 Persistent cAMP-signals triggered by internalized G-protein-coupled receptors. *PLoS Biology* **7** e1000172. (<https://doi.org/10.1371/journal.pbio.1000172>)
- Chang W, Tu C, Cheng Z, Rodriguez L, Chen TH, Gassmann M, Bettler B, Margeta M, Jan LY & Shoback D 2007 Complex formation with the Type B gamma-aminobutyric acid receptor affects the expression and signal transduction of the extracellular calcium-sensing receptor. Studies with HEK-293 cells and neurons. *Journal of Biological Chemistry* **282** 25030–25040. (<https://doi.org/10.1074/jbc.M700924200>)
- Chattopadhyay N & Brown EM 2006 Role of calcium-sensing receptor in mineral ion metabolism and inherited disorders of calcium-sensing. *Molecular Genetics and Metabolism* **89** 189–202. (<https://doi.org/10.1016/j.ymgme.2006.07.003>)
- Collins BM, McCoy AJ, Kent HM, Evans PR & Owen DJ 2002 Molecular architecture and functional model of the endocytic AP2 complex. *Cell* **109** 523–535. ([https://doi.org/10.1016/S0092-8674\(02\)00735-3](https://doi.org/10.1016/S0092-8674(02)00735-3))
- Conigrave AD & Ward DT 2013 Calcium-sensing receptor (CaSR): pharmacological properties and signaling pathways. *Best Practice and*

- Research: *Clinical Endocrinology and Metabolism* **27** 315–331. (<https://doi.org/10.1016/j.beem.2013.05.010>)
- Fan G, Goldsmith PK, Collins R, Dunn CK, Krapcho KJ, Rogers KV & Spiegel AM 1997 N-linked glycosylation of the human Ca²⁺ receptor is essential for its expression at the cell surface. *Endocrinology* **138** 1916–1922. (<https://doi.org/10.1210/endo.138.5.5131>)
- Feinstein TN, Yui N, Webber MJ, Wehbi VL, Stevenson HP, King JD Jr, Hallows KR, Brown D, Bouley R & Vilardaga JP 2013 Noncanonical control of vasopressin receptor type 2 signaling by retromer and arrestin. *Journal of Biological Chemistry* **288** 27849–27860. (<https://doi.org/10.1074/jbc.M112.445098>)
- Ferrandon S, Feinstein TN, Castro M, Wang B, Bouley R, Potts JT, Gardella TJ & Vilardaga JP 2009 Sustained cyclic AMP production by parathyroid hormone receptor endocytosis. *Nature Chemical Biology* **5** 734–742. (<https://doi.org/10.1038/nchembio.206>)
- Firek AE, Kao PC & Heath H, 3rd 1991 Plasma intact parathyroid hormone (PTH) and PTH-related peptide in familial benign hypercalcemia: greater responsiveness to endogenous PTH than in primary hyperparathyroidism. *Journal of Clinical Endocrinology and Metabolism* **72** 541–546. (<https://doi.org/10.1210/jcem-72-3-541>)
- Gama L & Breitwieser GE 1998 A carboxyl-terminal domain controls the cooperativity for extracellular Ca²⁺ activation of the human calcium sensing receptor. A study with receptor-green fluorescent protein fusions. *Journal of Biological Chemistry* **273** 29712–29718. (<https://doi.org/10.1074/jbc.273.45.29712>)
- Gama L, Wilt SG & Breitwieser GE 2001 Heterodimerization of calcium sensing receptors with metabotropic glutamate receptors in neurons. *Journal of Biological Chemistry* **276** 39053–39059. (<https://doi.org/10.1074/jbc.M105662200>)
- Geng Y, Mosyak L, Kurinov I, Zuo H, Sturchler E, Cheng TC, Subramanyam P, Brown AP, Brennan SC, Mun HC, *et al.* 2016 Structural mechanism of ligand activation in human calcium-sensing receptor. *Elife* **5** e13662. (<https://doi.org/10.7554/eLife.13662>)
- Gorvin CM, Cranston T, Hannan FM, Rust N, Qureshi A, Nesbit MA & Thakker RV 2016 A G-protein subunit- α 11 loss-of-function mutation, Thr54Met, causes familial hypocalciuric hypercalcemia type 2 (FHH2). *Journal of Bone and Mineral Research* **31** 1200–1206. (<https://doi.org/10.1002/jbmr.2778>)
- Gorvin CM, Hannan FM, Cranston T, Valta H, Makitie O, Schalin-Jantti C & Thakker RV 2017a Cinacalcet rectifies hypercalcemia in a patient with familial hypocalciuric hypercalcemia type 2 (FHH2) caused by a germline loss-of-function Galphai1 mutation. *Journal of Bone and Mineral Research* **33** 32–41. (<https://doi.org/10.1002/jbmr.3241>)
- Gorvin CM, Hannan FM, Howles SA, Babinsky VN, Piret SE, Rogers A, Freidin AJ, Stewart M, Paudyal A, Hough TA, *et al.* 2017b Galphai1 mutation in mice causes hypocalcemia rectifiable by calcilytic therapy. *JCI Insight* **2** e91103. (<https://doi.org/10.1172/jci.insight.91103>)
- Gorvin CM, Babinsky VN, Malinauskas T, Nissen PH, Schou AJ, Hanyaloglu AC, Siebold C, Jones EY, Hannan FM & Thakker RV 2018a A calcium-sensing receptor mutation causing hypocalcemia disrupts a transmembrane salt bridge to activate β -arrestin biased signaling. *Sci Signal* **11** ean3714. (<https://doi.org/10.1126/scisignal.aan3714>)
- Gorvin CM, Rogers A, Hastoy B, Tarasov AI, Frost M, Sposini S, Inoue A, Whyte MP, Rorsman P, Hanyaloglu AC, *et al.* 2018b AP2 σ mutations impair calcium-sensing receptor trafficking and signaling revealing an endosomal pathway that spatially directs G-protein selectivity. *Cell Reports* **22** 1054–1066. (<https://doi.org/10.1016/j.celrep.2017.12.089>)
- Grant MP, Stepanchick A, Cavanaugh A & Breitwieser GE 2011 Agonist-driven maturation and plasma membrane insertion of calcium-sensing receptors dynamically control signal amplitude. *Science Signaling* **4** ra78. (<https://doi.org/10.1126/scisignal.2002208>)
- Grant MP, Stepanchick A & Breitwieser GE 2012 Calcium signaling regulates trafficking of familial hypocalciuric hypercalcemia (FHH) mutants of the calcium sensing receptor. *Molecular Endocrinology* **26** 2081–2091. (<https://doi.org/10.1210/me.2012-1232>)
- Grant MP, Cavanaugh A & Breitwieser GE 2015 14-3-3 proteins buffer intracellular calcium sensing receptors to constrain signaling. *PLoS ONE* **10** e0136702. (<https://doi.org/10.1371/journal.pone.0136702>)
- Hannan FM & Thakker RV 2013 Calcium-sensing receptor (CaSR) mutations and disorders of calcium, electrolyte and water metabolism. *Best Practice and Research: Clinical Endocrinology and Metabolism* **27** 359–371. (<https://doi.org/10.1016/j.beem.2013.04.007>)
- Hannan FM, Nesbit MA, Zhang C, Cranston T, Curley AJ, Harding B, Fratter C, Rust N, Christie PT, Turner JJ, *et al.* 2012 Identification of 70 calcium-sensing receptor mutations in hyper- and hypo-calcaemic patients: evidence for clustering of extracellular domain mutations at calcium-binding sites. *Human Molecular Genetics* **21** 2768–2778. (<https://doi.org/10.1093/hmg/dds105>)
- Hannan FM, Howles SA, Rogers A, Cranston T, Gorvin CM, Babinsky VN, Reed AA, Thakker CE, Bockenhauer D, Brown RS, *et al.* 2015 Adaptor protein-2 sigma subunit mutations causing familial hypocalciuric hypercalcaemia type 3 (FHH3) demonstrate genotype-phenotype correlations, codon bias and dominant-negative effects. *Human Molecular Genetics* **24** 5079–5092. (<https://doi.org/10.1093/hmg/ddv226>)
- Haucke V & De Camilli P 1999 AP-2 recruitment to synaptotagmin stimulated by tyrosine-based endocytic motifs. *Science* **285** 1268–1271. (<https://doi.org/10.1126/science.285.5431.1268>)
- Hjalm G, MacLeod RJ, Kifor O, Chattopadhyay N & Brown EM 2001 Filamin-A binds to the carboxyl-terminal tail of the calcium-sensing receptor, an interaction that participates in CaR-mediated activation of mitogen-activated protein kinase. *Journal of Biological Chemistry* **276** 34880–34887. (<https://doi.org/10.1074/jbc.M100784200>)
- Hofer AM & Brown EM 2003 Extracellular calcium sensing and signalling. *Nature Reviews Molecular Cell Biology* **4** 530–538. (<https://doi.org/10.1038/nrm1154>)
- Holstein DM, Berg KA, Leeb-Lundberg LM, Olson MS & Saunders C 2004 Calcium-sensing receptor-mediated ERK1/2 activation requires Galphai2 coupling and dynamin-independent receptor internalization. *Journal of Biological Chemistry* **279** 10060–10069. (<https://doi.org/10.1074/jbc.M312039200>)
- Hu J, Reyes-Cruz G, Chen W, Jacobson KA & Spiegel AM 2002 Identification of acidic residues in the extracellular loops of the seven-transmembrane domain of the human Ca²⁺ receptor critical for response to Ca²⁺ and a positive allosteric modulator. *Journal of Biological Chemistry* **277** 46622–46631. (<https://doi.org/10.1074/jbc.M207100200>)
- Hu J, McLarnon SJ, Mora S, Jiang J, Thomas C, Jacobson KA & Spiegel AM 2005 A region in the seven-transmembrane domain of the human Ca²⁺ receptor critical for response to Ca²⁺. *Journal of Biological Chemistry* **280** 5113–5120. (<https://doi.org/10.1074/jbc.M413403200>)
- Hu J, Jiang J, Costanzi S, Thomas C, Yang W, Feyen JH, Jacobson KA & Spiegel AM 2006 A missense mutation in the seven-transmembrane domain of the human Ca²⁺ receptor converts a negative allosteric modulator into a positive allosteric modulator. *Journal of Biological Chemistry* **281** 21558–21565. (<https://doi.org/10.1074/jbc.M603682200>)
- Huang C, Wu Z, Hujer KM & Miller RT 2006 Silencing of filamin A gene expression inhibits Ca²⁺-sensing receptor signaling. *FEBS Letters* **580** 1795–1800. (<https://doi.org/10.1016/j.febslet.2006.02.035>)
- Huang Y, Cavanaugh A & Breitwieser GE 2011 Regulation of stability and trafficking of calcium-sensing receptors by pharmacologic chaperones. *Advances in Pharmacology* **62** 143–173. (<https://doi.org/10.1016/B978-0-12-385952-5.00007-5>)
- Irannejad R, Tomshine JC, Tomshine JR, Chevalier M, Mahoney JP, Steyaert J, Rasmussen SG, Sunahara RK, El-Samad H, Huang B, *et al.*

- 2013 Conformational biosensors reveal GPCR signalling from endosomes. *Nature* **495** 534–538. (<https://doi.org/10.1038/nature12000>)
- Jackson LP, Kelly BT, McCoy AJ, Gaffry T, James LC, Collins BM, Honing S, Evans PR & Owen DJ 2010 A large-scale conformational change couples membrane recruitment to cargo binding in the AP2 clathrin adaptor complex. *Cell* **141** 1220–1229. (<https://doi.org/10.1016/j.cell.2010.05.006>)
- Jean-Alphonse F, Bowersox S, Chen S, Beard G, Puthenveedu MA & Hanyaloglu AC 2014 Spatially restricted G protein-coupled receptor activity via divergent endocytic compartments. *Journal of Biological Chemistry* **289** 3960–3977. (<https://doi.org/10.1074/jbc.M113.526350>)
- Jung SY, Kwak JO, Kim HW, Kim DS, Ryu SD, Ko CB & Cha SH 2005 Calcium sensing receptor forms complex with and is up-regulated by caveolin-1 in cultured human osteosarcoma (Saos-2) cells. *Experimental and Molecular Medicine* **37** 91–100. (<https://doi.org/10.1038/emmm.2005.13>)
- Kelly BT, McCoy AJ, Spate K, Miller SE, Evans PR, Honing S & Owen DJ 2008 A structural explanation for the binding of endocytic dileucine motifs by the AP2 complex. *Nature* **456** 976–979. (<https://doi.org/10.1038/nature07422>)
- Kifor O, Diaz R, Butters R, Kifor I & Brown EM 1998 The calcium-sensing receptor is localized in caveolin-rich plasma membrane domains of bovine parathyroid cells. *Journal of Biological Chemistry* **273** 21708–21713. (<https://doi.org/10.1074/jbc.273.34.21708>)
- Kifor O, MacLeod RJ, Diaz R, Bai M, Yamaguchi T, Yao T, Kifor I & Brown EM 2001 Regulation of MAP kinase by calcium-sensing receptor in bovine parathyroid and CaR-transfected HEK293 cells. *American Journal of Physiology: Renal Physiology* **280** F291–F302. (<https://doi.org/10.1152/ajprenal.2001.280.2.F291>)
- Kirchhausen T, Owen D & Harrison SC 2014 Molecular structure, function, and dynamics of clathrin-mediated membrane traffic. *Cold Spring Harbor Perspectives in Biology* **6** a016725. (<https://doi.org/10.1101/cshperspect.a016725>)
- Kotowski SJ, Hopf FW, Seif T, Bonci A & von Zastrow M 2011 Endocytosis promotes rapid dopaminergic signaling. *Neuron* **71** 278–290. (<https://doi.org/10.1016/j.neuron.2011.05.036>)
- Leach K, Wen A, Davey AE, Sexton PM, Conigrave AD & Christopoulos A 2012 Identification of molecular phenotypes and biased signaling induced by naturally occurring mutations of the human calcium-sensing receptor. *Endocrinology* **153** 4304–4316. (<https://doi.org/10.1210/en.2012-1449>)
- Leach K, Wen A, Cook AE, Sexton PM, Conigrave AD & Christopoulos A 2013 Impact of clinically relevant mutations on the pharmacoregulation and signaling bias of the calcium-sensing receptor by positive and negative allosteric modulators. *Endocrinology* **154** 1105–1116. (<https://doi.org/10.1210/en.2012-1887>)
- Li D, Opas EE, Tuluc F, Metzger DL, Hou C, Hakonarson H & Levine MA 2014 Autosomal dominant hypoparathyroidism caused by germline mutation in GNA11: phenotypic and molecular characterization. *Journal of Clinical Endocrinology and Metabolism* **99** E1774–E1783. (<https://doi.org/10.1210/jc.2014-1029>)
- Lorenz S, Frenzel R, Paschke R, Breitwieser GE & Miedlich SU 2007 Functional desensitization of the extracellular calcium-sensing receptor is regulated via distinct mechanisms: role of G protein-coupled receptor kinases, protein kinase C and beta-arrestins. *Endocrinology* **148** 2398–2404. (<https://doi.org/10.1210/en.2006-1035>)
- Mannstadt M, Harris M, Bravenboer B, Chitturi S, Dreierink KM, Lambright DG, Lim ET, Daly MJ, Gabriel S & Juppner H 2013 Germline mutations affecting Galpha11 in hypoparathyroidism. *New England Journal of Medicine* **368** 2532–2534. (<https://doi.org/10.1056/NEJMc1300278>)
- Marx SJ 2015 Letter to the editor: distinguishing typical primary hyperparathyroidism from familial hypocalciuric hypercalcemia by using an index of urinary calcium. *Journal of Clinical Endocrinology and Metabolism* **100** L29–L30. (<https://doi.org/10.1210/jc.2014-4221>)
- Miedlich SU, Gama L, Seuwen K, Wolf RM & Breitwieser GE 2004 Homology modeling of the transmembrane domain of the human calcium sensing receptor and localization of an allosteric binding site. *Journal of Biological Chemistry* **279** 7254–7263. (<https://doi.org/10.1074/jbc.M307191200>)
- Mingione A, Verdelli C, Ferrero S, Vaira V, Guarnieri V, Scillitani A, Vicentini L, Balza G, Beretta E, Terranegra A, et al. 2017 Filamin A is reduced and contributes to the CASR sensitivity in human parathyroid tumors. *Journal of Molecular Endocrinology* **58** 91–103. (<https://doi.org/10.1530/JME-16-0184>)
- Nesbit MA, Hannan FM, Howles SA, Babinsky VN, Head RA, Cranston T, Rust N, Hobbs MR, Heath H 3rd & Thakker RV 2013a Mutations affecting G-protein subunit alpha11 in hypercalcemia and hypocalcemia. *New England Journal of Medicine* **368** 2476–2486. (<https://doi.org/10.1056/NEJMoa1300253>)
- Nesbit MA, Hannan FM, Howles SA, Reed AA, Cranston T, Thakker CE, Gregory L, Rimmer AJ, Rust N, Graham U, et al. 2013b Mutations in AP2S1 cause familial hypocalciuric hypercalcemia type 3. *Nature Genetics* **45** 93–97. (<https://doi.org/10.1038/ng.2492>)
- Obermannova B, Banghova K, Sumnik Z, Dvorakova HM, Betka J, Fencel E, Kolouskova S, Cinek O & Lebl J 2009 Unusually severe phenotype of neonatal primary hyperparathyroidism due to a heterozygous inactivating mutation in the CASR gene. *European Journal of Pediatrics* **168** 569–573. (<https://doi.org/10.1007/s00431-008-0794-y>)
- Pearce SH, Trump D, Wooding C, Besser GM, Chew SL, Grant DB, Heath DA, Hughes IA, Paterson CR, Whyte MP, et al. 1995 Calcium-sensing receptor mutations in familial benign hypercalcemia and neonatal hyperparathyroidism. *Journal of Clinical Investigation* **96** 2683–2692. (<https://doi.org/10.1172/JCI118335>)
- Pearce SH, Bai M, Quinn SJ, Kifor O, Brown EM & Thakker RV 1996a Functional characterization of calcium-sensing receptor mutations expressed in human embryonic kidney cells. *Journal of Clinical Investigation* **98** 1860–1866. (<https://doi.org/10.1172/JCI118987>)
- Pearce SH, Williamson C, Kifor O, Bai M, Coulthard MG, Davies M, Lewis-Barned N, McCredie D, Powell H, Kendall-Taylor P, et al. 1996b A familial syndrome of hypocalcemia with hypercalciuria due to mutations in the calcium-sensing receptor. *New England Journal of Medicine* **335** 1115–1122. (<https://doi.org/10.1056/NEJM199610103351505>)
- Petrel C, Kessler A, Dauban P, Dodd RH, Rognan D & Ruat M 2004 Positive and negative allosteric modulators of the Ca²⁺-sensing receptor interact within overlapping but not identical binding sites in the transmembrane domain. *Journal of Biological Chemistry* **279** 18990–18997. (<https://doi.org/10.1074/jbc.M400724200>)
- Pi M, Spurney RE, Tu QS, Hinson T & Quarles LD 2002 Calcium-sensing receptor activation of Rho involves filamin and Rho-guanine nucleotide exchange factor. *Endocrinology* **143** 3830–3838. (<https://doi.org/10.1210/en.2002-220240>)
- Pi M, Oakley RH, Gesty-Palmer D, Cruickshank RD, Spurney RE, Luttrell LM & Quarles LD 2005 Beta-arrestin- and G protein receptor kinase-mediated calcium-sensing receptor desensitization. *Molecular Endocrinology* **19** 1078–1087. (<https://doi.org/10.1210/me.2004-0450>)
- Pidasheva S, Grant M, Canaff L, Ercan O, Kumar U & Hendy GN 2006 Calcium-sensing receptor dimerizes in the endoplasmic reticulum: biochemical and biophysical characterization of CASR mutants retained intracellularly. *Human Molecular Genetics* **15** 2200–2209. (<https://doi.org/10.1093/hmg/ddl145>)
- Piret SE, Gorvin CM, Pagnamenta AT, Howles SA, Cranston T, Rust N, Nesbit MA, Glaser B, Taylor JC, Buchs AE, et al. 2016 Identification of a G-protein subunit-alpha11 gain-of-function mutation, Val340Met, in a family with autosomal dominant hypocalcemia type 2 (ADH2). *Journal of Bone and Mineral Research* **31** 1207–1214. (<https://doi.org/10.1002/jbmr.2797>)

- Pollak MR, Brown EM, Chou YH, Hebert SC, Marx SJ, Steinmann B, Levi T, Seidman CE & Seidman JG 1993 Mutations in the human Ca(2+)-sensing receptor gene cause familial hypocalciuric hypercalcemia and neonatal severe hyperparathyroidism. *Cell* **75** 1297–1303. ([https://doi.org/10.1016/0092-8674\(93\)90617-Y](https://doi.org/10.1016/0092-8674(93)90617-Y))
- Rappoport JZ & Simon SM 2003 Real-time analysis of clathrin-mediated endocytosis during cell migration. *Journal of Cell Science* **116** 847–855. (<https://doi.org/10.1242/jcs.00289>)
- Rasmussen SG, Choi HJ, Rosenbaum DM, Kobilka TS, Thian FS, Edwards PC, Burghammer M, Ratnala VR, Sanishvili R, Fischetti RE, *et al.* 2007 Crystal structure of the human beta2 adrenergic G-protein-coupled receptor. *Nature* **450** 383–387. (<https://doi.org/10.1038/nature06325>)
- Riccardi D & Brown EM 2010 Physiology and pathophysiology of the calcium-sensing receptor in the kidney. *American Journal of Physiology: Renal Physiology* **298** F485–F499. (<https://doi.org/10.1152/ajprenal.00608.2009>)
- Schrage R, Schmitz AL, Gaffal E, Annala S, Kehraus S, Wenzel D, Bullesbach KM, Bald T, Inoue A, Shinjo Y, *et al.* 2015 The experimental power of FR900359 to study Gq-regulated biological processes. *Nature Communications* **6** 10156. (<https://doi.org/10.1038/ncomms10156>)
- Shukla AK, Xiao K & Lefkowitz RJ 2011 Emerging paradigms of beta-arrestin-dependent seven transmembrane receptor signaling. *Trends in Biochemical Sciences* **36** 457–469. (<https://doi.org/10.1016/j.tibs.2011.06.003>)
- Shukla AK, Westfield GH, Xiao K, Reis RI, Huang LY, Tripathi-Shukla P, Qian J, Li S, Blanc A, Oleskie AN, *et al.* 2014 Visualization of arrestin recruitment by a G-protein-coupled receptor. *Nature* **512** 218–222. (<https://doi.org/10.1038/nature13430>)
- Standfuss J, Edwards PC, D'Antona A, Fransen M, Xie G, Oprian DD & Schertler GF 2011 The structural basis of agonist-induced activation in constitutively active rhodopsin. *Nature* **471** 656–660. (<https://doi.org/10.1038/nature09795>)
- Sun J & Murphy E 2010 Calcium-sensing receptor: a sensor and mediator of ischemic preconditioning in the heart. *American Journal of Physiology: Heart and Circulatory Physiology* **299** H1309–H1317. (<https://doi.org/10.1152/ajpheart.00373.2010>)
- Takasaki J, Saito T, Taniguchi M, Kawasaki T, Moritani Y, Hayashi K & Kobori M 2004 A novel Galphaq/11-selective inhibitor. *Journal of Biological Chemistry* **279** 47438–47445. (<https://doi.org/10.1074/jbc.M408846200>)
- Thomsen AR, Hvidtfeldt M & Brauner-Osborne H 2012 Biased agonism of the calcium-sensing receptor. *Cell Calcium* **51** 107–116. (<https://doi.org/10.1016/j.ceca.2011.11.009>)
- Uckun-Kitapci A, Underwood LE, Zhang J & Moats-Staats B 2005 A novel mutation (E767K) in the second extracellular loop of the calcium sensing receptor in a family with autosomal dominant hypocalcemia. *American Journal of Medical Genetics Part A* **132A** 125–129. (<https://doi.org/10.1002/ajmg.a.30403>)
- Vargas-Poussou R, Huang C, Hulin P, Houillier P, Jeunemaitre X, Paillard M, Planelles G, Dechaux M, Miller RT & Antignac C 2002 Functional characterization of a calcium-sensing receptor mutation in severe autosomal dominant hypocalcemia with a Bartter-like syndrome. *Journal of the American Society of Nephrology* **13** 2259–2266. (<https://doi.org/10.1097/01.ASN.0000025781.16723.68>)
- Volpe A, Guerriero A, Marchetta A, Caramaschi P & Furlani L 2009 Familial hypocalciuric hypercalcemia revealed by chondrocalcinosis. *Joint Bone Spine* **76** 708–710. (<https://doi.org/10.1016/j.jbspin.2009.02.001>)
- Ward DT, Brown EM & Harris HW 1998 Disulfide bonds in the extracellular calcium-polyvalent cation-sensing receptor correlate with dimer formation and its response to divalent cations in vitro. *Journal of Biological Chemistry* **273** 14476–14483. (<https://doi.org/10.1074/jbc.273.23.14476>)
- Watanabe S, Fukumoto S, Chang H, Takeuchi Y, Hasegawa Y, Okazaki R, Chikatsu N & Fujita T 2002 Association between activating mutations of calcium-sensing receptor and Bartter's syndrome. *Lancet* **360** 692–694. ([https://doi.org/10.1016/S0140-6736\(02\)09842-2](https://doi.org/10.1016/S0140-6736(02)09842-2))
- Wehbi VL, Stevenson HP, Feinstein TN, Calero G, Romero G & Vilardaga JP 2013 Noncanonical GPCR signaling arising from a PTH receptor-arrestin-Gbetagamma complex. *PNAS* **110** 1530–1535. (<https://doi.org/10.1073/pnas.1205756110>)
- Wu H, Wang C, Gregory KJ, Han GW, Cho HP, Xia Y, Niswender CM, Katritch V, Meiler J, Cherezov V, *et al.* 2014 Structure of a class C GPCR metabotropic glutamate receptor 1 bound to an allosteric modulator. *Science* **344** 58–64. (<https://doi.org/10.1126/science.1249489>)
- Zhang C, Zhang T, Zou J, Miller CL, Gorkhali R, Yang JY, Schillmiller A, Wang S, Huang K, Brown EM, *et al.* 2016 Structural basis for regulation of human calcium-sensing receptor by magnesium ions and an unexpected tryptophan derivative co-agonist. *Science Advances* **2** e1600241. (<https://doi.org/10.1126/sciadv.1600241>)

Received in final form 16 March 2018

Accepted 29 March 2018

Accepted Preprint published online 29 March 2018