THEMATIC REVIEW

SULFATION PATHWAYS

Sources and biological activities of marine sulfated steroids

Francisca Carvalho^{1,2}, Marta Correia-da-Silva^{1,2}, Emília Sousa^{1,2}, Madalena Pinto^{1,2} and Anake Kijjoa^{2,3}

¹Laboratory of Organic and Pharmaceutical Chemistry, Department of Chemical Sciences, Faculty of Pharmacy, University of Porto, Porto, Portugal

²Interdisciplinary Centre of Marine and Environmental Research (CIIMAR), Terminal de Cruzeiros do Porto de Leixões, Matosinhos, Portugal

³ICBAS-Instituto de Ciências Biomédicas Abel Salazar, University of Porto, Porto, Portugal

Correspondence should be addressed to M Correia-da-Silva: m_correiasilva@ff.up.pt

This paper is part of a thematic section on Sulfation Pathways. The guest editors for this section were Jonathan Wolf Mueller and Paul Foster.

Abstract

Marine environment is rich in structurally unique molecules and can be an inspiring source of novel drugs. Currently, six marine-derived drugs are in the market with FDA approval and several more are in the clinical pipeline. Structurally diverse and complex secondary metabolites have been isolated from the marine world and these include sulfated steroids. Biological activities of nearly 150 marine sulfated steroids reported from 1978 to 2017 are compiled and described, namely antimicrobial, antitumor, cardiovascular and antifouling activities. Structure–activity relationship for each activity is discussed.

Key Words

- ▶ marine
- ▶ sulfates
- ▶ steroids
- ▶ antitumor
- ▶ antimicrobial

*Journal of Molecular
Endocrinology*
(2018) **61**, T211–T231

Introduction

The marine biota represents one of the most unexplored extreme environments in the world. Oceans cover about 70% of the Earth's surface (Giddings & Newman 2015) and approximately one half of the Earth's species is present in the marine world (Glaser & Mayer 2009). Marine biota has a higher taxonomic diversity composition than its terrestrial counterpart (Appeltans *et al.* 2012, Pimm 2012), which combined with its unique environmental conditions and organisms, offers an enormous resource of novel compounds for industrial development such as pharmaceuticals, food ingredients, cosmeceuticals, drug delivery systems and industrial enzymes (Martins *et al.* 2014).

In the past 50 years, progressive improvements have been made in exploration of new marine habitats, leading to the isolation of thousands of unique marine natural

products (Martins *et al.* 2014). Currently, there are six U.S. Food and Drug Administration (FDA)-approved marine-derived drugs in the market and several compounds in different phases of the clinical trials (Mayer *et al.* 2017). The first marine drug launched in the market was a synthetic analog of the natural spongothymidine nucleoside originally isolated from a Caribbean marine sponge (Cytosar-U, FDA approved for cancer in 1969). Following, in 2004, omega-3 fatty acids, isolated from fish oils (Lovaza), granted the FDA approval as an anti-hypertriglyceridemia as well as a synthetic derivative (Prialt) inspired by a naturally occurring peptide, isolated from the venom of the cone snail *Conus magus*, for the management of severe chronic pain associated with cancer, AIDS and neuropathies. More recently, FDA has approved a macrolide (Halaven, 2010), derived from

the sponge *Halichondria okadai*, for metastatic breast cancer; an antibody–drug conjugate (Adcetris, 2011), isolated from the sea hare *Dolabella auricularia*, for use in Hodgkin’s lymphoma and anaplastic large cell lymphoma and an alkaloid (Yondelis, 2015), isolated from a tunicate, for ovarian cancer and soft tissue sarcoma (Mayer *et al.* 2010, Jaspars *et al.* 2016).

Within the marine-derived compounds which are in clinical trials, a marine steroid (squalamine, Fig. 1) is in phase III of drug development. Squalamine is a sulfated aminosterol derived from the internal organs (primarily the liver) of the dogfish shark and although was first described as an antimicrobial agent against fungi, protozoa and gram-negative and gram-positive bacteria (Moore *et al.* 1993) is now recognized as an anti-angiogenic drug with a novel intracellular mechanism of action. The drug acts against the development of aberrant neovascularization by inhibiting multiple growth factors, including vascular endothelial growth factor (VEGF) (Connolly *et al.* 2006). Clinical evidence has shown that these growth factors play a role in angiogenesis and ocular neovascular disease. OHR Pharmaceuticals Inc. evaluated the safety and effectiveness of squalamine for the treatment of neovascular age-related macular degeneration (IMPACT phase II study), showing an improvement in visual function when used in combination with an anti-VEGF agent (<https://www.ohrpharmaceutical.com/media-center/press-releases/detail/446/ohr-pharmaceutical-presents-data-from-ohr-102-phase-ii>, October 28, 2017). Presently, a phase III trial of squalamine is underway (NCT02727881, <https://clinicaltrials.gov/ct2/show/NCT02727881>, October 28, 2017). Squalamine has been targeted of several reviews highlighting its potential as a drug candidate (Abd *et al.* 2017, Hussain & Ciulla 2017), which puts forward the potential of sterol sulfates derived from marine organisms as models for development of new drugs.

Sulfur, in the form of sulfate salt, is the third most abundant element in seawater (Pinet 2011). So, it is not a surprise that sulfated metabolites are extensively distributed in marine organisms (Kornprobst *et al.*

1998). Sulfated compounds are found in a wide variety of organisms, from prokaryotes to multicellular species, being flavonoids the phenolic small molecules with more sulfated derivatives described from plants and seagrass (Correia-da-Silva *et al.* 2014). In marine biota, several sulfated secondary metabolites from different chemical classes (e.g., terpenoids, carotenoids, steroids, alkaloids, etc.) have been identified (Kornprobst *et al.* 1998). Sulfated steroids constitute one of the most numerous classes of these secondary metabolites; however, their biological functions are still poorly understood (Stonik 2001). Some of them are known for their use in chemical protection against predators (Morinaka *et al.* 2009), pathogenic (Smyrniotopoulos *et al.* 2015) or fouling organisms (Iorizzi *et al.* 1995). Other marine products, structurally close to bile acids and alcohols of higher animals, were described to facilitate food absorption and digestion in some invertebrates (Levina *et al.* 2004). Some sulfated steroids were also found to inhibit the digestive enzymes of marine molluscs, *endo*-1,3- β -D-glucanases, therefore functioning as a chemical defence mechanism (Zvyagintseva *et al.* 1986, Makarieva *et al.* 1995, Guzii *et al.* 2008). While regulation of sulfation and desulfation are vital processes for vertebrate endocrine system (Mueller *et al.* 2015), little is known concerning the influence of sulfated steroids on endocrine system of marine organisms. Considering that sulfated steroid derivatives can act by interaction with nuclear receptors (genomic pathways) or membrane surface (non-genomic) (Falkenstein *et al.* 2000, Fiorucci *et al.* 2012), it is possible to explain the interference with endocrine systems (Mueller *et al.* 2015). It is interesting to add that some sulfated steroids were found as pheromones in marine species (Fine & Sorensen 2005, Stacey 2015). The hypothesis of intervention at the endocrine level may also be corroborated if we take into account the intervention of some sulfated marine steroids in hormone-dependent human tumor cell lines (Malvarenko *et al.* 2015).

The investigation of biological/pharmacological activities of these marine-derived sulfated steroids can lead to novel structures for the treatment of important pathologies. This review covers the biological activities of 142 marine sulfated steroids reported from 1978 to 2017. The majority of reported marine sulfated steroids are secondary metabolites of sterols, a subgroup of steroid compounds, which present a C-3 hydroxyl group and an aliphatic side chain at C-17 position additionally to the cyclopentane perhydrophenanthrene core (Moss 1989). The most important biological activities found in sulfated steroids can be categorized as antimicrobial, antitumor

Figure 1
Structure of squalamine.

and cardiovascular. However, other activities were also reported, namely antifouling, showing their potential for different applications other than therapeutic drugs. Some structure–activities relationships (SAR) were also tentatively established.

Therapeutic applications

Antimicrobial activity

Antibacterial

Several reasons contribute to the imperative need for new antibiotics. The fact that infectious diseases still represent one of the greatest threats to human health worldwide is due to the rapid development of resistance by bacteria to therapeutic agents available (Spellberg et al. 2011, Butler et al. 2017).

Thirteen marine sulfated sterols (**1–13**, Fig. 2), mostly isolated from sponges, were shown to have antibacterial activity. Sponges are sessile organisms that endure over the years, due to their capability to engaging in chemical battles for defence or food competition against other animals, making them a reservoir of diverse and unique compounds. The study of their metabolites is of great importance to the discovery of new solutions to combat the development resistance by bacteria.

Carboxylated sterol 3 β -sulfates (compounds **1–3**), isolated from the marine sponge *Toxadocia zumi*, showed antimicrobial activity against *Bacillus subtilis* and *Staphylococcus aureus* at 100 μ g/disk and 50 μ g/disk, respectively (Nakatsu et al. 1983), whereas the respective 3 β -hydroxy sterols were not active, suggesting that 3 β -sulfate group was necessary for the antibacterial activity (Nakatsu et al. 1983).

Compound **4**, isolated from the starfish *Luidia clathrata*, also inhibited the growth of both bacterial

species (*B. subtilis* and *S. aureus*) at 50 μ g/disk (Iorizzi et al. 1995). The monosulfate polymastiamide A (**5**), isolated from the marine sponge *Polymastia boletiformis*, was the first example of a marine natural product with an uncommon side chain containing an amide bond linking the steroid part to a non-proteinaceous amino acid. Compound **5** showed an *in vitro* activity against *S. aureus* at 100 μ g/disk (Kong & Andersen 1993).

Two disulfated sterols (**6** and **7**), isolated from the brittle star *Ophioderma longicauda* (Riccio et al. 1985), were tested for their activity against *S. aureus*, and the results were compared with other sulfated and non-sulfated sterols. While non-sulfated sterols exhibited inhibitory activity at 200 μ g/disk, sulfated sterols **6** and **7** were much more potent against *S. aureus* (20 μ g/disk) (Andersson et al. 1989).

Fusetani and coworkers first described isolation of halistanol trisulfate (**8**) from the marine sponge *Halichondria* cf. *moorei* Bergquist and have found that this compound was able to inhibit the growth of gram-positive and gram-negative bacteria. However, the acid hydrolysis product of **8** exhibited no antimicrobial activity showing that the sulfate groups were necessary for this activity (Fusetani et al. 1981). More recently, **8** was also found to have a bactericidal activity against *S. aureus*, with a minimum inhibitory activity (MIC) value of 512 μ g/mL (Marinho et al. 2012). Moreover, compound **8** not only exhibited antibacterial activity against *Streptococcus mutans* CI (clinical isolated from an active caries) and *S. mutans* UA159 strains at 3.0 μ g/mL, but also inhibited the biofilm formation by these strains approximately 85 and 99%, respectively (Lima et al. 2014). From the sponge *Topsentia* sp., five trisulfated sterols, topsentiasterol sulfates A–E (**9–13**), were isolated and shown to be active against *Pseudomonas aeruginosa* and *Escherichia coli* at 10 μ g/disk (Fusetani et al. 1994). Compounds **9–12** were

Figure 2
Antibacterial sulfated steroids.

the first examples of marine sulfated sterols possessing a butenolide or a furan functionality at the end of the side chain (Fusetani *et al.* 1994).

From the aforementioned 13 antibacterial sulfated sterols, it is possible to observe that six were trisulfated sterols, five were monosulfated and only two were disulfated sterols. Some studies have highlighted the sulfate groups as crucial groups for the activity (Fusetani *et al.* 1981, Nakatsu *et al.* 1983, Andersson *et al.* 1989).

Antifungal

There are four major classes of antifungal agents available for clinicians today: the allylamines, the azoles, the echinocandins and the polyenes (Lockhart & Warnock 2015). Due to the host toxicity and interactions with other drugs of the currently used antifungal agents, and the significant morbidity and mortality associated with invasive fungal infections in immune-compromised patients, new agents with different mechanism of action and with extended activity against pathogens are needed (Perfect 2016).

Antifungal activity was described for 26 sulfated steroids (**5**, **8**, **12–35**, Fig. 3), namely against *Candida albicans*, *Mortierella remannianus*, *Saccharomyces cerevisiae*, *Aspergillus flavus* and marine/plant fungal pathogens (*Pythium ultimum*, *Cladosporium cucumerinum*, *Pyricularia oryzae*, *Lindra thallasiae*).

Several sulfated sterols such as **5**, **8**, **12–16** have the ability to inhibit *Candida* sp. Compound **5** and another monosulfated sterol conjugated with an amino acid, **14**, exhibited inhibitory activity against *C. albicans* at 75 µg/disk (Kong & Andersen 1993) and at 100 µg/disk (Smyrniotopoulos *et al.* 2015), respectively. Topsentasterol trisulfates D and E (**12** and **13**, Fig. 3) also showed activity against *C. albicans* at 10 µg/disk (Fusetani *et al.* 1994), while **8** exhibited activity against two strains of *Candida* sp., i.e. sensitive (*C. parapsilosis*) and azole resistant (*C. krusei*) to antibiotics belonging to the class of azoles. Halistanol trisulfate (**8**) was more active against *C. krusei* (IC₅₀ of 26 µg/mL) than *C. parapsilosis* (IC₅₀ of 60 µg/mL) (Kossuga *et al.* 2007). Compounds **12** and **13** also showed antifungal activity against *M. remannianus* (at 10 µg/disk) (Fusetani *et al.* 1994). Eurysterols A and B (**15** and **16**, Fig. 3), two monosulfated sterols isolated from a marine sponge of the genus *Euryspongia*, were tested for their inhibitory activity against amphotericin B-resistant strains of *C. albicans*. Despite their structural similarities, eurysterol A (**15**) was more active (MIC value of 16 µg/mL)

than eurysterol B (**16**) (MIC value of 63 µg/mL), allowing to postulate that the presence of the double bond in **16** caused a decrease of the activity when compared to **15** (Boonlarpradab & Faulkner 2007).

Some sulfated steroids, including the monosulfates **17–22** and the trisulfate **23**, also showed growth inhibitory activity against the yeast *Saccharomyces cerevisiae*: anti-yeast tests were carried out for acanthosterol I (**17**) and acanthosterol J (**18**) and both compounds showed growth inhibitory activity against *S. cerevisiae* strain A364A and its mutants (Tsukamoto *et al.* 1998). Compound **17** had an inhibitory zone of 7 mm against the strain A364A while **18** showed 11 mm effect (paper disks impregnated with 0.1 mg of each compound), and both of them showed higher inhibitory zone against the mutant than the wild-type strains (Tsukamoto *et al.* 1998). Four sulfated steroids were examined for their fluconazole reversal activity by a checkerboard assay that measures the combination treatment effect of two agents on a microbe, and the fractional inhibitory concentration (FIC) value was used to define the effect as synergistic (≤ 0.5), additive (0.51–1.0), indifferent (1.1–2.0) or antagonist (< 2.0). Two *S. cerevisiae* strains that overexpress *C. albicans* CDR1 (DSY 415) and MDR1 (DSY 415) efflux pumps were used in tests with capisterones A (**19**) and B (**20**), that significantly enhanced fluconazole activity in the CDR1 and MDR1 efflux pump-overexpressing strains, with FIC values of 0.33 and 0.08 for **19** and 0.33 and 0.15 for **20** (Li *et al.* 2006). Geodisterol-3-O-sulfite (**21**) and 29-demethylgeodisterol-3-O-sulfite (**22**), the first described naturally occurring sulfated sterols with an aromatic ring system, were isolated from the marine sponge *Topsentia* sp. These compounds were also found to enhance the activity of fluconazole in a *S. cerevisiae* strain overexpressing the *C. albicans* efflux pump MDR1 (FIC values of 0.08 and 0.15, respectively), as well as in a fluconazole-resistant *C. albicans* clinical isolate known to overexpress MDR1 (FIC values of 0.2 for both compounds) (DiGirolamo *et al.* 2009). A trisulfated sterol having a rare side chain with a diethyl substitution at C-24, Sch 575867 (**23**), isolated from a marine sponge belonging to the family Astroscleridae, was found to exhibit the antifungal activity with MIC value of 15 µg/mL against a modified strain of the wild-type *S. cerevisiae*, PM503 (Yang *et al.* 2003a).

Three glycosylated sterols, minutosides A (**24**) and B (**25**), and pycnopodioside (**26**), were active against *A. flavus* with inhibition zones of 5–10 mm at the highest concentration of 20–60 µg/disk. Compound **25**, the first example of a natural steroidal xyloside containing an amide function in the aglycone, showed activity only at

Figure 3
Antifungal sulfated steroids.

the highest concentration (20–60 µg/disk), while **24** and **26** exhibited activity at 20–60 µg/disk (inhibition zones of 5–7.5 mm) (Chludil & Maier 2005). The desulfated analogs of **24** and **26** were inactive against *A. flavus* at all tested concentrations, allowing to conclude that the presence of a sulfate group in the aglycone moiety could play an important role in the antifungal activity against *A. flavus* (Chludil & Maier 2005).

The growth of four plant pathogenic fungi, *P. ultimum*, *C. cucumerinum*, *P. oryzae* and the marine-derived *L. thallasiae*, was inhibited by a few sulfated sterols. Compound **5** exhibited *in vitro* activity against *P. ultimum* at 25 µg/disk (Kong & Andersen 1993). The two amino acid-conjugated sterols **14** and **27**, isolated from the sponge *Polymastia boletiformis*, showed inhibition zones of 10 and 8 mm at 30 and 60 µg/disk, respectively, against *C. cucumerinum* (Smyrniotopoulos et al. 2015). Several glycosylated sterol sulfates were also reported with antifungal activity against *C. cucumerinum*. Anasteroside A (**28**, Fig. 3), an asterosaponin (a 3 β ,6 α -dihydroxysteroid with a $\Delta^{9(11)}$ unsaturation, a sulfate group at C-3, and a carbohydrate chain with four to six monosaccharide units attached to C-6) which was isolated from the starfish *Anasterias minuta*, showed an inhibition zone of approximately 12 mm (at 10 µg/disk) against *C. cucumerinum*, while anasteroside B, with a shorter side chain and lack of the hydroxyl group at C-20, was inactive at all tested concentrations (Chludil et al. 2002). To evaluate the influence of the oligosaccharide moiety in the antifungal activity, versicoside A (**29**), which is structurally similar to anasterosides A and B, but with a different side chain, was enzymatically hydrolyzed to the triglycoside forberoside H and the pentaglycoside thornasteroside A (**30**). While the hexaglycosylated sterol **29** and the pentaglycosylated sterol **30** were active (inhibition zone of approximately 8 mm and 5 mm at 10 µg/disk, respectively), the triglycosylated sterol forberoside H, showed no antifungal activity in all concentration ranges (Chludil et al. 2002). Desulfation of versicoside A (**29**) by solvolysis afforded an inactive saponin. The results obtained in this work suggested that the side chain in the steroidal aglycone moiety, the structure of the sugar portion and the presence of a sulfate group at C-3, might play an important role in the antifungal activity of these sulfated steroidal saponins (Chludil et al. 2002). Other three sulfated glycosylated compounds, isolated from the starfish *A. minuta*, were studied. Compounds **24** and **26** exhibited antifungal activity against *C. cucumerinum* (inhibition zones of 7–10 mm at 10–60 µg/disk) (Chludil & Maier 2005),

and compound **25** showed activity only at the highest concentration (20–60 µg/disk) against *C. cucumerinum* (inhibition zones of 3–4 mm) (Chludil & Maier 2005). The desulfated analogs of **24** and **26** were inactive against *C. cucumerinum* at all tested concentrations, allowing to conclude that the presence of a sulfate group in the aglycone moiety could play an important role in the antifungal activity against *C. cucumerinum* (Chludil & Maier 2005).

The plant pathogenic fungus *P. oryzae* has been used as a test for the primary screening of antineoplastic and antifungal agents (Tang et al. 2005a), and five asterosaponins isolated from the starfish *Culcita novaeguineae* exhibited antifungal activity in this screening. Asterosaponins **31** and **32** showed significant antifungal inhibitory activity with minimum morphological deformation concentration (MMDC) values of 8 and 4 µg/mL, respectively (Tang et al. 2005a). Thornasteroside A (**30**), asterosaponin **33**, marthasteroside A₁ (**34**) and regularoside A (**35**) exhibited moderate activity inducing morphological deformation of *P. oryzae* mycelia with MMDC values of 64, 16, 32 and 64 µg/mL, respectively (Tang et al. 2006).

Compounds **19** and **20**, isolated from the tropical green alga *Penicillus capitatus*, showed potent inhibitory activity against the marine fungal pathogen *L. thallasiae*, with median lethal dose (LD₅₀) concentrations of 0.03 and 0.94 µg/mL, respectively (Puglisi et al. 2004).

From the 26 antifungal sulfated steroids, 13 were isolated from sponges, 11 from starfishes and only two were obtained from a marine alga. It is possible to observe that almost all sulfated steroids with antifungal activity are monosulfated with a sulfate group in position C-3 (mainly with β -configuration) and almost half of them are glycosylated (eight asterosaponins and three monoglycosides). It is noteworthy to mention that none of the glycosylated steroids have the sulfate group on the glycosidic portion.

These antifungal studies were mostly *in vitro* preliminary tests and only few used resistance models. Overall, these studies put forward the potential of sulfated steroids as antifungal agents.

Antiviral

Several sulfated flavonoids and other polyphenols have been highlighted for their antiviral activity (Sousa et al. 2008, Correia-da-Silva et al. 2014). Twenty-five marine sulfated sterols (**8**, **36–59**, Fig. 4) were studied against several viruses: human immunodeficiency virus (HIV-1 and HIV-2), feline leukemia virus (FeLV), polio virus

Figure 4
Antiviral sulfated steroids.

(PV and PV-3), junin virus (JV), respiratory syncytial virus (RSV) and herpes simplex virus (HSV-1 and HSV-2), as discussed below.

Weinbersterol disulfate A (**36**), which was isolated from the marine sponge *Petrosia weinbergi*, exhibited HIV-inhibitory activity with half maximum effective

concentration (EC₅₀) value around 1.0 µg/mL (Sun *et al.* 1991). Protection against the cytopathic effects of HIV-1 infection was evaluated in the National Cancer Institute (NCI) primary screen for several sulfated sterols, namely: sterols with sulfates on A and B rings; sterols with oxygen or sulfate substituent on the D ring and sterols with

sulfation at C-21 (McKee et al. 1994). Promising activity was obtained with four trisulfated sterols with a common 2α , 3β and 6α -trisulfate substitution pattern (McKee et al. 1994), i.e. compound **8** ($EC_{50}=6\mu\text{M}$), ibisterol trisulfate (**37**, $EC_{50}=13\mu\text{M}$) isolated from the Caribbean sponge *Topsentia* sp. (McKee et al. 1993), halistanol trisulfate F (**38**, $EC_{50}=3\mu\text{M}$), and halistanol trisulfate G (**39**, $EC_{50}=6\mu\text{M}$), isolated from the marine sponge *Pseudoaxinissa digitata* (Bifulco et al. 1994). Compounds **8**, **37**, **38** and **39** were also tested against a strain of HIV-2, which generally showed up to 70–80% protection (McKee et al. 1994). Cytopathic effects of HIV-1 infection were also obtained for four disulfated sterols (**40–43**), which were isolated from *Tremaster novaecaledoniae*, and have 3β and 6α substitution pattern and additional sites of oxygenation located on the sterol side chain. Compounds **40–43** exhibited antifungal activity with EC_{50} between 13 and $48\mu\text{M}$ (McKee et al. 1994). It is interesting to note that sterols with sulfate groups in the A and B rings were the most active, whereas sterols with oxygenation in the D ring did not showed antifungal activity (McKee et al. 1994).

Haplosamates A (**44**) and B (**45**), isolated from a Philippine sponge *Xestospongia* sp., represent the first examples of marine sterols with the sulfamate functionality and an unprecedented six-membered ether ring. Compounds **44** and **45** inhibited HIV type 1 integrase with IC_{50} values (Qureshi & Faulkner 1999) higher ($50\mu\text{g/mL}$ and $15\mu\text{g/mL}$, respectively) than those of ibisterol sulfates B (**46**) and C (**47**) ($2.3\mu\text{g/mL}$ and $1.8\mu\text{g/mL}$, respectively), also isolated from the same sponge (Lerch & Faulkner 2001).

Clathsterol (**48**), isolated from the Red Sea sponge *Clathria* sp., was found to inhibit an activity against HIV-1 reverse transcriptase at $10\mu\text{g/mL}$ (Rudi et al. 2001).

Two hydroxylated sterols disulfates, weinbersterol disulfates A (**36**) and B (**49**), which were isolated from the sponge *Petrosia weinbergi*, showed an *in vitro* activity against FeLV (EC_{50} of 4.0 and $5.2\mu\text{g/mL}$ respectively). The cyclopropane-containing side chain of these compounds shows the side chain diversity of the sterols from sponges (Sun et al. 1991).

Compounds **50–53**, which only differ in the side chain, were isolated from brittle stars (D'Auria et al. 1987, Roccatagliata et al. 1996) and were tested for their inhibitory effects on the replication of RNA (PV, JV, RSV) and DNA (HSV-1) viruses (Roccatagliata et al. 1996). Compounds **50** and **52**, with a Δ^{22} double bond, showed a marked RSV inhibitory activity (75% inhibition of RSV replication at $40\mu\text{g/mL}$) and were also active against PV (65% reduction of the cytopathic effect at

$68\mu\text{g/mL}$); however, both exhibited weaker activity against the other viruses investigated (<50% reduction of virus plaque formation). Compound **53**, with $\Delta^{24(28)}$ double bond, inhibited 54% of JV replication, but was less active against the other viruses (<50% reduction of virus plaque formation). On the contrary, **51** was weakly active to all the viruses tested (Roccatagliata et al. 1996). Even though HSV-1 was described as the less sensitive virus to these sulfated sterols (Roccatagliata et al. 1996), four sulfated sterols (**8**, **54**, **55**, **56**) were recently tested against HSV-1. Monosulfated sterol **54**, extracted from the China sea starfish *Asterina pectinifera* (Peng et al. 2010), and the polyhydroxylated sterol asterosaponin P2 (**55**), with the sulfate group only in the side chain, isolated from the Far-Eastern starfish *Patiria (Asterina) pectinifera* (Kicha et al. 2000), exhibited activity against HSV-1, with MIC values of 0.2 and $0.07\mu\text{M}$, respectively (Peng et al. 2010). Similarly, compound **8** and halistanol trisulfate C (**56**) were tested for their activity against HSV-1 (KOS strain) and both showed activity with IC_{50} values of 5.63 and $6.09\mu\text{g/mL}$, respectively (Guimaraes et al. 2013).

Three sulfated sterols (**57–59**) found in the Antarctic brittle star *Astrotoma agassizii* (Roccatagliata et al. 1998), were tested against three pathogenic viruses of humans, HSV-2, JV and PV-3. Compounds **58** and **59** exhibited marked HSV-2 inhibitory activity (IC_{50} values of 18.4 and $24.3\mu\text{g/mL}$), while **57** was active at the highest tested concentration ($80\mu\text{g/mL}$) with an IC_{50} of $51.5\mu\text{g/mL}$. However, only **59** presented inhibitory activity (IC_{50} of $67.4\mu\text{g/mL}$) toward JV virus, while **58** was only active against PV-3 at $80\mu\text{g/mL}$ (IC_{50} of $58.5\mu\text{g/mL}$). Therefore, these researchers concluded that sterols with sulfate groups at C-21 and C-2/C-3 were very effective against HSV-2, PV-3 and JV viruses (Comin et al. 1999).

From the 25 compounds with the antiviral activity, more than half of them were disulfated. It is noteworthy to observe that disulfated sterols with the sulfate groups in C-3 and C-21 positions (**50–53**, and **59**) were active against PV viruses while almost all trisulfated sterols (**8**, **37–39**) possessed anti-HIV-1 activity. Moreover, it is interesting to highlight that the majority of the marine antiviral sulfated sterols were isolated from sponges, and only a few are from brittle stars and starfishes.

Anti-parasitic

Halistanol trisulfate **8** (Fig. 2) previously described for its antibacterial activity, was shown to inhibit adenosyl phosphoribosyl transferase in a dose-dependent manner, with 92% inhibition at $25\mu\text{g/mL}$ (IC_{50} of $2.87\mu\text{g/mL}$).

The inhibitory activity of the desulfated analog of **8** was also tested, but no activity was observed. To study the influence of sulfate counter-cations, different sulfate salts of Na⁺, K⁺ and Mg²⁺ were submitted to the assay, and no activity was exhibited. Therefore, the presence of sulfate groups and their spatial position were proved to be essential for the enzymatic inhibitory activity (Kossuga et al. 2007).

Antitumor activity

Nearly 80 marine sulfated steroids with potential antitumor activity have been reported and are compiled in [Supplementary Table 1](#) (see section on [supplementary data](#) given at the end of this article) in chronological order ([Supplementary data](#)). Among these, 52 marine sulfated steroids were investigated as potential antitumor agents after 2000, whereas only 23 were described before the year 2000 ([Supplementary Table 1](#)). Most part of the studies focused on the growth inhibitory effects of the tumor cells, namely against BEL-7402, K-562, Hep-G2, SK-Mel2, KB, PC-3, SNU-C5, RPMI-795, HCT-116, HT-29, Hela, QGY-770 and JURCAT cancer cell lines. It is interesting to mention that molecular targets/mechanisms have been explored and disclosed mainly in the last decade and will be following presented.

Apoptosis

The monosulfate glycosylated sterol, leptaochotensoside A (**60**, [Fig. 5A](#)), isolated from the starfish *Leptasterias ochotensis* (Malyarenko et al. 2015), was found to significantly reduce the colony formation of breast cancer T-47D cells using a soft agar method with 48% inhibition of the colony formation at 50 µM. This compound also reduced the epidermal growth factor (EGF)-induced colony formation of mouse epidermal JB6 Cl41 cells up to 44% relative to control (200 µM) (Malyarenko et al. 2015). The cancer preventive action of **60** was investigated through a non-genomic mechanism of regulation of mitogen-activated protein kinase (MAPK) signaling pathway and was showed to inhibit effectively the EGF-induced phosphorylation of extracellular signal-regulated protein (ERK1/2) and mitogen- and stress-activated protein (MSK-1) kinases. MAPK pathways are evolutionarily conserved kinase modules that link extracellular signals to the machinery that controls fundamental cellular processes such as growth, proliferation, differentiation, migration and apoptosis (Malyarenko et al. 2015).

Astrosterioside D (**61**, [Fig. 5A](#)), an asterosaponin with two ketones in the side chain, isolated from the edible Vietnamese starfish *Astropecten monacanthus* (Thao et al. 2013), was shown to induce apoptosis of HL-60, PC-3 and SNU-C5 cells via the inactivation of phosphatidylinositol 3-kinase (PI3K)/AKT and ERK 1/2 MAPK pathways and the downregulation of C-myc (Thao et al. 2014).

Asterosaponin novaeguinoside II (**62**, [Fig. 5A](#)), which contains an epoxide at C-22 in the side chain, was isolated from the starfish *Culcita novaeguineae* (Tang et al. 2005b). This compound was shown to induce apoptosis of human glioblastoma U87MG cells by the mitochondrial apoptotic pathway (Zhou et al. 2011).

Antitumor effects of **31** ([Fig. 5A](#)) against U87MG cells may result from interfering with cell cycle progression and inducing apoptosis, possibly by decreasing Bcl-2 protein expression (Cheng et al. 2006). Moreover, it was also found that **31** inhibited the proliferation of A549 human lung cancer cells through induction of endoplasmic reticulum stress-associated apoptosis (Zhao et al. 2011).

Asterosaponin archasterosides B (**63**, [Fig. 5A](#)), isolated from the Vietnamese starfish *Archaster typicus*, exhibited cytotoxic activity against HeLa and JB6 P+ Cl41 cell lines. This compound was also shown to induce basal AP-1- and p53-transcriptional activations in JB6 Cl41 (Kicha et al. 2010).

Coscinasterioside B (**64**, [Fig. 5A](#)), a glycosylated sterol monosulfate isolated from the starfish *Coscinasterias tenuispina* (Riccio et al. 1986), was investigated for its p53 inhibitory activity in a yeast two-hybrid test system. This compound was found to inhibit p53 up to 59% relative to control (Levina et al. 2010). Gamma irradiation and anticancer therapy induce p53-dependent apoptosis in several normal tissues where the p53 gene is highly expressed such as lymphoid tissues, hematopoietic organs, intestinal epithelia and testis. Thus, temporary suppression of p53 might be a workable therapeutic strategy to prevent adverse effects of cancer therapy and other drugs (Nayak et al. 2009).

Angiogenesis

Sokotrasterol sulfate (**65**, [Fig. 5B](#)), a trisulfated sterol isolated from the sponge of the family Halichondriidae (Makarieva et al. 1983), was evaluated as an agent that promotes endothelial sprouting. The compound showed a dose-dependent endothelial sprouting activity, which peaked at 5 mg/mL (approximately 0.9 sprouts per bead), and was also able to promote angiogenesis *in vivo* (vascular density of ±5 vessels per mm, at 60 µg/mL). The action of

Figure 5

Antitumor sulfated steroids. (A) Apoptosis. (B). Angiogenesis. (C) Antiproliferative. (D) Tubulin polymerization. (E) PKC ζ inhibition. (F) PTK inhibition. (G) GDPX inhibition. (H) MT1-MMP inhibition. (I) *PXR* agonism. (J) *FXR* antagonism.

65 is dependent on cyclooxygenase-2 (COX-2) activation, vascular endothelial growth factor (VEGF) induction and the $\alpha_v\beta_3$ integrin. It was also found that the sulfate groups played a critical role in angiogenic activity of **65** (Murphy et al. 2006).

Trisulfated lembasterols A (**66**) and B (**67**) (Fig. 5B), isolated from the marine sponge *Petrosia strongylata*, showed inhibitory activity against thymidine phosphorylase (at 41 and 45 μ M, respectively), an enzyme related to angiogenesis that has an important role in growth and metastasis of solid tumors (Aoki et al. 2002). Thymidine phosphorylase is expressed at higher levels in a wide variety of solid tumors than that in the adjacent non-neoplastic tissues. A derivative of **66** without the sulfate group was tested and did not show any inhibition even at 230 μ M, suggesting the importance of the sulfate group for the inhibitory activity against thymidine phosphorylase. Compounds **66** and **67** were the first thymidine phosphorylase inhibitors with a non-nucleoside skeleton (Aoki et al. 2002).

Antiproliferative

Two disulfated sterols **68** and **69** (Fig. 5C), first isolated from the *Ophiopholis aculeata* (Fedorov et al. 1994) and *Ophiura sarsi* (Levina et al. 1990), respectively, were investigated for their cytotoxic activity against Ehrlich carcinoma cells and mouse spleen lymphocytes, inhibitory activity of 3 H-thymidine incorporation into DNA, and the influence of both compounds on transmembrane transport of calcium ions in Ehrlich carcinoma (Aminin et al. 1995). Although they did not exhibit cytotoxic activity against the carcinoma cells, both compounds inhibited the incorporation of 3 H-thymidine around 60 and 50%, at 100 μ g/mL. This inhibition may reflect several mechanisms by which steroids affect macromolecular synthesis (Aminin et al. 1995). Ehrlich carcinoma cells are capable of accumulating radioactive calcium from external medium. Addition of sterols **68** and **69** at 100 μ g/mL to the incubation medium led to an increase of Ca^{2+} uptake in tumor cells of about 260 and 150% above the control, respectively. Compound **69** showed almost half of the activity to stimulate calcium transport compared to **68**, which can be related to the presence of an unsaturated side chain in compound **68** (Aminin et al. 1995).

Tubulin polymerization

Novaeguinosides A (**70**), B (**71**), C (**72**) and D (**73**) (Fig. 5D), isolated from the starfish *Culcita novaeguineae*,

were tested for *in vitro* activity of tubulin polymerization promotion (Tang et al. 2009). Compounds **70** and **72** contain sulfated side chains not previously found in other asterosaponins. On the other hand, the 26-carboxylic acid function in the side chains of **71** and **73** is conjugated with an amide derivative of taurine. Compounds **70**, **72** and **73** exhibited significant activity with P_e (percent end-point promote coefficient) values of 28, 42 and 23% and P_k (percent kinetic promote coefficient) values of 94, 149 and 85% at 12, 9 and 9 μ M, respectively. Compounds **70–73** also displayed *in vitro* cytotoxicity against leukemia K-562 and human hepatoma BEL-7402 cell lines with IC_{50} ranging from 0.7 to 9.5 μ M, **72** being the most potent. The author attributed their cytotoxic activity to the presence of the $\Delta^{9(11)}$ - $3\beta,6\alpha$ -dioxysteroidal aglycone with a sulfate group attached at C-3 and an oligosaccharide moiety at C-6 (Tang et al. 2009).

PKC ζ inhibition

The isoform ζ (zeta) of protein kinase C (PKC) has been implicated as an integral factor in several types of cancer (Whitson et al. 2009). The trisulfated sterols spheciosterol sulfates A-C (**74–76**, Fig. 5E) were isolated from the sponge *Spheciospongia* sp., along with the known sterol trisulfate topsentiasterol sulfate E (**13**). Compounds **74**, **75**, **76** and **13** inhibited not only PKC ζ with IC_{50} values of 1.6, 0.5, 0.1 and 1.2 μ M, respectively, but also NF- κ B activation, in a cell-based assay, with EC_{50} values of 12–64 μ M (Whitson et al. 2008).

Fibrosterol sulfates A (**77**) and B (**78**) (Fig. 5E), two sterol dimmers isolated from the sponge *Lissodendoryx (Acanthodoryx) fibrosa*, were shown to exhibit PKC ζ inhibition with IC_{50} values of 16 and 6 μ M, respectively. Both **77** and **78** composed of two cholestene monomers, with different configuration at C-3 and oxygenation at C-22 in only one monomer. Additionally, the number of sulfate groups in both sterols is also different being four in **77** and five in **78** (Whitson et al. 2009).

PTK inhibition

The oncogenic protein tyrosine kinase (PTK) comprises a family of enzymes that regulate cell growth and intracellular pathways. Halistanol trisulfate (**8**, Fig. 5F) showed activity against PTK pp60^{v-src} with an IC_{50} value of 4 μ M. Acid hydrolysis of **8** to remove the sulfate gave desulfated halistanol, which exhibited no activity. Kinetic studies revealed that **8** is a competitive inhibitor for the peptide-binding site of the enzyme and both competitive and non-competitive for ATP (Slate et al. 1994).

The disulfated sterol **79**, isolated from the green brittle star *Ophiarachna incrassata*, and compounds **7**, **51**, **80** and **81** (Fig. 5F) were found to inhibit the PTK pp60^{v-src} with IC₅₀ of 65, 31, 62, 11 and 12 μM, respectively (Fu et al. 1994). These sterol sulfate PTK inhibitors represented a different chemotype from other known marine natural products PTK inhibitors (Fu et al. 1994).

GDPX inhibition

Ophirapstanol trisulfate (**82**, Fig. 5G), isolated from a deep water marine sponge *Topsentia ophiraphidites*, exhibited significant inhibition in the guanosine diphosphate/G-protein RAS exchange assay (GDPX) (Gunasekera et al. 1994). This assay determines the ability of inhibitors to affect the exchange of guanine nucleotides bound to p21^{RAS} (mutant form of RAS protein). RAS is able to transform cells when it is in the GTP-bound conformation, and mutant forms of RAS lead to unregulated cell proliferation which are found in human cancer (Gunasekera et al. 1994).

MT1-MMP inhibition

Membrane-type matrix metalloproteinases (MT-MMPs) are known as key enzymes in tumor metastasis and inhibitors of MT-MMPs are believed to be potential anticancer drugs (Fujita et al. 2001). Compound **83** (Fig. 5H), a steroid monosulfate with a methylphosphate functionality isolated from the marine sponge *Cribrochalina sp.*, was found as a MT1-MMP inhibitor, showing an IC₅₀ value of 160 mg/mL (Fujita et al. 2001).

PXR agonism

Pregnane X Receptor (PXR) is a member of the nuclear receptors (NRs) superfamily involved in the expression of a wide family of genes (Imperatore et al. 2014). Solomonsterols A (**84**) and B (**85**) (Fig. 5I), two trisulfated sterols isolated from the marine sponge *Theonella swinhoei*, were the first examples of natural marine PXR agonists (Festa et al. 2011). Both sterols effectively increased the expression of two well characterized PXR target genes, CYP3A4 and MDR1, in a human hepatocyte cell line (Festa et al. 2011).

Phallusiasterol A monosulfate (**86**, Fig. 5I), the first sulfated sterol isolated from a tunicate, *Phallusia fumigata*, was shown to induce PXR transactivation in liver carcinoma cells HepG2 cells and stimulated the expression of the PXR target genes CYP3A4 and MDR1 in the same cell line (Imperatore et al. 2014). Recent

studies suggested that the feature and/or the shape of A/B ring junction is critical for the structure-PXR regulating activity relationship in the case of sulfated steroids, since two other compounds, one differing in the configuration at C-6, and another featuring a double bond Δ⁵⁽⁶⁾, lacked PXR modulatory activity (Imperatore et al. 2016).

FXR antagonism

The farnesoid-X-receptor (FXR) is a ligand-regulated transcription factor involved in the regulation of lipid and glucose homeostasis in mammals. A disulfated sterol **87** (Fig. 5J), isolated from the brittle star *Ophiopsis superba* (D'Auria et al. 1989), was tested as an antagonist of FXR and showed potent antagonist activity (80% of FXR transactivation inhibition at 10 μM) in HEpG2 cells transfected with FXR (Sepe et al. 2011).

From the 75 steroids that exhibited antitumor activity (Supplementary data) 41 were glycosylated, from which 28 were asterosaponins (with a sulfate in C-3β, 6α-penta- or hexasaccharide, and a Δ⁹⁽¹¹⁾ unsaturation) and the remaining 13 were mono- (seven), di (four) or triglycosylated (two). In only two cases the sulfate group was in the sugar portion. Concerning the 34 non-glycosylated steroids, four were dimers with more than three sulfate groups, 12 were trisulfated, eight were disulfated and ten were monosulfated (including one that had phosphate a phosphate group simultaneously). From the trisulfated compounds that exhibited antitumor activity, most of them have the three sulfate groups in the ring system, with the substitution pattern 2β-3α-6α-trisulfate.

Cardiovascular activity

Antihypertensive activity

Hymenosulfate (**88**, Fig. 6A), the first sulfated sterol isolated from a marine microalgae (Haptophyte *Hymenomonas sp.*), showed a potent Ca²⁺-releasing activity in sarcoplasmic reticulum, i.e., ten times more potent than caffeine (Kobayashi et al. 1989).

Halistanol disulfate B (**89**, Fig. 6B), isolated from the sponge *Pachastrella sp.*, was able to inhibit the endothelin converting enzyme (ECE) with an IC₅₀ of 2.1 μM, while the non-sulfated counterpart was totally inactive. ECE is associated with the conversion of an inactive intermediate residue into a potent vasoconstrictive peptide endothelin-1 (ET-1) and the overproduction of ET is associated with hypertension and renal failure (Patil et al. 1996).

of saponins, asterosaponins **30**, **34**, **91**, **92**, **93**, and **94** showed activity with hemolytic index values of 22.000, 11.400, 31.400, 21.500, 17.700, and 10.700, respectively, allowing to suspect that a sulfate group at C-3 is important for the hemolytic activity (Fusetani *et al.* 1984). From the starfish *Aphelasterias japonica*, quinovoside aphelasteroside C (**95**), aphelaketotriol (**96**), cheliferoside L1 (**97**), forberoside E3 (**98**), and 3-O-sulfothornasterol A (**99**) were isolated and tested for hemolytic activity against mouse erythrocytes (Ivanchina *et al.* 2000). Steroids **95**, **96**, **97**, **98**, and **99** exhibited ED₅₀ values of 1.9×10^{-4} M, 4.0×10^{-5} M, 1.8×10^{-4} , 3.3×10^{-4} M, and 1.1×10^{-4} M, respectively (Ivanchina *et al.* 2000).

Triseramide (**100**, Fig. 6D), a sulfonated sterol, caused 50% hemolysis of erythrocytes at a minimal concentration of 1.2×10^{-4} M/L (Levina *et al.* 2004). Eight asterosaponins having in common a 3β -SO₃⁻, 6 α -penta- or hexasaccharide, $\Delta^{9(11)}$ unsaturation, a hydroxyl at C-20, and a carbonyl at C-23 of the side chain were also investigated.

Asterosaponins **31** and **32** showed hemolytic activity with ED₅₀ of 16 and 31 μ g/mL, respectively (Tang *et al.* 2005a).

It is possible to observe that the 14 marine sulfated sterols described with hemolytic activity are monosulfated, with only one compound with the sulfate group in a position other than at C-3 (compound **100**), and all present a carbonyl group in the side chain. The majority of the sulfated sterols with hemolytic activity are glycosylated (eight asterosaponins and three monoglycosides).

Sulfated sterols with antihypertensive and antiplatelet activities were isolated from marine sponges (with exception of **88** which was isolated from a microalga), while the sulfated sterols with hemolytic activity were obtained from different species of starfishes.

Anti-inflammatory

The anti-inflammatory activity was reported for six 'classical' asterosaponins, with a sulfate group at C-3, a carbohydrate portion, with five or six sugars, at C-6, a $\Delta^{9(11)}$ unsaturation, and a carbonyl group at C-24 of the side chain (Hwang *et al.* 2011, Thao *et al.* 2013). Compounds **29** and **30** (Fig. 7) inhibited the production of nitric oxide (NO) in Raw 264.7 cells (Hwang *et al.* 2011). When compared to control cells, treatment with **30** in a range of 0.1–4 μ M showed a reduction in NO production up to 19%, while **29**, at 8 μ M, presented an inhibition of 41% (Hwang *et al.* 2011). Four asterosaponins, astrosterosides A (**101**), C (**102**) and D (**61**), which were isolated for the first time from the edible Vietnamese starfish *Astropecten monacanthus* (Thao *et al.* 2013), and marthasteroside B (**103**) (Fig. 7), were reported for their effects on the inflammatory response of bone marrow-derived dendritic cells, by measuring the production of three pro-inflammatory cytokines, tumor necrosis factor (TNF- α), interleukin 6 (IL-6) and interleukin 6 subunit p40 (IL-12 p40) (Thao *et al.* 2013). Compound **61** exhibited potent inhibitory effects on the production of TNF- α , IL-6 and IL-12 p40 with IC₅₀ values of 1.2, 3.5 and 0.60 μ M,

Figure 7
Anti-inflammatory sulfated steroids.

respectively. Compounds **101** and **103** presented significant inhibitory effects on IL-6 production with IC_{50} values of 3.17 and 4.37 μ M, respectively, while **102** showed moderate inhibitory effects with IC_{50} of 30.5 μ M. The author concluded that the presence of an α,β -unsaturated carbonyl in the side chains of **101** and **103** and the two ketone groups in **61** were important in the inhibitory effects of these compounds (Thao et al. 2013).

All sulfated steroids described with anti-inflammatory activity (Fig. 7) were asterosaponins that were isolated from starfishes. With exception of **103**, all anti-inflammatory sulfated steroids also displayed antitumor activity, which could increase the potential of these compounds as models for new antitumor agents (Rayburn et al. 2009).

Others

Inhibition of BACE1

From an undescribed species of *Topsentia*, topsentinol K trisulfated (**104**, Fig. 8A), a rare 24-isopropyl sulfated sterol was isolated. Compound **104** inhibited the aspartic protease BACE1-mediated cleavage of amyloid precursor protein in a dose-dependent manner (IC_{50} value of 1.2 μ M), although in a detergent-dependent manner suggestive of non-specific aggregation (Dai et al. 2010). The activity of **104** was attributed to the presence of the sulfate esters, since the desulfated compounds did not show activity (Dai et al. 2010).

Anti-osteoblast proliferation

The high prevalence of osteoporosis has been regarded as a major public health problem all over the world. The effects on osteoblastic proliferation of some sulfated sterols isolated from the starfish *Asterias amurensis* were evaluated in the *in vitro* model UMR-106 cell lines. Compound **105** (Fig. 8B) significantly promoted the osteoblastic proliferation, in a range of concentrations of 0.01–100 μ M, while **106** (Fig. 8B) was found to promote osteoblastic proliferation at higher concentrations. In contrast, a non-glycosylated non-sulfated derivative of **105** and a sulfated and non-glycosylated compound, also isolated from the starfish, exhibited no activity, showing that the β -D-xylose residue of saponins is necessary for

activity (Liu et al. 2008). Both sulfated saponins have in common the presence of the sulfate group at C-15 α , which is rare among the sulfated sterols reported, two hydroxyl groups at C-3 β and C-6 α , and a β -D-xylose in the side chain.

Environmental applications

Antifouling activity

Fouling on ships represents serious economic and environmental problems. These problems are associated with frictional resistance resulting in generated roughness which leads to speed loss and accelerated fuel consumption, in addition to increasing costs and emission of harmful gases as well as money and time for cleaning and maintenance (Schultz 2007). Biofouling in ships may also lead to transport of non-native species, causing a major human health and environmental impact (Abbott et al. 2000). Thus, the search for antifouling products becomes a fast growing research field and some sulfated compounds have been reported as potential nontoxic antifouling agents (Almeida et al. 2017, Vilas-Boas et al. 2017).

Carboxylated sterol sulfates (**1–3**, Fig. 9), showed antifouling activity against *Salmacinia tribanchiata* at 10 μ g/mL (Nakatsu et al. 1983). Compounds **4**, **107**, and **108** (Fig. 9), isolated from the starfish *Luidia clathrata*, were able to inhibit the attachment of barnacle larvae at 0.2 mg/mL, while asterosaponins **30**, **103**, ophydianoside F (**109**), and regularoside B (**110**) (Fig. 9) showed complete inhibition of larvae attachment at the same concentration (Iorizzi et al. 1995). This study was the first demonstration that bioactive compounds from echinoderms may play a role in the prevention of fouling. It is believed that echinoderms have evolved an effective mechanism of protection from fouling (Iorizzi et al. 1995). Compound **8** showed antifouling activity, and no toxicity, against cypris larvae of the barnacle *Balanus amphitrite*, with an IC_{50} of 5.0 μ g/mL (Tsukamoto et al. 1997, Fusetani 2004).

Goniopectenosides A–C (**111–113**, Fig. 9), three asterosaponins that possess the same pentasaccharide moiety linked to C-6 of the sterol nucleus and with a 22-oxo functionality in the side chain, were isolated from the starfish *Goniopecten demonstrans*, and were found to

Figure 8
(A) Sulfated steroid with BACE1 inhibitory activity. (B) Sulfated steroids with anti-osteoblast activity.

group at C-3, to polyhydroxysteroids monosulfates, with both sulfate and hydroxyl groups. Monosulfates of glycosylated steroids were also found with the carbohydrate moieties in the ring system or in the side chain and few with the carbohydrates in both. The most common of these steroidal glycosides sulfates were the classical asterosaponins, with the sulfate group at C-3 and the carbohydrate chain of five or six sugars attached at C-6. An unusual compound, which is an inhibitor of MT1-MMP, having both sulfate and phosphate groups in the ring system, was also isolated (Fujita *et al.* 2001).

Disulfated steroids frequently have one sulfate group in the ring system (usually at C-3) and other in the side chain (usually at C-21). However, some sulfated steroids presented both sulfate groups in the ring system, with a large structural diversity, e.g. both sulfates in ring A, one sulfate in ring A and another in ring B, and one sulfate in ring A and a sulfonamide in ring D. Disulfated steroids were mainly simultaneously hydroxylated and sulfated and few were glycosylated.

In the case of trisulfated steroids, the three sulfate groups appear more frequently with the common 2 β , 3 α , 6 α -trisulfate substitution pattern and only few presented sulfation at C-2, C-3 and C-21.

These secondary metabolites showed a broad spectrum of complementary biological activities, namely antimicrobial and/or antifouling activities, and antitumor, cardiovascular and/or anti-inflammatory activities. Although more extensive assays were precluded by the limited availability of pure compounds, halistanol trisulfate (**8**) was the most well-studied sulfated steroid, followed by thornasteroside A (**30**). Both compounds are remarkable for their wide variety of bioactivities. New enzyme inhibitory activities are still being reported, for example, inhibitory activity against NAD⁺-dependent lysine deacetylase (KDAC) – sirtuins (SIRT) – of three halistanol sulfates was reported during the review process of this work (Nakamura *et al.* 2018).

It is also worth mentioning that due to their hydrophilic scaffold, sulfated steroids can reach intracellular targets and improve bioavailability, which represents a tremendous advantage in therapeutics. On the other hand, the nontoxicity associated to the sulfated molecules raises the probability of discovering novel and safer therapeutic agents (Correia-da-Silva *et al.* 2014). From the marine compounds currently in the market, only Prialt and Yondelis, became drugs without any modification of the original natural molecule, while all the others suffered lead optimization, in different steps of their development. Therefore, optimization of these

marine sulfated steroids by structural modifications or synthetic supply of unmodified natural molecule could be worth to explore to develop new solutions for some diseases as also for environmental problems.

Supplementary data

This is linked to the online version of the paper at <https://doi.org/10.1530/JME-17-0252>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was supported through national funds provided by FCT/MCTES – Foundation for Science and Technology from the Minister of Science, Technology and Higher Education (PIDDAC) and European Regional Development Fund (ERDF) through the COMPETE – Programa Operacional Factores de Competitividade (POFC) programme, under the projects PTDC/MAR-BIO/4694/2014 (reference POCI-01-0145-FEDER-016790; Project 3599 PPCDT – Promover a Produção Científica e Desenvolvimento Tecnológico e a Constituição de Redes Temáticas) and PTDC/AAG-TEC/0739/2014 (reference POCI-01-0145-FEDER-016793; Project 9471 RIDTI – Reforçar a Investigação, o Desenvolvimento Tecnológico e a Inovação) in the framework of the programme PT2020.

Acknowledgments

The author acknowledges the Strategic Funding UID/Multi/04423/2013 provided by FCT – Foundation for Science and Technology and European Regional Development Fund (ERDF), in the framework of the programme PT2020.

References

- Abbott A, Abel PD, Arnold DW & Milne A 2000 Cost-benefit analysis of the use of TBT: the case for a treatment approach. *Science of the Total Environment* **258** 5–19. ([https://doi.org/10.1016/S0048-9697\(00\)00505-2](https://doi.org/10.1016/S0048-9697(00)00505-2))
- Abd AJ, Kanwar RK & Kanwar JR 2017 Aged macular degeneration: current therapeutics for management and promising new drug candidates. *Drug Discovery Today* **22** 1671–1679. (<https://doi.org/10.1016/j.drudis.2017.1007.1010>)
- Almeida JR, Correia-da-Silva M, Sousa E, Antunes J, Pinto M, Vasconcelos V & Cunha I 2017 Antifouling potential of nature-inspired sulfated compounds. *Scientific Reports* **7** 42424. (<https://doi.org/10.1038/srep42424>)
- Aminin DL, Agafonova IG & Fedorov SN 1995 Biological activity of disulfated polyhydroxysteroids from the Pacific brittle star *Ophiopholis aculeata*. *Comparative Biochemistry and Physiology: Part C: Pharmacology, Toxicology and Endocrinology* **112** 201–204. ([https://doi.org/10.1016/0742-8413\(95\)00212-8](https://doi.org/10.1016/0742-8413(95)00212-8))
- Andersson L, Bohlin L, Iorizzi M, Riccio R, Minale L & Moreno-Lopez W 1989 Biological activity of saponins and saponin-like compounds from starfish and brittle-stars. *Toxicon* **27** 179–188. ([https://doi.org/10.1016/0041-0101\(89\)90131-1](https://doi.org/10.1016/0041-0101(89)90131-1))

- Aoki S, Naka Y, Itoh T, Furukawa T, Rachmat R, Akiyama S & Kobayashi M 2002 Lembehsterols A and B, novel sulfated sterols inhibiting thymidine phosphorylase, from the marine sponge *Petrosia strongylata*. *Chemical and Pharmaceutical Bulletin* **50** 827–830. (<https://doi.org/10.1248/cpb.50.827>)
- Appeltans W, Ah Yong Shane T, Anderson G, Angel Martin V, Artois T, Bailly N, Bamber R, Barber A, Bartsch I, Berta A, et al. 2012 The magnitude of global marine species diversity. *Current Biology* **22** 2189–2202. (<https://doi.org/10.1016/j.cub.2012.09.036>)
- Bifulco G, Bruno I, Minale L & Riccio R 1994 Novel HIV-inhibitory halistanol sulfates F-H from a marine sponge, *Pseudoaxinissa digitata*. *Journal of Natural Products* **57** 164–167. (<https://doi.org/10.1021/np50103a026>)
- Boonlarpapradab C & Faulkner DJ 2007 Eurysterols A and B, cytotoxic and antifungal steroidal sulfates from a marine sponge of the genus *Euryspongia*. *Journal of Natural Products* **70** 846–848. (<https://doi.org/10.1021/np060472c>)
- Butler MS, Blaskovich MAT & Cooper MA 2017 Antibiotics in the clinical pipeline at the end of 2015. *Journal of Antibiotics* **70** 3–24. (<https://doi.org/10.1038/ja.2016.72>)
- Cheng G, Zhang X, Tang HF, Zhang Y, Zhang XH, Cao WD, Gao DK, Wang XL & Jin BQ 2006 Asterosaponin 1, a cytostatic compound from the starfish *Culcita novaeguineae*, functions by inducing apoptosis in human glioblastoma U87MG cells. *Journal of Neuro-Oncology* **79** 235–241. (<https://doi.org/10.1007/s11060-006-9136-y>)
- Chludil HD & Maier MS 2005 Minutosides A and B, antifungal sulfated steroid xylosides from the patagonian starfish *Anasterias minuta*. *Journal of Natural Products* **68** 1279–1283. (<https://doi.org/10.1021/np050086f>)
- Chludil HD, Seldes AM & Maier MS 2002 Antifungal steroidal glycosides from the patagonian starfish *Anasterias minuta*: structure-activity correlations. *Journal of Natural Products* **65** 153–157. (<https://doi.org/10.1021/np010332x>)
- Comin MJ, Maier MS, Roccatagliata AJ, Pujol CA & Damonte EB 1999 Evaluation of the antiviral activity of natural sulfated polyhydroxysteroids and their synthetic derivatives and analogs. *Steroids* **64** 335–340. ([https://doi.org/10.1016/S0039-128X\(99\)00016-1](https://doi.org/10.1016/S0039-128X(99)00016-1))
- Connolly B, Desai A, Garcia CA, Thomas E & Gast MJ 2006 Squalamine lactate for exudative age-related macular degeneration. *Ophthalmology Clinics* **19** 381–391. (<https://doi.org/10.1016/j.ophc.2006.05.003>)
- Correia-da-Silva M, Sousa E & Pinto MM 2014 Emerging sulfated flavonoids and other polyphenols as drugs: nature as an inspiration. *Medicinal Research Reviews* **34** 223–279. (<https://doi.org/10.1002/med.21282>)
- D'Auria MV, Riccio R, Minale L, La Barre S & Pusset J 1987 Novel marine steroid sulfates from Pacific ophiuroids. *Journal of Organic Chemistry* **52** 3947–3952. (<https://doi.org/10.1021/jo00227a001>)
- D'Auria MV, Riccio R, Uriarte E, Minale L, Tanaka J & Higa T 1989 Isolation and structure elucidation of seven new polyhydroxylated sulfated sterols from the ophiuroid *Ophiolepis superba*. *Journal of Organic Chemistry* **54** 234–239. (<https://doi.org/10.1021/jo00262a049>)
- Dai J, Sorribas A, Yoshida WY, Kelly M & Williams PG 2010 Topsentinols, 24-Isopropyl steroids from the marine sponge *Topsentia* sp. *Journal of Natural Products* **73** 1597–1600. (<https://doi.org/10.1021/np100374b>)
- DiGirolamo JA, Li X-C, Jacob MR, Clark AM & Ferreira D 2009 Reversal of fluconazole resistance by sulfated sterols from the marine sponge *Topsentia* sp. *Journal of Natural Products* **72** 1524–1528. (<https://doi.org/10.1021/np900177m>)
- Falkenstein E, Tillmann HC, Christ M, Feuring M & Wehling M 2000 Multiple actions of steroid hormones – a focus on rapid, nongenomic effects. *Pharmacological Reviews* **52** 513–556. ([https://doi.org/0031-6997/00/5204-0513\\$03.00/0](https://doi.org/0031-6997/00/5204-0513$03.00/0))
- Fedorov SN, Levina EV, Kalinovsky AI, Dmitrenok PS & Stonik VA 1994 Sulfated steroids from Pacific brittle stars *Ophiopholis aculeata*, *Ophiura sarsi*, and *Stegophiura brachiactis*. *Journal of Natural Products* **57** 1631–1637. (<https://doi.org/10.1021/np50114a004>)
- Festa C, De Marino S, D'Auria MV, Bifulco G, Renga B, Fiorucci S, Petek S & Zampella A 2011 Solomonsterols A and B from *Theonella swinhoei*. The first example of C-24 and C-23 sulfated sterols from a marine source endowed with a PXR agonistic activity. *Journal of Medicinal Chemistry* **54** 401–405. (<https://doi.org/10.1021/jm100968b>)
- Fine JM & Sorensen PW 2005 Biologically relevant concentrations of petromyzonol sulfate, a component of the sea lamprey migratory pheromone, measured in stream water. *Journal of Chemical Ecology* **31** 2205–2210. (<https://doi.org/10.1007/s10886-005-6745-4>)
- Fiorucci S, Distrutti E, Bifulco G, D'Auria MV & Zampella A 2012 Marine sponge steroids as nuclear receptor ligands. *Trends in Pharmacological Sciences* **33** 591–601. (<https://doi.org/10.1016/j.tips.2012.08.004>)
- Fu X, Schmitz FJ, Lee RH, Papkoff JS & Slate DL 1994 Inhibitors of protein tyrosine kinase pp60v-src: sterol sulfates from the brittle star *Ophiarachna incrassata*. *Journal of Natural Products* **57** 1591–1594. (<https://doi.org/10.1021/np50113a023>)
- Fujita M, Nakao Y, Matsunaga S, Seiki M, Itoh Y, van Soest RWM, Heubes M, Faulkner DJ & Fusetani N 2001 Isolation and structure elucidation of two phosphorylated sterol sulfates, MT1-MMP inhibitors from a marine sponge *Cribrachalina* sp.: revision of the structures of haplosamates A and B. *Tetrahedron* **57** 3885–3890. ([https://doi.org/10.1016/S0040-4020\(01\)00259-9](https://doi.org/10.1016/S0040-4020(01)00259-9))
- Fusetani N 2004 Biofouling and antifouling. *Natural Products Reports* **21** 94–104. (<https://doi.org/10.1039/b302231p>)
- Fusetani N, Matsunaga S & Konosu S 1981 Bioactive marine metabolites II. Halistanol sulfate, an antimicrobial novel steroid sulfate from the marine sponge *Halichondria* cf. *moorei* bergquist. *Tetrahedron Letters* **22** 1985–1988. ([https://doi.org/10.1016/S0040-4039\(01\)92885-0](https://doi.org/10.1016/S0040-4039(01)92885-0))
- Fusetani N, Kato Y, Hashimoto K, Komori T, Itakura Y & Kawasaki T 1984 Biological activities of asterosaponins with special reference to structure-activity relationships. *Journal of Natural Products* **47** 997–1002. (<https://doi.org/10.1021/np50036a015>)
- Fusetani N, Takahashi M & Matsunaga S 1994 Topsentiasterol sulfates, antimicrobial sterol sulfates possessing novel side chains, from a marine sponge, *Topsentia* sp. *Tetrahedron* **50** 7765–7770. ([https://doi.org/10.1016/S0040-4020\(01\)85260-1](https://doi.org/10.1016/S0040-4020(01)85260-1))
- Giddings LA & Newman DJ 2015 Bioactive compounds from marine extremophiles. In *Bioactive Compounds from Marine Extremophiles*, pp 1–124. Eds SM Tiquia-Arashiro & M Mormile. Cham: Springer International Publishing.
- Glaser KB & Mayer AM 2009 A renaissance in marine pharmacology: from preclinical curiosity to clinical reality. *Biochemical Pharmacology* **78** 440–448. (<https://doi.org/10.1016/j.bcp.2009.04.015>)
- Guimaraes T, Quiroz CG, Borges CR, Oliveira SQ, Almeida MT, Bianco EM, Moritz MI, Carraro JL, Palermo JA, Cabrera G, et al. 2013 Anti HSV-1 activity of halistanol sulfate and halistanol sulfate C isolated from Brazilian marine sponge *Petromica citrina* (Demospongiae). *Marine Drugs* **11** 4176–4192. (<https://doi.org/10.3390/md11114176>)
- Gunasekera SP, Sennett SH, Kelly-Borges M & Bryant RW 1994 Ophirapstanol trisulfate, a new biologically active steroid sulfate from the deep water marine sponge *Topsentia* ophiraphidites. *Journal of Natural Products* **57** 1751–1754. (<https://doi.org/10.1021/np50114a024>)
- Guzii AG, Makarieva TN, Denisenko VA, Dmitrenok PS, Burtseva YV, Krasokhin VB & Stonik VA 2008 Topsentiasterol sulfates with novel iodinated and chlorinated side chains from the marine sponge *Topsentia* sp. *Tetrahedron Letters* **49** 7191–7193. (<https://doi.org/10.1016/j.tetlet.2008.10.007>)
- Hussain RM & Ciulla TA 2017 Emerging vascular endothelial growth factor antagonists to treat neovascular age-related macular

- degeneration. *Expert Opinion on Emerging Drugs* **22** 235–246. (<https://doi.org/10.1080/14728214.2017.1362390>)
- Hwang IH, Kim DW, Kim SJ, Min BS, Lee SH, Son JK, Kim CH, Chang HW & Na M 2011 Asterosaponins isolated from the starfish *Asterias amurensis*. *Chemical and Pharmaceutical Bulletin* **59** 78–83. (<https://doi.org/10.1248/cpb.59.78>)
- Imperatore C, D'Aniello F, Aiello A, Fiorucci S, D'Amore C, Sepe V & Menna M 2014 Phallusiasterols A and B: two new sulfated sterols from the Mediterranean tunicate *Phallusia fumigata* and their effects as modulators of the PXR receptor. *Marine Drugs* **12** 2066–2078. (<https://doi.org/10.3390/md12042066>)
- Imperatore C, Senese M, Aiello A, Luciano P, Fiorucci S, D'Amore C, Carino A & Menna M 2016 Phallusiasterol C, a new disulfated steroid from the Mediterranean tunicate *Phallusia fumigata*. *Marine Drugs* **14** 117. (<https://doi.org/10.3390/md14060117>)
- Iorizzi M, Bryan P, McClintock J, Minale L, Palagiano E, Maurelli S, Riccio R & Zollo F 1995 Chemical and biological investigation of the polar constituents of the starfish *Luidia clathrata*, collected in the Gulf of Mexico. *Journal of Natural Products* **58** 653–671. (<https://doi.org/10.1021/np50119a003>)
- Ivanchina NV, Kicha AA, Kalinovsky AI, Dmitrenok PS, Stonik VA, Riguera R & Jimenez C 2000 Hemolytic polar steroidal constituents of the starfish *Aphelasterias japonica*. *Journal of Natural Products* **63** 1178–1181. (<https://doi.org/10.1021/np000030f>)
- Jaspars M, De Pascale D, Andersen JH, Reyes F, Crawford AD & Ianora A 2016 The marine biodiversity pipeline and ocean medicines of tomorrow. *Journal of the Marine Biological Association of the United Kingdom* **96** 151–158. (<https://doi.org/10.1017/S0025315415002106>)
- Kicha AA, Ivanchina NV, Kalinovsky AI, Dmitrenok PS & Stonik VA 2000 Asterosaponin P2 from the Far-Eastern starfish *patiria* (asterina) pectinifera. *Russian Chemical Bulletin* **49** 1794–1795. (<https://doi.org/10.1007/BF02496356>)
- Kicha AA, Ivanchina NV, Huong TTT, Kalinovsky AI, Dmitrenok PS, Fedorov SN, Dyshlovoy SA, Long PQ & Stonik VA 2010 Two new asterosaponins, archasterosides A and B, from the Vietnamese starfish *Archaster typicus* and their anticancer properties. *Bioorganic and Medicinal Chemistry Letters* **20** 3826–3830. (<https://doi.org/10.1016/j.bmcl.2010.04.005>)
- Kobayashi Ji, Ishibashi M, Nakamura H, Ohizumi Y & Hirata Y 1989 Hymenosulphate, a novel sterol sulphate with Ca-releasing activity from the cultured marine haptophyte *Hymenomonas* sp. *Journal of the Chemical Society, Perkin Transactions* **10** 101–103. (<https://doi.org/10.1039/p19890000101>)
- Kong F & Andersen RJ 1993 Polymastiamide A, a novel steroid/amino acid conjugate isolated from the Norwegian marine sponge *Polymastia boletiformis* (Lamarck, 1815). *Journal of Organic Chemistry* **58** 6924–6927. (<https://doi.org/10.1021/jo00076a073>)
- Kornprobst JM, Sallenave C & Barnathan G 1998 Sulfated compounds from marine organisms. *Comparative Biochemistry and Physiology: Part B: Biochemistry and Molecular Biology* **119** 1–51. ([https://doi.org/10.1016/S0305-0491\(97\)00168-5](https://doi.org/10.1016/S0305-0491(97)00168-5))
- Kossuga MH, Lira SPd, Nascimento AM, Gambardella MTP, Berlinck RGS, Torres YR, Nascimento GGF, Pimenta EF, Silva M, Thiemann OH, et al. 2007 Isolamento e atividades biológicas de produtos naturais das esponjas *Monanchora arbuscula*, *Aplysina* sp. *Petromica ciocalyptoides* e *Topsentia ophiraphidites*, da ascídia *Didemnum ligulum* e do octocoral *Carijoa riisei*. *Química Nova* **30** 1194–1202. (<https://doi.org/10.1590/S0100-40422007000500027>)
- Lerch ML & Faulkner DJ 2001 Unusual polyoxygenated sterols from a Philippines sponge *Xestospongia* sp. *Tetrahedron* **57** 4091–4094. ([https://doi.org/10.1016/S0040-4020\(01\)00296-4](https://doi.org/10.1016/S0040-4020(01)00296-4))
- Levina EV, Fedorov SN, Stonik VA, Andriyashchenko PV, Kalinovskii AI & Isakov VV 1990 Steroid compounds from ophiuroids II. Sulfated steroids from *Ophiura sarsi* and *Ophiura leptocentia*. *Chemistry of Natural Compounds* **26** 408–411. (<https://doi.org/10.1007/BF00598993>)
- Levina EV, Kalinovskii AI, Dmitrenok PS, Prokof'eva NG, Andriyashchenko PV & Stonik VA 2004 Triseramide is a new steroid conjugate from the starfish *Astropecten triseriatus*. *Doklady Biochemistry and Biophysics* **396** 171–173. (<https://doi.org/10.1023/B:DOBI.0000033521.54237.83>)
- Levina EV, Aminin DL, Koval'chuk SN, Kozhemiako VB, Dyshlovoi SA, Kalinovskii AI & Dmitrenok PS 2010 Polar steroids from *Solaster endeca* starfish and the physiological activity of polar steroids from three starfish species. *Bioorganicheskaya Khimiya* **36** 251–258. (<https://doi.org/10.1134/S1068162010020135>)
- Li X-C, Jacob MR, Ding Y, Agarwal AK, Smillie TJ, Khan SI, Nagle DG, Ferreira D & Clark AM 2006 Capisterones A and B, which enhance fluconazole activity in *Saccharomyces cerevisiae*, from the marine green alga *Penicillus capitatus*. *Journal of Natural Products* **69** 542–546. (<https://doi.org/10.1021/np050396y>)
- Lima BdA, Lira SPd, Kossuga MH, Gonçalves RB, Berlinck RGS & Kamiya RU 2014 Halistanol sulfate A and rodriguesins A and B are antimicrobial and antibiofilm agents against the cariogenic bacterium *Streptococcus mutans*. *Revista Brasileira de Farmacognosia* **24** 651–659. (<https://doi.org/10.1016/j.bjp.2014.11.002>)
- Liu HW, Li JK, Zhang DW, Zhang JC, Wang NL, Cai GP & Yao XS 2008 Two new steroidal compounds from starfish *Asterias amurensis* Lutken. *Journal of Asian Natural Products Research* **10** 521–529. (<https://doi.org/10.1080/10286020801966674>)
- Lockhart SR & Warnock DW 2015 Antifungal agents and susceptibility test methods. In *Manual of Clinical Microbiology*, edn 17. Eds James H Jorgensen, Michael A Pfaller, Karen C Carroll, Guido Funke, Marie Louise Landry, Sandra S Richter and David W Warnock. American Society of Microbiology.
- Makarieva TN, Shubina LK, Kalinovsky AI, Stonik VA & Elyakov GB 1983 Steroids in Porifera. II. Steroid derivatives from two sponges of the family Halichondriidae. Sokotrasterol sulfate, a marine steroid with a new pattern of side chain alkylation. *Steroids* **42** 267–281. ([https://doi.org/10.1016/0039-128X\(83\)90039-9](https://doi.org/10.1016/0039-128X(83)90039-9))
- Makarieva TN, Stonik VA, D'Yachuk OG & Dmitrenok AS 1995 Annasterol sulfate, a novel marine sulfated steroid, inhibitor of glucanase activity from the deep water sponge *Poecillastra laminaris*. *Tetrahedron Letters* **36** 129–132. ([https://doi.org/10.1016/0040-4039\(94\)02192-E](https://doi.org/10.1016/0040-4039(94)02192-E))
- Malyarenko TV, Malyarenko Vishchuk OS, Ivanchina NV, Kalinovsky AI, Popov RS & Kicha AA 2015 Four new sulfated polar steroids from the Far Eastern starfish *Leptasterias ochotensis*: structures and activities. *Marine Drugs* **13** 4418–4435. (<https://doi.org/10.3390/md13074418>)
- Marinho PR, Simas NK, Kuster RM, Duarte RS, Fracalanza SE, Ferreira DE, Romanos MT, Muricy G, Giambiagi-Demarval M & Laport MS 2012 Antibacterial activity and cytotoxicity analysis of halistanol trisulphate from marine sponge *Petromica citrina*. *Journal of Antimicrobial Chemotherapy* **67** 2396–2400. (<https://doi.org/10.1093/jac/dks229>)
- Marino Simona D, Iorizzi M, Zollo F, Amsler Charles D, Greer Stephen P & McClintock James B 2000 Three new asterosaponins from the starfish *Gonioplecten demonstrans*. *European Journal of Organic Chemistry* **2000** 4093–4098. ([https://doi.org/10.1002/1099-0690\(200012\)2000:24<4093::AID-EJOC4093>3.0.CO;2-M](https://doi.org/10.1002/1099-0690(200012)2000:24<4093::AID-EJOC4093>3.0.CO;2-M))
- Martins A, Vieira H, Gaspar H & Santos S 2014 Marketed marine natural products in the pharmaceutical and cosmeceutical industries: tips for success. *Marine Drugs* **12** 1066–1101. (<https://doi.org/10.3390/md12021066>)
- Mayer AM, Glaser KB, Cuevas C, Jacobs RS, Kem W, Little RD, McIntosh JM, Newman DJ, Potts BC & Shuster DE 2010 The odyssey of marine pharmaceuticals: a current pipeline perspective. *Trends in Pharmacological Sciences* **31** 255–265. (<https://doi.org/10.1016/j.tips.2010.02.005>)
- Mayer AM, Nguyen M, Kalwajtys P, Kerns H, Newman DJ & Glaser KB 2017 The marine pharmacology and pharmaceuticals pipeline in

2016. *FASEB Journal* **31** 818.811. (<https://doi.org/10.1096/fj.1530-6860>)
- McKee TC, Cardellina JH, Tischler M, Snader KM & Boyd MR 1993 Ibisterol sulfate, a novel HIV-inhibitory sulfated sterol from the deep water sponge *Topsentia* sp. *Tetrahedron Letters* **34** 389–392. ([https://doi.org/10.1016/0040-4039\(93\)85083-9](https://doi.org/10.1016/0040-4039(93)85083-9))
- McKee TC, Cardellina JH, Riccio R, D'Auria MV, Iorizzi M, Minale L, Moran RA, Gulakowski RJ & McMahon JB 1994 HIV-inhibitory natural products. 11. Comparative studies of sulfated sterols from marine invertebrates. *Journal of Medicinal Chemistry* **37** 793–797. (<https://doi.org/10.1021/jm00032a012>)
- Moore KS, Wehrli S, Roder H, Rogers M, Forrest JN, McCrimmon D & Zasloff M 1993 Squalamine: an aminosterol antibiotic from the shark. *PNAS* **90** 1354–1358. (<https://doi.org/10.1073/pnas.90.4.1354>)
- Morinaka BI, Pawlik JR & Molinski TF 2009 Amaroxocanes A and B: sulfated dimeric sterols defend the Caribbean coral reef sponge *Phorbas amaranthus* from fish predators. *Journal of Natural Products* **72** 259–264. (<https://doi.org/10.1021/np800652v>)
- Moss GP 1989 The nomenclature of steroids. *European Journal of Biochemistry* **186** 429–458. (<https://doi.org/10.1111/j.1432-1033.1989.tb15228.x>)
- Mueller JW, Gilligan LC, Idkowiak J, Arlt W & Foster PA 2015 The regulation of steroid action by sulfation and desulfation. *Endocrine Reviews* **36** 526–563. (<https://doi.org/10.1210/er.2015-1036>)
- Murphy S, Larrivee B, Pollet I, Craig KS, Williams DE, Huang XH, Abbott M, Wong F, Curtis C, Conrads TP, et al. 2006 Identification of sokotrasterol sulfate as a novel proangiogenic steroid. *Circulation Research* **99** 257–265. (<https://doi.org/10.1161/01.RES.0000233316.17882.33>)
- Nakamura F, Kudo N, Tomachi Y, Nakata A, Takemoto M, Ito A, Tabei H, Arai D, de Voogd N, Yoshida M, et al. 2018 Halistanol sulfates I and J, new SIRT1–3 inhibitory steroid sulfates from a marine sponge of the genus *Halichondria*. *Journal of Antibiotics* **71** 273–278. (<https://doi.org/10.1038/ja.2017.145>)
- Nakatsu T, Walker RP, Thompson JE & Faulkner DJ 1983 Biologically-active sterol sulfates from the marine sponge *Toxadocia zumi*. *Experientia* **39** 759–761. (<https://doi.org/10.1007/BF01990312>)
- Nayak SK, Panesar PS & Kumar H 2009 p53-Induced apoptosis and inhibitors of p53. *Current Medicinal Chemistry* **16** 2627–2640. (<https://doi.org/10.2174/092986709788681976>)
- Patil AD, Freyer AJ, Breen A, Carte B & Johnson RK 1996 Halistanol disulfate B, a novel sulfated sterol from the sponge *Pachastrella* sp.: inhibitor of endothelin converting enzyme. *Journal of Natural Products* **59** 606–608. (<https://doi.org/10.1021/np9601770>)
- Peng Y, Zheng J, Huang R, Wang Y, Xu T, Zhou X, Liu Q, Zeng F, Ju H, Yang X, et al. 2010 Polyhydroxy steroids and saponins from China sea starfish *Asterina pectinifera* and their biological activities. *Chemical and Pharmaceutical Bulletin* **58** 856–858. (<https://doi.org/10.1248/cpb.58.856>)
- Perfect JR 2016 Is there an emerging need for new antifungals? *Expert Opinion on Emerging Drugs* **21** 129–131. (<https://doi.org/10.1517/14728214.2016.1155554>)
- Pimm SL 2012 Biodiversity: not just lots of fish in the sea. *Current Biology* **22** R996–R997. (<https://doi.org/10.1016/j.cub.2012.09.028>)
- Pinet PR 2011 The properties of seawater. In *Invitation to Oceanography*, edn 5, Eds: Molly Steinbach, Dean W DeChambeau, and Megan R Turner pp 134–186. Canada: Jones & Bartlett Learning.
- Puglisi MP, Tan LT, Jensen PR & Fenical W 2004 Capisterones A and B from the tropical green alga *Penicillus capitatus*: unexpected antifungal defenses targeting the marine pathogen *Lindra thalassiae*. *Tetrahedron* **60** 7035–7039. (<https://doi.org/10.1016/j.tet.2003.10.131>)
- Qureshi A & Faulkner DJ 1999 Haplosamates A and B: new steroidal sulfamate esters from two Haplosclerid sponges. *Tetrahedron* **55** 8323–8330. ([https://doi.org/10.1016/S0040-4020\(99\)00465-2](https://doi.org/10.1016/S0040-4020(99)00465-2))
- Rayburn ER, Ezell SJ & Zhang R 2009 Anti-inflammatory agents for cancer therapy. *Molecular and Cellular Pharmacology* **1** 29–43. (<https://doi.org/10.4255/mcpharmacol.09.05>)
- Riccio R, D'Auria MV & Minale L 1985 Unusual sulfated marine steroids from the ophiuroid *Ophioderma longicaudum*. *Tetrahedron* **41** 6041–6046. ([https://doi.org/10.1016/S0040-4020\(01\)91445-0](https://doi.org/10.1016/S0040-4020(01)91445-0))
- Riccio R, Iorizzi M & Minale L 1986 Starfish saponins XXX. Isolation of sixteen steroidal glycosides and three polyhydroxysteroids from the Mediterranean starfish *Coscinasterias tenuispina*. *Bulletin des Sociétés Chimiques Belges* **95** 869–893. (<https://doi.org/10.1002/bcsb.19860950912>)
- Roccatagliata AJ, Maier MS, Seldes AM, Pujol CA & Damonte EB 1996 Antiviral sulfated steroids from the ophiuroid *Ophioplocus januarii*. *Journal of Natural Products* **59** 887–889. (<https://doi.org/10.1021/np960171a>)
- Roccatagliata AJ, Maier MS & Seldes AM 1998 New sulfated polyhydroxysteroids from the Antarctic ophiuroid *Astrothoma agassizii*. *Journal of Natural Products* **61** 370–374. (<https://doi.org/10.1021/np970429c>)
- Rudi A, Yosief T, Loya S, Hizi A, Schleyer M & Kashman Y 2001 Clathsterol, a novel anti-HIV-1 RT sulfated sterol from the sponge *Clathria* species. *Journal of Natural Products* **64** 1451–1453. (<https://doi.org/10.1021/np010121s>)
- Schultz MP 2007 Effects of coating roughness and biofouling on ship resistance and powering. *Biofouling* **23** 331–341. (<https://doi.org/10.1080/08927010701461974>)
- Sepe V, Bifulco G, Renga B, D'Amore C, Fiorucci S & Zampella A 2011 Discovery of sulfated sterols from marine invertebrates as a new class of marine natural antagonists of farnesoid-X-receptor. *Journal of Medicinal Chemistry* **54** 1314–1320. (<https://doi.org/10.1021/jm101336m>)
- Slate DL, Lee RH, Rodriguez J & Crews P 1994 The marine natural product, halistanol trisulfate, inhibits pp60v-src protein tyrosine kinase activity. *Biochemical and Biophysical Research Communications* **203** 260–264. (<https://doi.org/10.1006/bbrc.1994.2176>)
- Smyrniotopoulos V, Rae M, Soldatou S, Ding Y, Wolff CW, McCormack G, Coleman CM, Ferreira D & Tasdemir D 2015 Sulfated steroid-amino acid conjugates from the Irish marine sponge *Polymastia boletiformis*. *Marine Drugs* **13** 1632–1646. (<https://doi.org/10.3390/md13041632>)
- Sousa M, da Silva MC & Pinto M 2008 Sulfated flavonoids: nature playing with the hydrophilic-hydrophobic balance. In *Natural Products: Chemistry, Biochemistry and Pharmacology*, edn 1, pp 392–416. Ed G Brahmachari. New Delhi, India: Narosa Publishing House.
- Spellberg B, Blaser M, Guidos RJ, Boucher HW, Bradley JS, Eisenstein BI, Gerding D, Lynfield R, Reller LB, Rex J, et al. 2011 Combating antimicrobial resistance: policy recommendations to save lives. *Clinical Infectious Diseases* **52** (Supplement 5) 397–428. (<https://doi.org/10.1093/cid/cir153>)
- Stacey N 2015 Hormonally derived pheromones in teleost fishes. In *Fish Pheromones and Related Cues*, pp 33–88. Eds PW Sorensen & BD Wisenden. Iowa, USA and Oxford, UK: John Wiley & Sons, Inc.
- Stonik VA 2001 Marine polar steroids. *Russian Chemical Reviews* **70** 673–715. (<https://doi.org/10.1070/RC2001v070n08ABEH000679>)
- Sun HH, Gross SS, Gunasekera M & Koehn FE 1991 Weinbersterol disulfates A and B, antiviral steroid sulfates from the sponge *Petrosia weinbergi*. *Tetrahedron* **47** 1185–1190. ([https://doi.org/10.1016/S0040-4020\(01\)86375-4](https://doi.org/10.1016/S0040-4020(01)86375-4))
- Tang HF, Yi YH, Li L, Sun P, Zhang SQ & Zhao YP 2005a Three new asterosaponins from the starfish *Culcita novaeguineae* and their bioactivity. *Planta Medica* **71** 458–463. (<https://doi.org/10.1055/s-2005-871215>)
- Tang HF, Yi YH, Li L, Sun P, Zhang SQ & Zhao YP 2005b Bioactive asterosaponins from the starfish *Culcita novaeguineae*. *Journal of Natural Products* **68** 337–341. (<https://doi.org/10.1021/np0401617>)

- Tang HF, Yi YH, Li L, Sun P, Zhang SQ & Zhao YP 2006 Asterosaponins from the starfish *Culcita novaeguineae* and their bioactivities. *Fitoterapia* **77** 28–34. (<https://doi.org/10.1016/j.fitote.2005.07.009>)
- Tang HF, Cheng G, Wu J, Chen XL, Zhang SY, Wen AD & Lin HW 2009 Cytotoxic asterosaponins capable of promoting polymerization of tubulin from the starfish *Culcita novaeguineae*. *Journal of Natural Products* **72** 284–289. (<https://doi.org/10.1021/np8004858>)
- Thao NP, Cuong NX, Luyen BTT, Thanh NV, Nhiem NX, Koh Y-S, Ly BM, Nam NH, Kiem PV, Minh CV, et al. 2013 Anti-inflammatory asterosaponins from the starfish *Astropecten monacanthus*. *Journal of Natural Products* **76** 1764–1770. (<https://doi.org/10.1021/np400492a>)
- Thao NP, Luyen BT, Kim EJ, Kang HK, Kim S, Cuong NX, Nam NH, Kiem PV, Minh CV & Kim YH 2014 Asterosaponins from the starfish *Astropecten monacanthus* suppress growth and induce apoptosis in HL-60, PC-3, and SNU-C5 human cancer cell lines. *Biological and Pharmaceutical Bulletin* **37** 315–321. (<https://doi.org/10.1248/bpb.113-00705>)
- Tsakamoto S, Kato H, Hirota H & Fusetani N 1997 Antifouling terpenes and steroids against barnacle larvae from marine sponges. *Biofouling* **11** 283–291. (<https://doi.org/10.1080/08927019709378337>)
- Tsakamoto S, Matsunaga S, Fusetani N & van Soest RW 1998 Acanthosterol sulfates A–J: ten new antifungal steroidal sulfates from a marine sponge *Acanthodendrilla* sp. *Journal of Natural Products* **61** 1374–1378. (<https://doi.org/10.1021/np980178n>)
- Ushiyama S, Umaoka H, Kato H, Suwa Y, Morioka H, Rotinsulu H, Losung F, Mangindaan RE, de Voogd NJ, Yokosawa H, et al. 2012 Manadosterols A and B, sulfonated sterol dimers inhibiting the Ubc13-Uev1A interaction, isolated from the marine sponge *Lissodendryx fibrosa*. *Journal of Natural Products* **75** 1495–1499. (<https://doi.org/10.1021/np300352u>)
- Vilas-Boas C, Sousa E, Pinto M & Correia-da-Silva M 2017 An antifouling model from the sea: a review of 25 years of zosteric acid studies. *Biofouling* **33** 927–942. (<https://doi.org/10.1080/08927014.08922017.01391951>)
- Whitson EL, Bugni TS, Chockalingam PS, Concepcion GP, Harper MK, He M, Hooper JNA, Mangalindan GC, Ritacco F & Ireland CM 2008 Sphecosterol sulfates, PKC ζ inhibitors from a Philippine sponge *Spheciospongia* sp. *Journal of Natural Products* **71** 1213–1217. (<https://doi.org/10.1021/np8001628>)
- Whitson EL, Bugni TS, Chockalingam PS, Concepcion GP, Feng X, Jin G, Harper MK, Mangalindan GC, McDonald LA & Ireland CM 2009 Fibrosterol sulfates from the Philippine sponge *Lissodendoryx* (*Acanthodoryx*) *fibrosa*: sterol dimers that inhibit PKC ζ . *Journal of Organic Chemistry* **74** 5902–5908. (<https://doi.org/10.1021/jo900844r>)
- Yang SW, Chan TM, Pomponi SA, Chen G, Loeberberg D, Wright A, Patel M, Gullo V, Pramanik B & Chu M 2003a Structure elucidation of a new antifungal sterol sulfate, Sch 575867, from a deep-water marine sponge (Family: *Astroscleridae*). *Journal of Antibiotics* **56** 186–189. (<https://doi.org/10.7164/antibiotics.56.186>)
- Yang SW, Buivich A, Chan TM, Smith M, Lachowicz J, Pomponi SA, Wright AE, Mierzwa R, Patel M, Gullo V, et al. 2003b A new sterol sulfate, Sch 572423, from a marine sponge, *Topsentia* sp. *Bioorganic and Medicinal Chemistry Letters* **13** 1791–1794. ([https://doi.org/10.1016/S0960-894X\(03\)00260-9](https://doi.org/10.1016/S0960-894X(03)00260-9))
- Zhao Y, Zhu C, Li X, Zhang Z, Yuan Y, Ni Y, Liu T, Deng S, Zhao J & Wang Y 2011 Asterosaponin 1 induces endoplasmic reticulum stress-associated apoptosis in A549 human lung cancer cells. *Oncology Reports* **26** 919–924. (<https://doi.org/10.3892/or.2011.1358>)
- Zhou J, Cheng G, Cheng G, Tang HF & Zhang X 2011 Novaeguinoside II inhibits cell proliferation and induces apoptosis of human brain glioblastoma U87MG cells through the mitochondrial pathway. *Brain Research* **1372** 22–28. (<https://doi.org/10.1016/j.brainres.2010.11.093>)
- Zvyagintseva TN, Makar'eva TN, Stonik VA & Elyakova LA 1986 Sulfated steroids of sponges of the family *Halichondriidae* – Natural inhibitors of endo-(1 $\rightarrow$ 3)- β -D-glucanases. *Chemistry of Natural Compounds* **22** 66–71. (<https://doi.org/10.1007/BF00574583>)

Received in final form 21 December 2017

Accepted 3 January 2018